

8<sup>th</sup> World Congress on

## PLANT GENOMICS AND PLANT SCIENCE

August 10-11, 2018 Osaka, Japan

**Mutated and wild type *Gossypium* Universal Stress Protein-2 (*GUSP-2*) gene confers resistance to stresses in *Escherichia coli*, *Pichia pastoris* and cotton plant****Nadeem Hafeez**

Center for Excellence in Molecular Biology, Pakistan

*Gossypium arboreum* is considered to be a rich source of stress responsive genes and EST data base revealed that most of its genes are uncharacterized. The full length *Gossypium* Universal Stress Protein-2 (*GUSP-2*) gene (510 bp) was cloned in *Escherichia coli*, *Pichia pastoris* and *Gossypium hirsutum*, characterized and point mutated at three positions separately at 352-354, lysine-60 to proline (*M1-usp-2*) and 214-216, aspartic acid-26 to serine (*M2-usp-2*) and 145-147, lysine-3 to proline (*M3-usp-2*) to study its role in abiotic stress tolerance. It was found that heterologous expression of one mutant (*M1-usp-2*) provided enhanced tolerance against salt and osmotic stresses, recombinant cells have higher growth up to 10<sup>-5</sup> dilution in spot assay as compared to *W-usp-2* (wild type *GUSP-2*), *M2-usp-2* and *M3-usp-2* genes. *M1-usp-2* in *Pichia pastoris* transcript profiling exhibited significant expression (7.1-fold) to salt and (9.7) and osmotic stresses. *M1-usp-2* gene was also found to enhance drought tolerance and significant expression (8.7) in CIM-496-*Gossypium hirsutum* transgenic plants. However, little tolerance against heat and cold stresses both in recombinant yeast and bacterial cells was observed. The results from our study concluded that activity of *GUSP-2* was enhanced in *M1-usp-2* but wipe out in *M2-usp-2* and *M3-usp-2* response remained almost parallel to *W-usp-2*. Further, it was predicted through in silico analysis that *M1-usp-2*, *W-usp-2* and *M3-usp-2* may be directly involved in stress tolerance or function as signaling molecule to activate the stress adaptive mechanism.

chnadeemhfz@gmail.com