

24th Global Dentists and Pediatric Dentistry Annual Meeting

June 11-12, 2018 | London, UK

Study of effect of Polyvinylpyrrolidone-iodine (PVP-I) 2% as an anti-oedematous agent in third molar surgery

Kanwaldeep Singh Soodan
M M University, India

A single blind randomized control trial was carried out on 50 healthy outpatients who required surgical removal of Mandibular third molars under local anesthesia were selected. The patients were divided into 2 groups (n=25), the treatment (PVP-I) and control group (normal saline). The treatment group patients were irrigated using PVP-I 2% (w/v) (Betadine, Win- Medicare, India) during bone guttering and tooth sectioning. The control group patients were irrigated with saline (sodium chloride 0.9%, w/v; Parental Drugs, India) only. Procedures that exceeded more than 1 hour were excluded from the study. Using Pederson difficulty index, patients with moderately difficulty index were chosen. All parameters for swelling were recorded preoperatively, on the first, second and seventh postoperative days for both procedures. The data were statistically analyzed using SPSS (version 22.0) software. Independent t-test was applied for Operative time in minutes and the two groups matched ($p > 0.05$) for operative time. For change in swelling, T-tests was applied and we found increase in swelling in saline group which was highly significant for change from preoperative to day 2 ($p = .005$) and from preoperative to day 7 ($p\text{-value} < .001$). Mean for Pederson index for Betadine and saline group was found out to be same ($P = 1$). PVP-I 2% was found out to be significantly reducing swelling as compared to saline suggesting that it acts as an anti-oedematous agent in mandibular third molar surgery.

dr.kanwal@rediffmail.com