

18th World Congress on Clinical Nursing & Practice

September 21-22, 2018 | Prague, Czech Republic

Self-care agency and associated factors in patients with systemic lupus erythematosus

Hui Yang

Sichuan University, China

Aim: The aim of this study was to estimate the level of self-care agency and explore its associated factors in patients with systemic lupus erythematosus (SLE).

Method: A cross-sectional study was conducted between July and October 2016 and all patients were from a tertiary general hospital in Southwest China. The self-care agency was assessed by the exercise of self-care agency scale (ESCA). Pain and fatigue were assessed by the visual analog scale (VAS). The disease activity was measured using Systemic lupus erythematosus disease activity index 2000 (SLEDAI-2K). Physical and mental health were evaluated by the physical (PCS) and mental component summary (MCS) of the instrument SF-36, version 2. Multivariate regression analysis was performed to explore the associated factors of self-care agency.

Result: A total of 123 patients were recruited. The mean of ESCA was 86.29. In univariate analysis, self-care agency of patients has the difference in gender, work status, educational level, household income and disease activity ($p<0.05$). Additionally, higher BMI, higher level of fatigue and worse mental health was found in patients with lower self-care agency ($p<0.05$). The stepwise multivariate regression analysis showed that male ($p=0.001$), lower educational level ($p=0.003$), lower household income ($p<0.001$) and worse mental health ($p<0.001$) could predict lower self-care agency.

Conclusion: These results indicated that the patients with SLE had a middle level of self-care agency. The lower level of self-care agency was assessed with male, lower educational level, lower household income and worse mental health. These findings pointed out the importance of targeted intervention to enhance the self-care agency in patients with SLE.

13281005120@163.com

Notes: