OPEN ACCESS Freely available online

ISSN: 2157-7560
Journal of
Vaccines & Vaccination

Opinion Article

Advanced Antigen Delivery Strategies for Improved Immune Activation

Mirela Kovacen*

Department of Vaccine Science, University of Ljubljana, Ljubljana, Slovenia

DESCRIPTION

Antigen delivery systems are specialized technologies developed to transport immune-stimulating substances into the body in a controlled and effective manner. Their main purpose is to improve the exposure of antigens to immune cells and support the development of a specific immune response. An antigen may consist of a protein, peptide, carbohydrate, nucleic acid product, or another biological structure that can be recognized by the immune system. Although some antigens can stimulate immunity when administered directly, their biological activity may be reduced by rapid degradation, poor tissue distribution, or limited uptake by antigen-presenting cells. Delivery systems are therefore being studied as a means of protecting antigens and improving their interaction with the cells responsible for immune recognition.

The immune system relies on specialized cells, including dendritic cells, macrophages, and B lymphocytes, to recognize foreign biological structures. Dendritic cells are particularly important because they can capture antigens, process them, and present selected fragments to T lymphocytes. This process initiates a sequence of cellular interactions that can lead to antibody production and cellular immune activity. The effectiveness of an antigen delivery system depends partly on its ability to place the antigen in a biological environment where these processes can occur efficiently.

Lipid-based carriers represent one of the major categories being examined for antigen transport. Liposomes, for example, consist of lipid membranes that can surround and protect biological materials. Their composition can be modified to influence stability, cellular interaction, and distribution after administration. Lipid nanoparticles provide another type of carrier and can be prepared with different lipid components depending on the material being transported. Their physical properties may influence how they enter cells and how their contents become available for immune processing.

Polymeric carriers are also being investigated for controlled antigen release. These systems can be produced from biodegradable materials and formed into particles, capsules, or other structures. As the carrier gradually breaks down, the antigen can be released over a defined period. This approach may help maintain antigen exposure for longer periods than some free-antigen formulations. Polymer selection can influence degradation rate, particle size, antigen loading, and biological compatibility, making formulation design an important part of development.

Protein-based particles provide another approach to antigen administration. Certain proteins can naturally assemble into organized structures that allow antigens to be displayed on their surfaces. Repeated presentation of an antigen can influence interactions with immune receptors and may support strong antibody responses. These structures can be designed without using complete infectious organisms, making them useful for investigating vaccine approaches that require controlled antigen presentation.

Nanoparticle-based systems have received considerable research attention because their size, surface characteristics, and composition can be adjusted. Particles within suitable size ranges may be efficiently taken up by dendritic cells and macrophages. Surface modifications can also affect interactions with biological membranes and proteins. Some delivery systems are designed to respond to environmental conditions, such as changes in acidity or enzyme activity, allowing researchers to examine controlled release at specific cellular locations.

The administration route is another factor that influences antigen delivery. Intramuscular injection is widely used for many vaccines, while subcutaneous and intradermal administration can provide different patterns of tissue exposure. Intradermal delivery is of interest because the skin contains several types of immune cells. Nasal and oral delivery systems are being investigated for mucosal vaccination, with the goal of stimulating immune responses at surfaces that frequently encounter microorganisms. Oral delivery, however, requires protection against digestive enzymes and acidic conditions that can damage sensitive antigens.

Correspondence to: Mirela Kovacen, Department of Vaccine Science, University of Ljubljana, Ljubljana, Slovenia, E-mail: mirela.kovacen@uvl-research.si

Received: 20-Apr-2026, Manuscript No. JVV-26-32839; **Editor assigned:** 22-Apr-2026, Pre QC No. JVV-26-32839 (PQ); **Reviewed:** 06-May-2026, QC No. JVV-26-32839; **Revised:** 13-May-2026, Manuscript No. JVV-26-32839 (R); **Published:** 20-May-2026, DOI: 10.35248/2157-7560.26.17.662.

Citation: Kovacen M (2026). Advanced Antigen Delivery Strategies for Improved Immune Activation. J Vaccines Vaccin. 17:662.

Copyright: © 2026 Kovacen M. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Antigen delivery systems also have applications in therapeutic vaccination research. In cancer studies, researchers may use tumor-associated antigens to stimulate immune recognition of abnormal cells. A delivery carrier can assist in transporting these antigens to antigen-presenting cells and supporting their processing. Similar approaches are being examined for infectious diseases, where selected components of pathogens can be delivered without administering the complete infectious organism.

CONCLUSION

Research in antigen delivery continues to combine findings from immunology, pharmaceutical science, materials science, and molecular biology. Scientists are evaluating systems that can improve antigen protection, cellular uptake, controlled release, and immune presentation while maintaining acceptable safety profiles. Future development will depend on producing delivery platforms that are effective, reproducible, stable, affordable, and suitable for practical manufacturing. As understanding of immune-cell behavior and biomaterial interactions increases, antigen delivery technologies may contribute to improved preventive vaccines and therapeutic immune interventions across a range of diseases.