



Advanced Glycation End Products: Biochemical Formation and Impact on Human Health

Olivia Turner*

Department of Biochemistry, Redwood University, Vancouver, Canada

DESCRIPTION

Advanced Glycation End products, commonly referred to as AGEs, are a diverse group of molecules formed through non-enzymatic reactions between reducing sugars and proteins, lipids, or nucleic acids. This process, known as glycation, occurs naturally in the body and is accelerated under conditions of elevated glucose levels. Over time, the accumulation of these compounds contributes to cellular damage and plays a significant role in the development of various chronic diseases, particularly those associated with aging and metabolic imbalance. The formation of AGEs begins with an initial reaction between a sugar molecule and a free amino group on a protein or lipid, resulting in the formation of a Schiff base. This intermediate undergoes further rearrangement to form more stable amadori products. Through a series of complex chemical transformations, these early glycation products eventually become irreversible advanced glycation end products. The rate of AGE formation is influenced by factors such as glucose concentration, oxidative stress, and the presence of reactive intermediates.

AGEs can accumulate both endogenously and exogenously. Within the body, their formation is closely linked to metabolic conditions such as diabetes, where chronic hyperglycemia accelerates glycation reactions. Exogenous sources of AGEs include dietary intake, particularly foods that are processed or cooked at high temperatures, such as grilling, frying, or roasting. These cooking methods promote chemical reactions that generate significant amounts of AGEs, which can be absorbed into the bloodstream upon consumption.

Once formed, AGEs can alter the structure and function of biological molecules. Proteins modified by glycation often lose their normal function and become more resistant to degradation. This can lead to the accumulation of damaged proteins within tissues, impairing cellular processes. In structural proteins such as collagen, glycation increases stiffness and

reduces elasticity, contributing to tissue damage and aging-related changes.

A key mechanism through which AGEs exert their effects is by interacting with specific cell surface receptors, most notably the receptor for advanced glycation end products, commonly abbreviated as RAGE. Binding of AGEs to RAGE activates intracellular signaling pathways that promote inflammation and oxidative stress. These responses can damage cells and tissues, contributing to the progression of chronic diseases such as cardiovascular disease, kidney disease, and neurodegenerative disorders.

In diabetes, AGEs play a central role in the development of complications affecting multiple organ systems. Elevated blood glucose levels accelerate AGE formation, leading to damage in blood vessels, nerves, and organs such as the kidneys and eyes. In the vascular system, AGEs contribute to endothelial dysfunction, reducing the ability of blood vessels to regulate blood flow. This increases the risk of atherosclerosis and cardiovascular events. In the kidneys, AGE accumulation can impair filtration function, leading to diabetic nephropathy. Similarly, in the eyes, AGEs are involved in the development of diabetic retinopathy by damaging retinal blood vessels.

The body has natural defense mechanisms to limit AGE accumulation, including enzymatic systems that degrade reactive intermediates and remove damaged molecules. However, these systems can become overwhelmed under conditions of chronic metabolic stress, leading to increased AGE levels. Aging is also associated with a gradual decline in the efficiency of these protective mechanisms, resulting in higher AGE accumulation over time.

CONCLUSION

Advanced glycation end products are bioactive molecules formed through complex chemical reactions that have significant effects on cellular function and overall health. Their accumulation is

Correspondence to: Olivia Turner, Department of Biochemistry, Redwood University, Vancouver, Canada, E-mail: olivia.turner.research@redwooduniv.ca

Received: 24-Feb-2026, Manuscript No. DCRS-26-31318; **Editor assigned:** 26-Feb-2026, Pre QC No. DCRS-26-31318 (PQ); **Reviewed:** 12-Mar-2025, QC No. DCRS-26-31318; **Revised:** 19-Mar-2025, Manuscript No. DCRS-26-31318 (R); **Published:** 26-Mar-2025, DOI: 10.35841/2572-5629.26.11.268

Citation: Turner O (2026). Advanced Glycation End Products: Biochemical Formation and Impact on Human Health. Diabetes Case Rep. 11:268.

Copyright: © 2026 Turner O. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

influenced by metabolic conditions, dietary habits, and aging processes. The significance of AGEs extends beyond metabolic diseases, as they are also implicated in aging and various degenerative conditions. Their ability to modify proteins, induce inflammation, and generate oxidative stress makes them important contributors to tissue damage across multiple systems.

Understanding the role of AGEs provides valuable insight into the mechanisms underlying chronic diseases and highlights the importance of preventive strategies. By addressing factors that contribute to AGE formation and exploring therapeutic interventions, it is possible to reduce their impact and improve long-term health outcomes.