



Allelic Variation: Genetic Differences Shaping Biological Diversity and Drug Response

Naomi Carter*

Department of Human Genetics, University of Toronto, Toronto, Canada

DESCRIPTION

Allelic variation refers to differences in Deoxyribonucleic Acid (DNA) sequences that occur at the same position within corresponding chromosomes. Alternative forms of a gene are called alleles and differences among these alleles can influence physical traits, biochemical processes, disease susceptibility and responses to medicines. Such variation is a natural feature of human populations and contributes to differences among individuals. Some allelic changes have little or no observable effect, whereas others can modify the amount, structure, or activity of a protein.

A gene can contain several alleles within a population. These forms may differ by a single nucleotide or by larger sequence changes. Single-nucleotide variants involve the replacement of one DNA base with another. Insertions and deletions can add or remove one or more bases, while larger structural changes may alter sections of chromosomes. The biological consequence depends on where the alteration occurs and whether it affects gene expression or protein production. Allelic variation can occur within coding regions that provide instructions for protein synthesis. A nucleotide change may substitute one amino acid for another, potentially changing protein folding, stability, or activity. Some substitutions have minimal effects because the altered amino acid has similar chemical properties. Other changes can strongly influence protein behavior. Variants that introduce a premature stop signal may produce shortened proteins, while changes affecting essential functional regions can reduce or eliminate protein activity.

Variation can also occur outside coding sequences. Regulatory regions control when, where and how strongly genes are expressed. An allele in such a region may increase or decrease transcription without changing the protein sequence itself. Changes in promoters, enhancers, splice-related regions and untranslated segments can therefore influence the amount of a protein produced by cells. This type of variation is important because differences in protein quantity may affect physiological

processes and medication response. Allelic variation contributes to differences in drug metabolism. Genes encoding cytochrome P450 enzymes contain many variants that can change enzyme activity. *Cytochrome P450 2D6*, *CYP2C19* and *CYP2C9* are notable examples. Individuals carrying different alleles may process particular medicines at different rates. Reduced enzyme activity can increase exposure to some drugs, while increased activity can lower exposure or alter the conversion of certain prodrugs into active compounds.

Drug transport is also influenced by allelic differences. Transport proteins move medicines across cellular membranes and participate in absorption, distribution and elimination. Variants in transporter genes may change protein expression or function. As a result, the concentration of a medicine in blood or tissues can differ between individuals. The clinical importance depends on the medicine, the specific allele and other factors affecting drug disposition. Alleles can affect drug targets as well. Receptors, enzymes, ion channels and signaling proteins may contain sequence differences that alter their structure or activity. A medicine that acts on one of these targets may therefore produce different effects in individuals carrying different alleles. In oncology, genetic differences within tumor cells can influence whether a particular molecular target is present or active.

Immune-related allelic variation can influence susceptibility to selected adverse drug reactions. Certain human leukocyte antigen variants are associated with increased risk of severe immune responses to particular medicines. When clinical guidance recommends genetic testing before treatment, identification of a relevant allele can help clinicians select an alternative medicine and reduce avoidable risk. Modern DNA sequencing and genotyping methods allow scientists and clinicians to identify many types of allelic variation. Targeted genotyping can examine selected variants, while broader sequencing can detect changes across numerous genes. Bioinformatic tools compare DNA sequences, classify variants and connect genetic findings with available biological and

Correspondence to: Naomi Carter, Department of Human Genetics, University of Toronto, Toronto, Canada, E-mail: naomi.carter@geneticsmail.ca

Received: 29-Aug-2025, Manuscript No. JPP-25-32625; **Editor assigned:** 01-Sept-2025, PreQC No. JPP-25-32625 (PQ); **Reviewed:** 15-Sept-2025, QC No. JPP-25-32625; **Revised:** 22-Sept-2025, Manuscript No. JPP-25-32625 (R); **Published:** 29-Sept-2025, DOI: 10.35248/2153-0645.25.16.148

Citation: Carter N (2025) Allelic Variation: Genetic Differences Shaping Biological Diversity and Drug Response. J Pharmacogenom Pharmacoproteomics. 16:148.

Copyright: © 2025 Carter N. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

clinical information. Interpretation requires attention to the quality of the sequence data and the evidence supporting the functional effect of each variant.

Allelic variation is especially relevant to precision medicine because genetic differences can contribute to variation in treatment outcomes. Pharmacogenetic testing can identify selected alleles that influence drug metabolism, transport, target activity, or immune reactions. When validated clinical recommendations are available, such information can support medication selection or dose decisions. Genetic findings should still be considered together with kidney function, liver function, age, concurrent medicines, adherence and disease

characteristics. Allelic variation represents one of the major sources of genetic diversity within human populations. Differences in DNA can influence protein structure, gene expression, disease susceptibility, medication metabolism, immune responses and treatment outcomes. Understanding these variations helps connect genetic information with observable biological characteristics and clinical decisions. As genomic technologies become more accessible, systematic interpretation of alleles can contribute to improved diagnosis, safer medication use and more individualized approaches to healthcare.