

Antitumor Potential of Lactaptin

Vladimir A Richter*, Anna A Vaskova, Olga A Koval and Elena V Kuligina

Institute of Chemical Biology and Fundamental Medicine, Siberian Branch of the Russian Academy of Sciences, Lavrentiev av. 8, Novosibirsk, 630090, Russia

Abstract

A number of naturally occurring proteins that selectively induce apoptosis of cancer cells were recently used to develop novel anticancer drug substances. Human milk is a source of multiple bioactive peptides, some of which are activated by proteolysis. Lactaptin is an 8.6 kDa proteolytic fragment of kappa-casein isolated from human milk. This mini review describes apoptotic properties, mechanism of action and antitumor activity of lactaptin.

Keywords: Lactaptin; Antitumor; Proteins

Introduction

Human milk is known to contain proteins and peptides that kill tumor cells – HAMLET (Human α -lactalbumin made lethal to tumor cells) [1], lactoferrin, lactoferricin [2] and lactaptin. The potential applications of HAMLET, lactoferrin and lactoferricin as pharmaceutical drug candidates and clinical nutrition in the overall management of cancer have been examined in the review Chen et al. [3]. This mini review describes a little-known human milk protein lactaptin.

Origin of Lactaptin

We isolated and characterized the pro-apoptotic peptide from human milk, that was capable of reducing cell viability and inducing apoptosis in cultured tumor cells. We named this peptide lactaptin. Lactaptin was identified as 8.6 kDa proteolytic fragment of human kappa-casein, composed of 74 amino acid residues (residues 57–134 of kappa-casein) [4]. A series of recombinant analogues of lactaptin was constructed but only one of them, RL2, containing the complete amino acid sequence of lactaptin, effectively induced cell death in various human and mouse tumour cells *in vitro* while having no effect on the viability of nonmalignant MSC cells [5].

Mechanism of Lactaptin Apoptotic Action

We have analyzed biochemical markers of RL2 induced apoptosis of cancer cells. The activation of initiator and effector caspases as well as mitochondrial membrane potential and cytoplasm membrane changes were analyzed using flow cytometry and Western-blot methods. We have found that RL2 induced apoptotic death of tumor cells was accompanied by phosphatidylserine exposure on the plasma membrane surface. It was also shown that RL2 has induced dissipation of mitochondrial membrane potential and resulted in activation of initiator caspases 8, 9 and effector caspase 7 [6]. We demonstrated that RL2 downregulates Bcl-2 expression and induces p53-independent cell death. RL2 penetrates both into cancer and non-transformed cells. Identification of the cellular targets of the lactaptin analogue revealed that α / β -tubulin and α -actinin-1 were RL2-bound proteins [7].

Antitumor Activity of Lactaptin

Mice hepatocarcinoma HA-1 and human breast adenocarcinoma MDA-MB-231 were selected for testing RL2 antitumor activity *in vivo*. Female A/Sn mice between 10 and 12 weeks of age (colony of the Institute of Cytology and Genetics SB RAS, Novosibirsk, Russia) weighing 20-25g were used for the *in vivo* experiments. Ascites HA-1 tumor cells were washed and suspended in saline and 0.1 ml of the

suspension containing 2×10^5 cells was inoculated subcutaneously on the dorsal side or intramuscularly in the right thigh of the A/Sn mice. To obtain artificial metastases, 1.5×10^5 tumor cells in 0.2 ml saline were injected into the mice intravenously in the lateral tail vein. Repetitive injections of RL2 (5-50 mg/kg) for 3-5 days effectively inhibited ascites and solid tumor transplant growth when administered intravenously or intraperitoneally, without obvious side effects. The solid tumor inhibitory effect of RL2 (5 i.v. injections, cumulative dose 125 mg/kg) was comparable with that of cyclophosphamide at a therapeutic dose (5 i.v. injections, cumulative dose 150 mg/kg). In combination therapy with cyclophosphamide, RL2 had an additive antitumor effect for ascites-producing tumors. Histomorphometric analysis indicated a three-fold reduction of spontaneous metastases in the liver of RL2-treated mice with solid tumor transplants in comparison with control animals. Repeated RL2 treatment substantially prolonged the lifespan of mice with intravenously injected tumor cells [8,9].

To test *in vivo* RL2 antitumor activity to human cancer cells a mouse xenograft model of breast cancer was used. A suspension of MDA-MB-231 cells in Matrigel was subcutaneously injected into 6–8-week-old female SCID mice to form solid tumors. Tumors volumes were monitored every 5 days. When tumors reached roughly 100 mm³, RL2 (40 mg/kg) was injected intravenously every 2 days, with a total of three injections. The selected dose of RL2 and the treatment regime were based on a scheme of therapy described previously [8]. Seven days after the last injection the tumours were excised and weighed. RL2 injections significantly delayed tumors growth compared to the control group. The rate of inhibition of tumors growth after RL2 therapy was calculated as 43% [7].

The antitumor effect of RL2 was also examined against primary human endometrial cancer cells. Primary cell cultures of malignant and normal human endometrium were performed by enzymatic digestion of endometrial tissue from biopsy material. Our results indicate that the recombinant analog of lactaptin, RL2, exerts cytotoxic effects against

*Corresponding author: Vladimir A. Richter, Institute of Chemical Biology and Fundamental Medicine, Siberian Branch of the Russian Academy of Sciences, Lavrentiev av. 8, Novosibirsk, 630090, Russia, Tel: +7-383-363-51-52; Fax: +7-383-363-51-53; E-mail: richter@niboch.nsc.ru

Received August 27 2015; Accepted September 30 2015; Published October 07 2015

Citation: Richter VA, Vaskova AA, Koval OA, Kuligina EV (2015) Antitumor Potential of Lactaptin. Biol Med (Aligarh) S2: 002. doi: 10.4172/0974-8369.S2-004

Copyright: © 2015 Richter VA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

primary hormone-dependent endometrial tumor cells *in vitro* with features of apoptosis [8].

Now the pre-clinical trials of the drug Lactaptin, created on the basis of RL2, have been successfully completed. Like other therapeutic proteins Lactaptin is evenly distributed in organs and tissues, which results in reducing its concentration in the tumor and the antitumor efficiency decrease. Selective delivery of Lactaptin to the tumor site could alleviate this problem and reduce the therapeutic dose by increasing the local concentration of the drug within the tumor to achieve a desired antitumor response.

To increase the antitumor lactaptin efficacy we have chosen tumor specific peptides as the molecular targeting agents and constructed different plasmid vectors producing fusion proteins consisting of tumor specific peptides and RL2. To obtain tumor specific peptides we applied a selection method using phage peptide libraries (Ph.D.[™]- Phage Display Peptide Library Kit, New England Biolabs). Two displaying peptides GLHTSATNLYLH and SGVYKVAYDWQH were selected to mice hepatocarcinoma A-1 (HA-1) and Lung adenocarcinoma (LA). The RL2 and fusion proteins were expressed in *E.coli* and purified. We investigated relative cytotoxic effects *in vitro* and tumor retention of obtained proteins *in vivo*.

Breast cancer MDA-MB-231 and MCF-7 cells showed a similar decrease in viability after RL2 and fusion proteins treatment. It was shown, that fusion proteins with targeted peptides are accumulated in HA-1 tumor comparing to RL2.

Conclusion

Thereby we demonstrated that recombinant analog of human milk peptide lactaptin RL2 effectively induced apoptosis of tumor cells *in vitro*, suppressed the growth of sensitive tumors and metastases *in vivo*. Lactaptin is not toxic for nonmalignant normal cells and animals and

is considered to be the basis for the development of antitumor drugs.

Source of Support

The Ministry of Education and Science of the Russian Federation, Contract No. 14.607.21.0063 (project unique identifier RFMEFI60714X0063).

References

1. Svensson M, Håkansson A, Mossberg AK, Linse S, Svanborg C (2000) Conversion of alpha-lactalbumin to a protein inducing apoptosis. Proc Natl Acad Sci USA 97: 4221-4226.
2. Yin CM, Wong JH, Xia J, Ng TB (2013) Studies on anticancer activities of lactoferrin and lactoferricin. Curr Protein Pept Sci 14: 492-503.
3. Chen HY, Mollstedt O, Tsai MH, Kreider RB (2014). Potential clinical applications of multi-functional milk proteins and peptides in cancer management. Curr Med Chem 21: 2424-2437.
4. Nekipelaya VV, Semenov DV, Potapenko MO, Kuligina EV, Kit Yu, et al. (2008) Lactaptin is a human milk protein inducing apoptosis of MCF-7 adenocarcinoma cells. Dokl Biochem Biophys 419: 58-61.
5. Semenov DV, Fomin AS, Kuligina EV, Koval OA, Matveeva VA, et al. (2010) Recombinant analogs of a novel milk pro-apoptotic peptide, lactaptin, and their effect on cultured human cells. Protein J 29: 174-180.
6. Fomin AS, Koval' OA, Semenov DV, Potapenko MO, et al. (2012) The analysis of biochemical markers of MCF-7 cells apoptosis induced by recombinant analog of lactaptin. Bioorg Khim 38: 92-98.
7. Koval OA, Tkachenko AV, Fomin AS, Semenov DV, Nushtaeva AA, et al. (2014) Lactaptin induces p53-independent cell death associated with features of apoptosis and autophagy and delays growth of breast cancer cells in mouse xenografts. PLoS One 7: 1-12.
8. Koval OA, Fomin AS, Kaledin VI, Semenov DV, Potapenko MO, et al. (2012) A novel pro-apoptotic effector lactaptin inhibits tumor growth in mice models. Biochimie 94: 2467-2474.
9. Koval OA, Sakaeva GR, Fomin AS, Nushtaeva AA, Semenov DV, et al. (2015) Sensitivity of endometrial cancer cells from primary human tumor samples to new potential anticancer peptide lactaptin. J Cancer Res Ther 11: 345-351.

This article was originally published in a special issue, **Cancer Biology and Therapeutics** handled by Editor. Tapasya Srivastava, University of Delhi South Campus, India