

Application of Harm Reduction Interventions to Recreational Drug Use

Fernández-Calderón Fermín*

Department of Clinical, Experimental and Social Psychology, University of Huelva, Spain

Harm reduction programs are strategies aimed to minimize harm associated with a given circumstance. One of these circumstances is drug consumption, where intervention under this approach is designed to minimize the potential risks associated with drug use rather than eliminate it [1,2]. Harm reduction aims to empower drug users to make informed and responsible decisions that help them to reduce the risk associated with their drug use.

Harm reduction arose during the 80s in the Netherlands and United Kingdom to reduce the negative consequences of drug injection [3]. Since then, it has been expanded worldwide and nowadays it represents a normalized and accepted intervention with drug users who have established a problematic relationship with the substance (mainly with opiates). Its activities include: methadone maintenance treatment, needle exchange programs or drug consumption rooms [4]. Moreover, in the 90s harm reduction started to be applied in recreational contexts and nightlife. In these contexts, risks are more linked to the acute effects of drugs (overdose) than substance use disorders and related problems. Among its strategies, it can be found: outreach activities, peer-education interventions or on-site drug checking [5]. Like many authors have shown [6,7] these interventions are well accepted and appreciated by recreational drug users.

Nevertheless, harm reduction interventions in recreational settings are less common than those aimed to problematic drug user, and are not accepted and normalized as an option for dealing with possible drug problems. Keeping in mind that most drug users do not develop Substance Use Disorders [8], this seems to be contradictory. That is, paradoxically, most drug users (recreational) are less likely to receive interventions aimed to mitigate the potential harms they expose to.

This difference in the application and acceptance of harm reduction strategies is rooted in the abstentionist/prohibitionist discourse on drugs, which is ideologically and culturally dominant [9]. The idea that emerges from this discourse is "drug user" as an addict or a person with problems, the idea that the use of drugs necessarily implies having problems, or developing them in the future (mechanic vision of drug use). From this point of view, harm reduction is accepted as an intervention approach with marginal drug users who have developed a problematic relationship with drugs. However, when it comes to

normalized and socially integrated users, the most common strategy is to separate them from consumption to prevent them from developing Substance Use Disorders.

In conclusion, harm reduction strategies are recommended for those recreational drug users that do not wish to give up their substance use. This approach is presented as a public health option grounded more in pragmatism and cost-benefit analysis than on moral issues. In this sense, harm reduction still has a long way to go, making its discourse and strategies visible [9]. Only then, could such strategies be placed as a normalized intervention within the different options for drug users' assistance.

References

1. Marlatt GA (1998) Basic principles and strategies of harm reduction. In Marlatt GA (ed) Harm Reduction: Pragmatic strategies for managing high-risk behaviors. New York: Guilford Press.
2. Marlatt GA, Witkiewitz K (2010) Update on harm-reduction policy and intervention research. *Annu Rev Clin Psychol* 6: 591-606.
3. Cook C, Bridge J, Stimson G V (2010) The diffusion of harm reduction in Europe and beyond. In Rhodes Ty, Hendrich D (eds) Harm reduction: evidence, impacts and challenges. European Monitoring Centre for Drugs and Drug Addiction 37-56.
4. European Monitoring Centre for Drugs and Drug Addiction (2010) Harm reduction: evidence, impacts and challenges. Publications Office of the European Union, Luxembourg.
5. Charlois T (2009) Safer Nightlife in Europe. 5th Meeting EXASS Network. Council of Europe's Pompidou Group.
6. Branigan P, Wellings K (1999) Acceptance of the Harm Minimization Message in London Clubs and Underground System. *Drugs: Education, Prevention and Policy* 6: 389-398.
7. Chinet L, Stéphan P, Zobel F, Halfon O (2007) Party drug use in techno nights: A field survey among French-speaking Swiss attendees. *Pharmacology, Biochemistry and Behavior* 86: 284-289.
8. European Monitoring Centre for Drugs and Drug Addiction (2013) European Drug Report 2013. Trends and Developments. Publications Office of the European Union, Luxembourg.
9. Faura R, García N (2013) El ocio nocturno y la reducción de riesgos. In: Martínez DP, Pallarés J (Eds) De riesgos y placeres. Manual para entender las drogas. Lleida: Milenio 169-179.

***Corresponding author:** Fernández-Calderón Fermín, Department of Clinical, Experimental and Social Psychology, University of Huelva, Spain, Tel: +0034959218438; Fax: +34954712619; E-mail: fermin.fernandez@dpes.uhu.es

Received February 24, 2014; **Accepted** February 26, 2014; **Published** February 28, 2014

Citation: Fernández-Calderón F (2014) Application of Harm Reduction Interventions to Recreational Drug Use. *J Socialomics* 3: e123. doi:10.4172/2167-0358.1000e123

Copyright: © 2014 Fernández-Calderón F. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.