

COVID-19 and Dermatology-A Short Note

Bhavna Joshi

Director, Atharva Multispeciality Ayurveda Hospital, India

Correspondence to: Bhavna Joshi, Director, Atharva Multispeciality Ayurveda Hospital, India, Email: drgaurang_joshi@yahoo.com

Received: July 07, 2020; **Accepted:** July 23, 2020; **Published:** July 29, 2020

Citation: Bhavna Joshi (2020) COVID-19 and dermatology-A Short Note. J Dermatit 8:121. doi: 10.35248/2684-1436.20.5.121.

Copyright: © 2020 Joshi B, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

The World Health Organization (WHO) proclaimed Coronavirus disease-2019 (COVID-19) as a pandemic illness brought about by SARS-CoV-2 (recently known as 2019-nCoV). [1]. By the nineteenth of April, 2 241 778 affirmed patients were accounted for worldwide with 152 552 COVID-19-related passings. The COVID-19 pandemic speaks to the most genuine wellbeing emergency confronting the cutting-edge world bringing about extraordinary endeavors to contain this pandemic and its results [1].

Virology

All the coronaviruses (CoVs) have a place with the 'Coronaviridae' family, which, comprises of a gathering of wrapped, positive-sense and single-stranded RNA infections with an all-encompassing field of unique roots. The CoVs are genotypically and serologically arranged into four genera: α , β , γ and δ -CoVs, of which, the α and β CoVs are equipped for tainting people. SARS-CoV-2 was perceived as an individual from β -CoVs with a genome grouping like the bat CoV and SARS-CoV-1 [2]. SARS-CoV-2 is portrayed by the nearness of explicit spike glycoprotein that shows solid restricting liking to the angiotensinogen-converting protein 2 (ACE2) receptors; in any case, it merits referencing that SARS-CoV-2 has 10-20 folds higher restricting proclivity to ACE2 receptors contrasted and SARS-CoV-1. For the most part, ACE2 might be communicated in various pieces of the human body including the lung, small digestive tract, colon, duodenum, kidney, testis, gallbladder, heart, throat and urinary bladder, delivering these organs a possible objective for the SARS-CoV-2. Strangely, ACE2 in the basal cell layer of the epidermis proceeding to the basal cell layer of the hair follicles, the smooth muscle cells that encase the sebaceous organs and the eccrine organs are available. There is a speculation that androgen receptors may assume a job in the seriousness of the COVID-19 patients. This theory depended on the lopsided pervasiveness of extreme COVID-19 among youngsters and grown-up females contrasted and grown-up guys. In addition, the ACE2 movement has appeared to diminish with the decrease of the androgen hormones. Be that as it may, this hypothesis isn't yet demonstrated.

Cutaneous appearances of COVID-19

A few viral contaminations are described by the nearness of related cutaneous appearances [3]. Cutaneous indication are appeared as erythematous rash, urticaria and chickenpox-like vesicles chiefly in the storage compartment with practically zero tingling. Skin sores were not connected with infection seriousness. COVID-19 patients may give urticarial ejection with no respiratory indications (hack or fever). On a similar hand, a COVID-19 quiet was misdiagnosed as dengue in Thailand as the vital introductions were skin rash with petechiae and low platelet check [4]. The cutaneous signs of COVID-19 are grouped into five significant clinical designs including pseudo-chilblain (19%), other vesicular ejections (9%), urticarial sores (19%), maculopapules (47%), and livedo or corruption (6%).

Effect on dermatology practice

By and large, skin is viewed as a piece of the resistant framework as it goes about as a shield against various natural upgrades. In this setting, trading off the trustworthiness of this obstruction by cuts, scratching, needling, pre-existing contaminations or illnesses of skin, wounds and consumes may encourage the intrusion by micro-organisms. Subsequently, dermatological patients might be at higher danger of SARS-CoV-2 either due the loss of the skin honesty by dermatological illnesses or the immunosuppressive treatment utilized in the administration of certain ailments.

Numerous patients may exploit the isolate period to do some restorative methodology thinking about that the isolate period is a fitting open door for recuperating and personal time of the skin; in any case, it ought to be noticed that fractionated ablative laser reemerging may bring about tight sections of dermal wounds taking into account possible contamination. Furthermore, Micro-needling particularly home ones may convey high danger of disease and viral autoinoculation.

Dermatologic malignancies

Malignant growth patients might be at a higher hazard for the improvement of irresistible sickness with a 3.5 folds expanded danger of creating COVID-19 unfavorable occasions including the requirement for mechanical ventilation as well as death because of their immunocompromised express that is related to the nature and the forcefulness of the neoplasm and the anti-cancer treatment. The patients follow with cutting edge basal cell carcinoma to hedgehog inhibitors during the COVID-19 pandemic. A report included 37 patients, of which, just three patients interfered with the treatment because of serious comorbidities, while all the rest of the patients either went to the outpatient center for the month to month remedy or utilized an altered treatment plan to drag out their treatment. It is vital to make reference to that there is no proof against the continuation of these medications during the current pandemic [5].

Then again, invulnerable checkpoint inhibitors (ICIs) are viewed as one of the promising treatment choices for patients with cutting edge melanoma. Vital, the helplessness of patients taking ICIs to irresistible infections has not been completely talked about in the writing. The utilization of ICIs during the current pandemic may speak to a matter of worry because of the possible cover between the SARS-CoV-2-related pneumonia and the aspiratory harmfulness of the ICIs (in any case, it ought to be referenced that this pneumological poisonousness is an uncommon occasion extending from 2.5% to 5% in the event of ICIs monotherapy to 7–10% if there

should be an occurrence of mix treatment). Besides, the synergism between the ICIs system of activity and the SARS-CoV-2 pathogenesis may potentiate the neurotic impact of COVID-19. In this setting, the utilization of ICIs ought to be painstakingly weighted by a case-by-case situation. Considering squamous cell carcinoma of the skin, similar safety measures (talked about previously) ought to be followed if immunomodulators are considered for treatment.

Word related skin issues among medicinal services laborers during COVID-19

A few skin difficulties may result from the drawn out utilization of individual defensive gear because of hyperhydration impact, grinding and unfavorably susceptible contact responses. The predominance of skin harm coming about because of wearing defensive gear was 97.0% among social insurance laborers including injuries of the nasal extension (83.1%), hands, cheek and temple. Moreover, exorbitant hand washing with cleansers and additionally disinfectants may cause contact dermatitis. In these settings, an accord of Chinese specialists on the assurance of the skin and mucous films was discharged suggesting that hand washing or gloves cleaning ought to be restricted to the accompanying minutes: before contacting the patients or any aseptic methodology and after presentation to body liquids, contacting the patient or any of the patients' things. Moreover, they suggested the utilization of ethanol for hand cleanliness and applying hand cream following each season of hand cleanliness if conceivable. Be that as it may, if there should be an occurrence of latex gloves it is prescribed to wear a cotton gloves underneath and to utilize lotions with skin glucocorticoid cream. So also, it is prescribed to utilize an appropriately fitting cover and goggles and apply lotions or gel at contact zones. The utilization of twofold assurance (benzalkonium chloride glue on the nasal scaffold followed by hydrocolloid dressing) before wearing the N95 veils is recommended.

Expected utilization of dermatological medications for the board of COVID-19

Numerous immunizations had been infused intralesionally to treat moles, as BCG, which, results in IL-12 increment and IL-4 diminishing and MMR antibody, which, is fit for invigorating the cell-mediated and humoral insusceptible reaction. Kids are portrayed by altogether lower dangers for creating serious type of COVID-19 with lower death rates contrasted with grown-up populace and this might be because of the worldwide dynamic viral vaccination of youngsters [6,7]. In light of these perceptions, the immunizations that have been utilized to treat viral skin sores can hypothetically be a potential treatment alternative of COVID-19; be that as it may, further examinations are required to affirm the job of antibodies in COVID 19 prophylaxis as well as treatment.

References

1. Puliatti S, Eissa A, Eissa R, Amato M, Mazzone E, et al. COVID-19 and urology: a comprehensive review of the literature. *BJU Int*. 2020;125(6):E7-E14.
2. Wu F, Zhao S, Yu B, Chen YM, Wang W, et al. A new coronavirus associated with human respiratory disease in China. *Nature*. 2020;579:265-269.
3. Memar O, Tying SK. Cutaneous viral infections. *J Am Acad Dermatol*. 1995;33(2): 279-287.
4. Joob B, Wiwanitkit V. COVID-19 can present with a rash and be mistaken for Dengue. *J Am Acad Dermatol*. 2020;82:e177.

5. Villani A, Fabbrocini G, Costa C, Scalvenzi M. Patients with advanced basal cell carcinomas in treatment with sonic hedgehog inhibitors during the coronavirus disease 2019 (COVID-19) period: management and adherence to treatment. *J Am Acad Dermatol*. 2020.
6. Thappa D, Chiramel M. Evolving role of immunotherapy in the treatment of refractory warts. *Indian Dermatol Online J*. 2016;7:364.
7. Dong Y, Mo X, Hu Y, Qi X, Jiang F, et al. Epidemiological characteristics of 2143 pediatric patients with 2019 coronavirus disease in china. *Pediatrics*. 2020; 16:e20200702.