OPEN ACCESS Freely available online

ISSN: 2572-5629
Diabetes Case Reports

Opinion Article

Curved Root Formation in Teeth: Developmental Deviation Known as Dilaceration

Yifang Su*

Laboratory of Oral Diseases, West China Hospital of Stomatology, Sichuan University, Chengdu, China

DESCRIPTION

Dilaceration is a developmental alteration in tooth structure characterized by an abrupt change in the normal alignment of a tooth's crown or root. This deviation results in a tooth that exhibits an unusual bend, which may occur in varying degrees of severity. The condition arises during the formative stages of odontogenesis when the developing dental tissues are still sensitive to external and internal disturbances. It is most commonly observed in permanent dentition, particularly involving maxillary incisors, though any tooth may be affected depending on the timing and nature of the disturbance.

The formation of teeth follows a carefully timed sequence in which enamel and dentin are deposited gradually while the tooth germ assumes its final orientation within the jaw. When a disturbance occurs during this period, especially involving displacement of a primary predecessor or direct impact to the jaw region in early childhood, the developing permanent tooth may change its orientation. As mineralization continues along this altered axis, the resulting structure preserves the curvature, leading to a permanent anatomical deviation.

One of the most recognized contributing factors is trauma to primary teeth. A forceful impact on a deciduous tooth can transmit energy to the underlying permanent tooth germ. Since the permanent successor is still in a formative state and not yet fully mineralized, it is vulnerable to positional alteration. This can result in an angular shift in either the crown or root depending on the stage of development at the time of injury. However, not all cases are associated with reported trauma, suggesting that other developmental disturbances or spatial constraints within the jaw may also contribute.

The morphological appearance of a tooth affected by this condition varies. In some cases, the crown may appear bent while the root remains relatively normal. In others, the crown is unaffected, but the root exhibits a pronounced curvature that becomes evident only after radiographic evaluation. The direction of the bend can differ as well, sometimes pointing labially, lingually, or mesially depending on the nature of the initial disturbance.

Detection of this condition is usually achieved through imaging methods. Standard radiographs often reveal an unexpected angulation that does not correspond to the typical anatomical pattern of the tooth. In more complex cases, advanced imaging techniques provide a clearer representation of the spatial orientation of the root system. Such evaluation is important when planning procedures involving affected teeth, as the irregular shape may complicate access and instrumentation.

Orthodontic considerations are also relevant when dealing with such teeth. Movement of a tooth with significant curvature may not follow predictable patterns due to altered root orientation within the bone. Forces applied during alignment procedures may distribute unevenly, which can affect movement efficiency and stability. In some situations, repositioning strategies may be adjusted, or alternative plans may be considered depending on the severity of the curvature and its effect on surrounding structures.

The prevalence of this condition is relatively low compared to other dental anomalies, yet it holds significant importance in diagnostic and procedural contexts. Studies indicate a higher occurrence in anterior teeth, particularly maxillary central incisors, likely due to their position and exposure to trauma during early developmental stages. Nonetheless, posterior teeth may also be affected, though less frequently.

Preventive considerations are mainly centered on reducing the likelihood of early childhood trauma. Protective supervision during activities that pose risk of falls or impact may indirectly reduce disturbances to developing permanent teeth. Early assessment following injury to primary teeth is important, as monitoring the underlying permanent tooth can help identify developmental changes at an early stage. Additionally, educating parents and caregivers about injury prevention and the importance of regular dental follow-ups can help minimize potential complications in permanent tooth development.

Correspondence to: Yifang Su, Laboratory of Oral Diseases, West China Hospital of Stomatology, Sichuan University, Chengdu, China, E-mail: s.yifang.dr.@ru.ch

Received: 27-May-2026, Manuscript No DCR-26-32204; **Editor assigned:** 29-May-2026, PreQC DCR-26-32204 (PQ); **Reviewed:** 12-Jun-2026, QC No DCR-26-32204; **Revised:** 19-Jun-2026, Manuscript No. DCR-26-32204 (R); **Published:** 26-Jun-2026, DOI: 10.35248/2161-1122.26.16.770

Citation: Su Y (2026). Curved Root Formation in Teeth: Developmental Deviation Known as Dilaceration. J Dentistry. 16:770.

Copyright: © 2026 Su Y. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

OPEN ACCESS Freely available online

CONCLUSION

Dilaceration represents a distinctive alteration in tooth morphology that can influence eruption patterns, internal anatomyandprocedural approaches in dentistry. Its recognition through imaging and careful evaluation plays an important role in ensuring appropriate handling of affected teeth within various aspects of dental care. From an educational standpoint, this condition highlights how sensitive tooth development is to environmental and positional influences. It also demonstrates the importance of careful interpretation of radiographic findings before undertaking any invasive procedures involving teeth with atypical morphology. Awareness of such variations assists in reducing procedural difficulties and supports more accurate planning in dental practice.