



# Epigenetic Pharmacology: Modulating Gene Activity Through Reversible Molecular Regulation

Sofia Nakamura\*

Department of Molecular Pharmacology, University of Tokyo, Tokyo, Japan

## DESCRIPTION

Pharmacological Epigenetics examines how medicines influence gene activity through molecular mechanisms that regulate Deoxyribonucleic Acid (DNA) and chromatin without changing the underlying DNA sequence. Epigenetic processes determine which genes are active or inactive in particular cells and tissues. These processes include DNA methylation, histone modification, chromatin remodeling and regulation by non-coding Ribonucleic Acid (RNA) molecules. Because epigenetic states can change in response to disease, environmental factors, cellular signals and medication exposure, they provide important targets for pharmacological intervention. DNA is organized within cells by association with proteins called histones. DNA and histones form chromatin, which can exist in relatively accessible or compact forms. The accessibility of chromatin influences whether transcription machinery can reach particular genes. Epigenetic mechanisms modify this organization and help determine patterns of gene expression. Pharmacological agents can alter these regulatory processes and consequently change the production of specific proteins.

DNA methylation is one of the best-characterized epigenetic mechanisms. It commonly involves the addition of a methyl group to cytosine residues at Cytosine-phosphate-Guanine (CpG) sites. DNA methyltransferase enzymes add these methyl groups, while other cellular processes can remove or modify them. Increased methylation in certain regulatory regions can reduce transcription, whereas reduced methylation may permit greater gene activity. Abnormal DNA methylation patterns occur in several diseases and can influence cellular behavior. DNA methyltransferases are therefore important pharmacological targets. Medicines that inhibit these enzymes can alter DNA methylation patterns and modify gene expression. Azacitidine and decitabine are examples of agents used in selected hematological conditions. Their activity involves interference with DNA methyltransferase function, producing changes in cellular gene regulation. Clinical use requires careful dosing and

monitoring because epigenetic processes participate in many normal cellular functions.

Histone modifications represent another major epigenetic mechanism. Histone proteins contain amino acid residues that can undergo chemical modifications such as acetylation, methylation, phosphorylation and ubiquitination. These modifications can influence chromatin structure and recruitment of regulatory proteins. Histone acetylation is often associated with increased transcriptional accessibility, while particular histone methylation patterns can either activate or repress gene expression depending on the residue involved. Histone acetyltransferases add acetyl groups to histones, whereas histone deacetylases remove them. Histone deacetylase inhibitors can increase histone acetylation and alter the expression of multiple genes. Several medicines in this class have been evaluated or approved for selected cancers and other conditions. Their effects can extend beyond histones because acetylation also occurs on non-histone proteins involved in cellular signaling and regulation.

Histone methylation is controlled by enzymes that add or remove methyl groups from specific histone residues. Histone methyltransferases and histone demethylases can therefore influence transcriptional states. Pharmacological inhibitors directed toward selected enzymes in these groups are being examined for diseases in which abnormal gene regulation contributes to cellular dysfunction. Chromatin remodeling complexes provide another regulatory layer. These protein complexes use energy from Adenosine Triphosphate (ATP) to reposition nucleosomes and alter access to DNA. Changes in chromatin remodeling can affect transcription, DNA repair, cell differentiation and other cellular activities. Mutations or altered activity within chromatin remodeling components have been observed in several cancers, making selected components potential pharmacological targets.

Non-coding RNAs also participate in epigenetic regulation. MicroRNAs can reduce the production of specific proteins by

**Correspondence to:** Sofia Nakamura, Department of Molecular Pharmacology, University of Tokyo, Tokyo, Japan, E-mail: sofia.nakamura@epigenpharm.com

**Received:** 29-Aug-2025, Manuscript No. JPP-25-32627; **Editor assigned:** 01-Sept-2025, PreQC No. JPP-25-32627 (PQ); **Reviewed:** 15-Sept-2025, QC No. JPP-25-32627; **Revised:** 22-Sept-2025, Manuscript No. JPP-25-32627 (R); **Published:** 29-Sept-2025, DOI: 10.35248/2153-0645.25.16.150

**Citation:** Nakamura S (2025) Epigenetic Pharmacology: Modulating Gene Activity Through Reversible Molecular Regulation. J Pharmacogenom Pharmacoproteomics. 16:150.

**Copyright:** © 2025 Nakamura S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

interacting with messenger RNA, while long non-coding RNAs can influence chromatin organization and transcription. Some medicines can affect the expression of non-coding RNAs indirectly through changes in signaling pathways. These RNA molecules are also being investigated as potential biomarkers of disease and treatment response. Cancer is a major area of epigenetic pharmacology. Cancer cells frequently contain abnormal patterns of DNA methylation, histone modification and chromatin organization. These changes can alter genes involved in cell division, differentiation, DNA repair and programmed cell death. Epigenetic medicines can modify these regulatory states and may restore the activity of genes that have become abnormally silenced.

Epigenetic mechanisms are also relevant to drug resistance. A cancer cell may change its gene expression without acquiring a

new DNA mutation, allowing it to survive exposure to a medicine. Alterations in DNA methylation, histone modifications, or chromatin accessibility can influence the expression of drug targets and survival pathways. Monitoring epigenetic changes can therefore provide information about changing treatment responses. Epigenetic pharmacology has potential applications beyond cancer. Abnormal gene regulation has been associated with neurological disorders, cardiovascular conditions, inflammatory diseases, metabolic disorders and immune dysfunction. However, because epigenetic mechanisms regulate many genes simultaneously, pharmacological intervention must be carefully evaluated to minimize unwanted effects on normal cellular processes.