OPEN ACCESS Freely available online

ISSN: 2155-9546
Journal of Aquaculture Research & Development

Short Communication

Erythrocyte Osmoregulation in Cultured Aquatic Animals

Charlotte Pessier*

Department of Fisheries, Auburn University, Auburn, AL, United States

DESCRIPTION

Erythrocytes are essential cellular components of the circulatory system in fish and other aquatic vertebrates. Their primary task is the transport of oxygen and carbon dioxide, but their physiological activities extend into several processes associated with acid base balance, ion exchange, cell volume regulation, and resistance to environmental stress [1]. In aquaculture systems, erythrocytes are continuously exposed to changes in water chemistry because fish may experience variation in salinity, temperature, dissolved oxygen, mineral concentration, and feeding conditions. Among these factors, osmotic variation has a direct influence on the movement of water across the erythrocyte membrane. The capacity of these cells to maintain an appropriate internal volume despite changes in the surrounding fluid is referred to as erythrocyte osmoregulation [2].

The internal environment of an erythrocyte contains dissolved salts, proteins, organic compounds, and other substances that contribute to osmotic pressure. When the concentration of dissolved particles outside the cell changes, water moves across the plasma membrane according to the difference in water potential. A sudden reduction in external osmotic concentration can cause water to enter the cell, producing cellular swelling. In contrast, exposure to a more concentrated external medium can result in water loss and cellular shrinkage. Excessive changes in volume can affect membrane stability, hemoglobin concentration, cellular shape, and the ability of erythrocytes to pass through narrow blood vessels. Fish therefore possess several cellular mechanisms that help maintain erythrocyte volume within a functional range [3].

The erythrocyte membrane contains lipid components and membrane proteins that regulate the movement of ions and water. Channels, transport proteins, and ion pumps contribute to the maintenance of intracellular electrolyte concentrations. Sodium and potassium gradients are particularly important because they influence water distribution and electrical properties across the membrane. The sodium potassium ATPase uses cellular energy to move sodium ions outward and potassium ions inward. This process contributes to the maintenance of intracellular ionic conditions and indirectly affects cell volume. Other transport pathways can modify chloride, bicarbonate, calcium, and hydrogen ion concentrations according to physiological requirements [4].

Erythrocyte volume regulation can occur through two general responses. Regulatory volume decrease takes place when cells swell because of water entry. In this situation, cells can lose intracellular ions and associated water, allowing their volume to move toward its previous condition. Regulatory volume increase occurs following cellular shrinkage and involves the uptake of ions and other osmotically active substances, followed by water entry. The effectiveness of these responses differs among fish species and can also vary according to developmental stage, acclimation history, and environmental conditions [5].

Aquaculture makes erythrocyte osmoregulation particularly relevant because cultured fish may be transferred between water sources with different salinity levels. Species that naturally inhabit estuaries or coastal environments often possess physiological mechanisms that permit movement between freshwater and seawater. Freshwater exposure generally creates a tendency for water to enter the body, whereas marine exposure creates a tendency for water to leave the body. Although whole-animal osmoregulation involves organs such as the gills, kidneys, and digestive tract, erythrocytes must independently preserve their cellular condition while circulating through these changing internal environments [6].

Changes in salinity can influence erythrocyte morphology as well as volume. Cells exposed to unsuitable osmotic conditions may become rounded, irregular, swollen, or contracted. Severe osmotic disturbance can damage the membrane and lead to hemolysis, releasing hemoglobin into the surrounding plasma. Hemolysis can reduce the number of functional circulating erythrocytes and interfere with oxygen transport. For cultured fish, excessive cellular damage may therefore contribute to reduced growth, impaired activity, lower stress tolerance, and poor survival when environmental conditions are not properly managed [7].

Correspondence to: Charlotte Pessier Department of Fisheries, Auburn University, Auburn, AL, United States, E-mail: charlotte.pessier@auburn.edu

Received: 03-Aug-2026, Manuscript No. JARD-26-32963; **Editor assigned:** 06-Aug-2026, PreQC No. JARD-26-32963 (PQ); **Reviewed:** 20-Aug-2026, QC No. JARD-26-32963; **Revised:** 27-Aug-2026, Manuscript No. JARD-26-32963 (R); **Published:** 03-Sep-2026, DOI: 10.35248/2155-9546.26.17.1136

Citation: Pessier C (2026). Erythrocyte Osmoregulation in Cultured Aquatic Animals. J Aquac Res Dev. 17.1136.

Copyright: © 2026 Pessier C. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Temperature can modify erythrocyte osmoregulatory responses because membrane fluidity and enzyme activity are temperature dependent. At lower temperatures, membrane lipid movement may decrease, potentially altering the activity of membrane transport systems. Higher temperatures can increase metabolic activity and modify the rates of ion movement and cellular reactions. Consequently, salinity and temperature should not always be considered independently when evaluating erythrocyte responses in aquaculture species. A salinity level tolerated under one temperature condition may produce a different cellular response under another [8].

Acid–base regulation is also closely associated with erythrocyte ion exchange. Fish erythrocytes participate in carbon dioxide transport and bicarbonate formation. Carbon dioxide generated by tissues enters the blood and is converted through reactions involving carbonic anhydrase. Bicarbonate and chloride exchange across the erythrocyte membrane helps maintain appropriate chemical balance while carbon dioxide is transported toward the respiratory surface. This activity demonstrates that ion transport across the erythrocyte membrane has consequences beyond simple volume control [9].

Oxygen availability may further influence erythrocyte physiology. Fish living in ponds, tanks, cages, and recirculating systems can experience temporary reductions in dissolved oxygen. During oxygen limitation, physiological adjustments may occur in circulation, respiration, and cellular metabolism. Erythrocytes may change their metabolic activity and ion transport behavior during such conditions. Since oxygen delivery depends on both hemoglobin function and adequate circulation, maintaining cellular volume and membrane integrity is important for effective gas transport [10].

CONCLUSION

Erythrocyte osmoregulation represents an important cellular component of aquatic animal physiology. Through controlled ion movement, membrane transport, volume adjustment, and metabolic regulation, erythrocytes maintain conditions that permit continued oxygen transport under changing environmental circumstances. Further investigation across cultured species can improve knowledge of species-specific responses and help aquaculture researchers establish suitable environmental conditions for healthy blood function and efficient production.

REFERENCES

1. Rodrigues S, Antunes SC, Nunes B, Correia AT. Histological alterations in gills and liver of rainbow trout (Oncorhynchus mykiss) after exposure to the antibiotic oxytetracycline. Environ Toxicol Pharmacol. 2017;53:164-176.
2. Wang J, He RZ, Lu GL, Luo HL, Lu DQ, Li AX. Vaccine-induced antibody level as the parameter of the influence of environmental salinity on vaccine efficacy in Nile tilapia. Fish Shellfish Immunol. 2018;82:522-30.
3. Burgos-Aceves MA, Lionetti L, Faggio C. Multidisciplinary haematology as prognostic device in environmental and xenobiotic stress-induced response in fish. Sci Total Envir. 2019;670:1170-1183.
4. Dawood MA, Moustafa EM, Gewaily MS, Abdo SE, AbdEl-Kader MF, SaadAllah MS, et al. Ameliorative effects of Lactobacillus plantarum L-137 on Nile tilapia (Oreochromis niloticus) exposed to deltamethrin toxicity in rearing water. Aquat Toxicol. 2020;219:105377.
5. Świacka K, Michnowska A, Maculewicz J, Caban M, Smolarz K. Toxic effects of NSAIDs in non-target species: a review from the perspective of the aquatic environment. Environ Pollut. 2021;273:115891.
6. Mugwanya M, Dawood MA, Kimera F, Sewilam H. Anthropogenic temperature fluctuations and their effect on aquaculture: A comprehensive review. Mar Pollut Bull. 2022;7(3):223-43.
7. Kumari A, Koner D, Lanong AS, Khongmawloh E, Snaitang R, Saha N. Changes in the expression of certain osmosensitive channel and transporter genes in primary hepatocytes of air-breathing catfish, Clarias magur: A strategy to adapt under osmotic stress. Aqua. 2023;562:738756.
8. Su C, Li J, Zhang M, Pan L, Wang Y, Ding Y, et al. Dietary cholesterol enhances osmoregulation, antioxidant defenses and immune response of Litopenaeus vannamei to alleviate the macromolecular damage induced by salinity stress. Aqua. 2023;563:738861.
9. Wambreuse N, Caulier G, Delroisse J, Eeckhaut I. Integration of Molecular and Cellular Immunity in Sea Cucumbers: A Comprehensive Review. Comp. Immunol. Rep. 2026:200315.
10. Li S, Li S, Yang Y, Wu Z. Dihydroartemisinin alleviates ammonia toxicity and autophagy in zebrafish (Danio rerio) gill. Aquac. Rep. 2024;34:101915.