OPEN ACCESS Freely available online

ISSN: 2572-5629
Diabetes Case Reports

Perspective

Etiological Considerations in the Development of Ameloblastic Fibro-Odontoma: Cellular and Developmental Perspectives

Lisa Danylak*

Department of Oral and Maxillofacial Pathology, Ridgeway University of Health Sciences, Melbourne, Australia

DESCRIPTION

Ameloblastic fibro-odontoma is an uncommon mixed odontogenic lesion that typically appears in younger individuals and contains both epithelial and mesenchymal components along with varying degrees of dental hard tissue formation. Understanding its causes requires careful consideration of odontogenic tissue development, signaling interactions during tooth formation, and localized disturbances that alter normal differentiation pathways. Although no single factor can be identified as solely responsible, multiple contributing biological influences have been described in scientific literature and clinical observations.

The lesion is believed to arise from remnants of the dental lamina or from pluripotent cells within the developing tooth germ. During normal odontogenesis, epithelial cells and ectomesenchymal cells coordinate closely through reciprocal signaling interactions. These interactions regulate stages such as bud formation, cap development, and bell stage differentiation. In cases where this coordination becomes altered, odontogenic cells may continue proliferating beyond typical developmental limits, resulting in a mixed lesion containing both soft tissue components and calcified dental structures. This abnormal continuation of developmental activity is considered a central mechanism in the formation of ameloblastic fibro-odontoma.

Another proposed factor involves developmental arrest or alteration during a specific stage of odontogenesis. If tooth development does not proceed beyond early formative stages, the retained tissue may retain embryonic characteristics. These cells may continue to proliferate slowly over time and form a mass containing various degrees of dental hard tissue differentiation. The presence of enamel-like and dentin-like structures within the lesion suggests that inductive interactions between epithelial and mesenchymal components remain partially active, even though they are no longer following normal developmental sequencing.

Hormonal and systemic influences have also been considered in discussions regarding etiology. Although direct causal relationships remain unconfirmed, some researchers suggest that growth-related hormonal activity during childhood and adolescence may influence the behavior of odontogenic remnants. Periods of rapid skeletal and dental development could potentially increase the responsiveness of residual odontogenic cells, contributing to abnormal growth patterns under certain conditions.

The role of cell signaling molecules such as growth factors and morphogens is another area of interest. These molecules regulate differentiation and proliferation during normal tooth formation. Disruptions in their expression or receptor activity may lead to continued activation of odontogenic pathways beyond their intended developmental timeframe. This may result in the formation of a lesion that retains the ability to generate dental tissues but lacks organized structural arrangement.

Immunological factors may indirectly influence lesion formation as well. Local immune responses within jaw tissues could modify cellular behavior through the release of cytokines and other signaling mediators. While not directly causative, such immune-mediated changes may alter the balance between cell proliferation and differentiation, creating conditions favorable for abnormal odontogenic growth.

It is also important to consider that ameloblastic fibro-odontoma may represent part of a spectrum of related odontogenic lesions rather than a completely distinct pathological entity. Some authors suggest that it may exist within a continuum that includes ameloblastic fibroma and odontoma, reflecting progressive stages of tissue differentiation within a single pathological process. In this context, its development may depend on timing and degree of maturation arrest or continuation during tooth formation.

Clinical observations support the idea that these lesions often develop during periods of active dental development, which further reinforces the association with odontogenic growth processes rather than external environmental carcinogenic influences. Most cases occur in younger age groups, aligning

Correspondence to: Lisa Danylak, Department of Oral and Maxillofacial Pathology, Ridgeway University of Health Sciences, Melbourne, Australia, E-mail: lisa.danylak@ridge.edu.au

Received: 27-May-2026, Manuscript No DCR-26-32200; **Editor assigned:** 29-May-2026, PreQC DCR-26-32200 (PQ); **Reviewed:** 12-Jun-2026, QC No DCR-26-32200; **Revised:** 19-Jun-2026, Manuscript No. DCR-26-32200 (R); **Published:** 26-Jun-2026, DOI: 10.35248/2161-1122.26.16.766

Citation: Danylak L (2026). Photobiomodulation in Contemporary Dental Practice: Expanding Clinical Applications for Tissue Repair and Patient Comfort. J Dentistry. 16:766.

Copyright: © 2026 Danylak L. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

with active phases of tooth formation when odontogenic tissues are still present and responsive.

CONCLUSION

The causes of ameloblastic fibro-odontoma appear to be multifactorial, involving developmental disturbances in odontogenic tissues, genetic expression variations, local environmental influences, and possible systemic factors that affect cell behavior during tooth formation. Rather than a single identifiable cause, the condition likely results from a combination of disrupted regulatory mechanisms that govern normal dental tissue development. Continued research into molecular signaling pathways and developmental biology of odontogenic tissues may provide further clarity regarding the origins of this lesion and its relationship to other mixed odontogenic entities.