

Facts, Challenges, Difficulties and Hopes in Single-Cell Biology: Physiopathological Studies

Francesco Salvatore^{1,2*}, Federica Cariati^{1,3} and Rossella Tomaiuolo^{1,3}

¹CEINGE-Biotecnologie Avanzate, via G. Salvatore 486, 80145 Naples, Italy

²IRCCS-Fondazione SDN, 80143 Naples, Italy

³Department of Molecular Medicine and Medical Biotechnologies, University of Naples Federico II, via Pansini 5, 80131 Naples, Italy

Abstract

Single-cell approaches are being increasingly used to unravel the many diverse mechanisms underlying biological processes that characterize each cell irrespective of the influx of other cells even within the same tissue. Consequently, the interference of metabolites and nervous stimuli emanating from the circulatory or nervous system in a higher organism like man is avoided. However, while the single-cell approach yields a wealth of data about single-cell metabolism and internal regulatory mechanisms, information about interactions and interrelations among similar or dissimilar cells may remain obscure. Starting from these considerations, here we summarize, without attempting to be exhaustive, some areas in which we think single-cell biological studies could be effective in translational medicine and in other areas of applied sciences. In this short review we describe the facts, challenges and perspectives related to these issues.

Keywords: Single cells; Single cancer cells; Pharmacology in single cells; Preimplantation genetic diagnosis; Preimplantation genetic screening

Commentary

Here we briefly recapitulate concepts and experimental findings that can enhance our understanding of physiological phenomena and/or their variations that lead to disease. Furthermore, we mention, albeit with a reductionist approach, the areas where breakthroughs have been made and where challenges remain, and what we may reasonably expect the future to bring.

Studies conducted with single cancer cells have been particularly fruitful, mainly as regards the capacity of these cells to act as early harbingers of the disease or of tumor relapse as well as a model to understand cancer pathogenesis, spreading and metastatization [1-4]. Furthermore, nucleic acid sequencing is starting to shed light on the molecular features of single cancer cells. Based on this information one may dissect events that belong to a specific cell population or organ whether healthy or diseased [5]. This applies to the diverse fields of physiopathology [6], particularly those of nanobiology [7] and immunology [8], and during development and tissue mosaicism [9].

Facts and perspectives are now emerging in the field of the three-D structural architecture of the genome [10,11], and in the field of the electrical characterization of single cells [12]. The latter aspect is particularly important for studies conducted with nervous, muscle and cardiac tissues using, for instance, such sophisticated patch-clamp, ion diffusion and the very recent microfluidic technique [13].

Also in the field of pharmacology, a single cell can be studied and measured *in vivo* and studies can be conducted on committed cells starting from stem cells [14]. Furthermore, studies based on pharmacokinetics and pharmacodynamics using different drugs can help to determine the exact cell target where the “bullet” will operate to modify an altered status to a possible normal status [15,16]. This approach will help to define the pathway through which a stem cell reaches the final cell-fate decision thereby overcoming the difficulty inherent in studies of a population of multiple-fate cells [17]. For example, heterogeneity is extensive in hematopoietic stem and progenitor cells as well as in multipotent progenitors [18,19]. Notably,

the root hair plant model in a single cell is generating data, also through a developmental approach, that have relevant consequences for the overall biological equilibrium of our planet [20].

A field in which single-cell analysis is already having a great impact on human health is the field of genome biology and medicine. In fact, single-cell genomics is being used in preimplantation genetic diagnosis, not only for aneuploidy screening but also to avoid single-gene disorders [21], chromosomal abnormalities [22], and to prevent transmission of X-linked disorders [23]. Preimplantation genetic diagnosis is an important feature of human reproduction not only because of ethical considerations but also in preventing the transmission of severe inherited disease in affected families [24]. In this context, ethic problems such as sex selection also with regard to social sexing should be addressed with a very open-minded philosophical attitude [25]. In the table are reported the most important facts, challenges, difficulties and hopes that appear to be the most fruitful areas to pursue future research in single-cell biomedicine.

To sum up, we prospectively illustrate some examples that are closely related to the contents of this Short Review: (i) The application of single-cell analysis in the field of cancer is rapidly expanding and promises to shed light on how a single cell (mainly a single stem cell) can become metastatic or lead to cancer progression and invasion. Furthermore, in cancer biology, it would in principle be more effective to dissect biomarkers in single-cell analysis in order to detect not only the early appearance of the transformation process but the tissue types from which the primary cancer originated; (ii) Single-cell analysis

***Corresponding author:** Francesco Salvatore, M.D., Ph.D., CEINGE-Biotecnologie Avanzate; via Gaetano Salvatore 486, 80145 Naples, Italy, Tel: +39 081 7463133; Fax: +39 081 7463650; E-mail: salvator@unina.it

Received August 10, 2015; **Accepted** September 10, 2015; **Published** September 12, 2015

Citation: Salvatore F, Cariati F, Tomaiuolo R (2015) Facts, Challenges, Difficulties and Hopes in Single-Cell Biology: Physiopathological Studies. Single Cell Biol 4: 121. doi:10.4172/2168-9431.1000121

Copyright: © 2015 Salvatore F, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

	Facts	Challenges	Difficulties	Hopes	References	
Cancer biology	<ul style="list-style-type: none"> Bulk tissue genomic analysis provides key insights into cancer biology but does not provide information about spatio-temporal cell interactions and the genetic events occurring during tumor progression 	<ul style="list-style-type: none"> To define clonal heterogeneity in tumor To detect mutations and perform gene-expression studies on 'cancer stem cells' To identify circulating tumor cells originating from a primary tumor 	<ul style="list-style-type: none"> Isolation of single tumor cells thereby minimizing their loss and genomic degradation Reduction of the noise generated during sampling due to multiple layers of cell heterogeneity in cancer 	<ul style="list-style-type: none"> To detect subclones of cancer cells in order to shed light on cell interactions, on the development and control of metastatic disease, drug resistance and disease progression To detect parts of cell components (e.g., exosomes, microvesicles) as biomarkers or tools for pathological studies 	1-5,26-29	
Cell population within tissue	<ul style="list-style-type: none"> Analysis of bulk tissue samples does not reveal cell-to-cell variations or rare cells that may play an important role in disease progression Single-cell sequencing (SCS) provides detailed comprehensive data about individual cells 	<ul style="list-style-type: none"> Neurobiology 	<ul style="list-style-type: none"> Neurons are one of the most morphologically diverse cell populations Single-cell RNA sequencing is a powerful unbiased approach with which to classify neurons based on their transcriptional profiles 	<ul style="list-style-type: none"> Single neuron RNA sequencing must be combined with other techniques (e.g., electrophysiology and microfluidics) to obtain a more comprehensive functional profile from a specific brain microenvironment 	<ul style="list-style-type: none"> SCS may be a novel approach with which to classify neuronal cell types and identify possible DNA diversity in neuronal cell populations 	30-33
		<ul style="list-style-type: none"> Immunology 	<ul style="list-style-type: none"> The immune system is constituted by a wide variety of cell types that work together in a highly integrated fashion to recognize and eliminate antigens 	<ul style="list-style-type: none"> The transcriptional heterogeneity within cell types in response to a variety of antigens 	<ul style="list-style-type: none"> Unbiased RNA SCS methods may be used to investigate heterogeneous transcriptional responses in immune cells after antigen activation to: <ol style="list-style-type: none"> Analyze cells that show variable responses when stimulated under different conditions <i>in vitro</i>; Identify multimodal gene expression patterns of differentially stimulated cells 	8,34-36
		<ul style="list-style-type: none"> Tissue mosaicism 	<ul style="list-style-type: none"> The traditional view of somatic tissues is that normal single cells have identical genomes. However, this dogma is beginning to be challenged by increasing evidence of genetic mosaicism in normal tissues that arises during normal development 	<ul style="list-style-type: none"> Most studies have analyzed bulk tissue samples. Therefore, controversy exists about the prevalence of mosaic mutations and whether they can simply be explained by technical error 	<ul style="list-style-type: none"> SCS methods may provide a novel approach with which to resolve cell-to-cell variations in normal tissues at an unprecedented genomic resolution 	9,37,38
3D genome architecture	<ul style="list-style-type: none"> The chromosome organization and its spatial nuclear organization are linked to the control of gene expression, DNA replication and repair The 3D genomic organization is cell type-specific 	<ul style="list-style-type: none"> The hierarchical 3D chromosome organization must be studied to fully understand the transcriptional behaviour of the gene 	<ul style="list-style-type: none"> Chromosome conformation capture methods (Hi-C) enables one to assess the probabilistic chromosome conformation averaged from millions of cells, but not to relate the chromosome conformation to the single genome activity pattern 	<ul style="list-style-type: none"> The recently developed single cell Hi-C protocol assesses the spatial organization of the entire genome in an individual nucleus and may provide structural information about individual cell variability 	10, 11, 39	
Pharmacology	<ul style="list-style-type: none"> To enhance drug discovery and development it is crucial to measure pharmacokinetics (PK) and pharmacodynamics (PD) <i>in vivo</i> at single cell level 	<ul style="list-style-type: none"> Single-cell PK measurements 	<ul style="list-style-type: none"> Traditional PK, which is based on the measurement of plasma concentration, does not provide information about the cellular heterogeneity of drug distribution 	<ul style="list-style-type: none"> The cellular location within tissue can modify the concentration time course of a drug and can significantly affect treatment response 	<ol style="list-style-type: none"> To follow drug distribution and uptake at single-cell level using intravital microscopy. In fact, multichannel imaging may be used to identify different cell populations and to measure drug uptake within these cells <i>in vivo</i>; Measurement of a drug concentration in different cell types within a tissue may be a means to understand and optimize local or targeted drug delivery 	14-17
		<ul style="list-style-type: none"> Single-cell PD measurements 	<ul style="list-style-type: none"> To establish drug efficacy one must measure and understand not just the drug-target engagement but also the downstream PD response 	<ul style="list-style-type: none"> The time-course of drug distribution and target interaction differs significantly at single-cell level 	<ol style="list-style-type: none"> To evaluate the downstream PD response at single-cell level through intravital microscopy: fluorescent genetic reporters may allow the real-time measurement of relevant cellular pathways To monitor the real-time distribution and single-cell uptake of drugs would provide an integrated PD/PK perspective 	
Preimplantation genetic diagnosis	<ul style="list-style-type: none"> Single gene mutation Chromosomal abnormality Mitochondrial DNA copy number 	<ul style="list-style-type: none"> A comprehensive approach is required for the simultaneous detection of monogenic and chromosomal disorders 	<ul style="list-style-type: none"> DNA contamination Single-cell based PCR artefacts: allele drop out, random failure of amplification of one allele at a locus, the richness of the locus in G/C bases, chimeric DNA molecules and copy error Biologically variable noise: the cell-cycle stage of the isolated cell 	<ul style="list-style-type: none"> To develop a genome-wide method using next-generation sequencing for single-cell genome characterization to accurately determine genomic variations, such as copy number variations and single or small nucleotide variations simultaneously 	21-25, 40-44	

Table: Facts, challenges, difficulties and hopes in single-cell biology.

of cell populations within a given tissue will reveal differences in the whole genome RNA transcriptome. The main cell differences that derive from gene expression analysis are pertinent not only to each cell type but above all to pathological aspects during a disease. These cells can also characterize a cell genomic mosaicism which raises additional questions; (iii) Pharmacological studies at both pharmacodynamic and pharmacokinetic level can be more precisely focused when single-cell analysis is characterized as illustrated in point i; (iv) Preimplantation genetic diagnosis could revolutionize the timing and hence the impact of molecular diagnosis and the efficiency of *in vitro* fertilization techniques. In fact, once the genome-wide method becomes optimized, it will be possible to evaluate a single gene disorder and chromosomal abnormalities simultaneously. In this context, next-generation sequencing may lead to the development of a genome-wide procedure for single-cell genomics.

In conclusion, we think that the era of single cell biology is still in its infancy and that the next few years will see unprecedented scientific progress along the lines prospected herein.

References

1. Wills QF, Mead AJ (2015) Application of single-cell genomics in cancer: promise and challenges. *Hum Mol Genet* 24: R74-84.
2. Navin N, Kendall J, Troge J, Andrews P, Rodgers L, et al. (2011) Tumour evolution inferred by single-cell sequencing. *Nature* 472: 90-94.
3. Lohr JG, Adalsteinsson VA, Cibulskis K, Choudhury AD, Rosenberg M, et al. (2014) Whole-exome sequencing of circulating tumor cells provides a window into metastatic prostate cancer. *Nat Biotechnol* 32: 479-484.
4. Morimoto A, Mogami T, Watanabe M, Iijima K, Akiyama Y, et al. (2015) High-Density Dielectrophoretic Microwell Array for Detection, Capture, and Single-Cell Analysis of Rare Tumor Cells in Peripheral Blood. *PLoS One* 10: e0130418.
5. Voet T, Kumar P, Van Loo P, Cooke SL, Marshall J, et al. (2013) Single-cell paired-end genome sequencing reveals structural variation per cell cycle. *Nucleic Acids Res* 41: 6119-6138.
6. Wang Y, Navin NE (2015) Advances and applications of single-cell sequencing technologies. *Mol Cell* 58: 598-609.
7. Shi X, Zhang X, Xia T, Fang X (2012) Living cell study at the single-molecule and single-cell levels by atomic force microscopy. *Nanomedicine (Lond)* 7: 1625-1637.
8. Satija R, Shalek AK (2014) Heterogeneity in immune responses: from populations to single cells. *Trends Immunol* 35: 219-229.
9. Machiela MJ, Zhou W, Sampson JN, Dean MC, Jacobs KB, et al. (2015) Characterization of large structural genetic mosaicism in human autosomes. *Am J Hum Genet* 96: 487-497.
10. Furlan-Magaril M, Várnai C, Nagano T, Fraser P (2015) 3D genome architecture from populations to single cells. *Curr Opin Genet Dev* 31: 36-41.
11. Nagano T, Lubling Y, Stevens TJ, Schoenfelder S, Yaffe E, et al. (2013) Single-cell Hi-C reveals cell-to-cell variability in chromosome structure. *Nature* 502: 59-64.
12. Mansor MA, Ahmad MR (2015) Single Cell Electrical Characterization Techniques. *Int J Mol Sci* 16: 12686-12712.
13. Chen J, Xue C, Zhao Y, Chen D, Wu MH, et al. (2015) Microfluidic impedance flow cytometry enabling high-throughput single-cell electrical property characterization. *Int J Mol Sci* 16: 9804-9830.
14. Vinegoni C, Dubach JM, Thurber GM, Miller MA, Mazitschek R, et al. (2015) Advances in measuring single-cell pharmacology in vivo. *Drug Discov Today* 20: 1087-1092.
15. Thurber GM, Reiner T, Yang KS, Kohler RH, Weissleder R (2014) Effect of small-molecule modification on single-cell pharmacokinetics of PARP inhibitors. *Mol Cancer Ther* 13: 986-995.
16. Thurber GM, Yang KS, Reiner T, Kohler RH, Sorger P, et al. (2013) Single-cell and subcellular pharmacokinetic imaging allows insight into drug action in vivo. *Nat Commun* 4: 1504.
17. Regot S, Hughey JJ, Bajar BT, Carrasco S, Covert MW (2014) High-sensitivity measurements of multiple kinase activities in live single cells. *Cell* 157: 1724-1734.
18. Weston W, Zayas J, Perez R, George J, Jurecic R (2014) Dynamic equilibrium of heterogeneous and interconvertible multipotent hematopoietic cell subsets. *Sci Rep* 4: 5199.
19. Wu J, Tzanakakis ES (2013) Deconstructing stem cell population heterogeneity: single-cell analysis and modeling approaches. *Biotechnol Adv* 31: 1047-1062.
20. Qiao Z, Libault M (2013) Unleashing the potential of the root hair cell as a single plant cell type model in root systems biology. *Front Plant Sci* 4: 484.
21. Van der Aa N, Zamani Esteki M, Vermeesch JR, Voet T (2013) Preimplantation genetic diagnosis guided by single-cell genomics. *Genome Med* 5: 71.
22. Fragouli E, Wells D (2012) Aneuploidy screening for embryo selection. *Semin Reprod Med* 30: 289-301.
23. Christofidou C, Sofocleous C, Vrettou C, Destouni A, Traeger-Synodinos J, et al. (2009) PGD for X-linked and gender-dependent disorders using a robust, flexible single-tube PCR protocol. *Reprod Biomed Online* 19: 418-425.
24. Harper JC, Wilton L, Traeger-Synodinos J, Goossens V, Moutou C, et al. (2012) The ESHRE PGD Consortium: 10 years of data collection. *Hum Reprod Update* 18: 234-247.
25. Eftekhari TE, Nejatizadeh AA, Rajaei M, Soleimani S, Fallahi S, et al. (2015) Ethical considerations in sex selection. *J Educ Health Promot* 4: 32.
26. McGranahan N, Swanton C (2015) Biological and therapeutic impact of intratumor heterogeneity in cancer evolution. *Cancer Cell* 27: 15-26.
27. Grün D, Lyubimova A, Kester L, Wiebrands K, Basak O, et al. (2015) Single-cell messenger RNA sequencing reveals rare intestinal cell types. *Nature* 525: 251-255.
28. Heitzer E, Auer M, Gasch C, Pichler M, Ulz P, et al. (2013) Complex tumor genomes inferred from single circulating tumor cells by array-CGH and next-generation sequencing. *Cancer Res* 73: 2965-2975.
29. Raposo G, Stoorvogel W (2013) Extracellular vesicles: exosomes, microvesicles, and friends. *J Cell Biol* 200: 373-83.
30. Sarin S, Antonio C, Tursun B, Hobert O (2009) The *C. elegans* Tailless/TLX transcription factor nhr-67 controls neuronal identity and left/right asymmetric fate diversification. *Development* 136: 2933-2944.
31. Darmanis S, Sloan SA, Zhang Y, Enge M, Caneda C, et al. (2015) A survey of human brain transcriptome diversity at the single cell level. *Proc Natl Acad Sci U S A* 112: 7285-7290.
32. Qiu S1, Luo S, Evgrafov O, Li R, Schroth GP, et al. (2012) Single-neuron RNA-Seq: technical feasibility and reproducibility. *Front Genet* 3: 124.
33. Pollen AA, Nowakowski TJ, Shuga J, Wang X, Leyrat AA, et al. (2014) Low-coverage single-cell mRNA sequencing reveals cellular heterogeneity and activated signaling pathways in developing cerebral cortex. *Nat. Biotechnol* 32: 1053-8.
34. Shay T, Kang J (2013) Immunological Genome Project and systems immunology. *Trends Immunol* 34: 602-609.
35. Jaitin DA, Kenigsberg E, Keren-Shaul H, Elefant N, Paul F, et al. (2014) Massively parallel single-cell RNA-seq for marker-free decomposition of tissues into cell types. *Science* 343: 776-779.
36. Kumar RM, Cahan P, Shalek AK, Satija R, DaleyKeyser AJ, et al. (2014) Deconstructing transcriptional heterogeneity in pluripotent stem cells. *Nature* 516: 56-61.
37. Biesecker LG, Spinner NB (2013) A genomic view of mosaicism and human disease. *Nat Rev Genet* 14: 307-320.
38. Samuels ME, Friedman JM (2015) Genetic mosaics and the germ line lineage. *Genes (Basel)* 6: 216-237.
39. van Berkum NL, Lieberman-Aiden E, Williams L, Imakaev M, Gnirke A, et al. (2010) Hi-C: a method to study the three-dimensional architecture of genomes. *J Vis Exp* (39) pii: 1869.
40. Dreesen J, Destouni A, Kourlaba G, Degn B, Mette WC, et al. (2014) Evaluation of PCR-based preimplantation genetic diagnosis applied to monogenic diseases: a collaborative ESHRE PGD consortium study. *Eur J Hum Genet* 22: 1012-1018.

41. Fuccio A, Iorio M, Amato F, Elce A, Ingino R, et al. (2011) A novel DHPLC-based procedure for the analysis of COL1A1 and COL1A2 mutations in osteogenesis imperfecta. *J Mol Diagn* 13: 648-656.
42. Konstantinidis M, Alfarawati S, Hurd D, Paolucci M, Shovelton J, et al. (2014) Simultaneous assessment of aneuploidy, polymorphisms, and mitochondrial DNA content in human polar bodies and embryos with the use of a novel microarray platform. *Fertil Steril* 10: 1385-1392.
43. Konstantinidis M, Prates R, Goodall NN, Fischer J, Tecson V, et al. (2015) Live births following Karyomapping of human blastocysts: experience from clinical application of the method. *Reprod Biomed Online* pii: S1472-6483(15)00259-X.
44. Tan Y, Yin X, Zhang S, Jiang H, Tan K, et al. (2014) Clinical outcome of preimplantation genetic diagnosis and screening using next generation sequencing. *Gigascience* 3: 30.