OPEN ACCESS Freely available online

ISSN: 2155-9546
Journal of Aquaculture Research & Development

Commentary

Fish Bile Acid Profiling: A Molecular Approach to Digestive Function and Aquaculture Nutrition

Yubexi Xie*

Department of Aquaculture, National Taiwan Ocean University, Keelung, Taiwan

DESCRIPTION

Fish bile acids are increasingly being examined as useful biochemical indicators of digestive activity, nutrient utilization, intestinal microbial interactions, and metabolic condition in cultured aquatic animals. Bile is produced mainly by the liver and stored in the gallbladder in many fish species before being released into the digestive tract. Its major components include bile acids, phospholipids, cholesterol, pigments, proteins, and electrolytes. Among these compounds, bile acids have attracted particular interest because they participate in lipid digestion and also function as signaling molecules that influence metabolic processes. Profiling these molecules can therefore provide information extending beyond conventional measurements of feed intake, growth rate, and feed conversion.

Bile acids facilitate the digestion and absorption of dietary lipids by interacting with fats and forming micellar structures within the intestinal environment. This process improves the accessibility of lipids to digestive enzymes such as lipases. Efficient lipid digestion is especially important in aquaculture because dietary oils and other lipid sources are frequently used to supply concentrated energy and essential fatty acids. Changes in dietary lipid composition can influence bile acid synthesis, secretion, recycling, and intestinal activity. Measuring bile acid profiles may therefore help researchers understand how fish respond to different feed formulations.

Fish species differ considerably in their bile acid composition. Some species contain bile acids that are closely associated with those found in mammals, whereas others possess distinctive compounds related to their evolutionary history and dietary ecology. Freshwater and marine species may also display differences in bile acid metabolism because environmental salinity, feeding habits, intestinal physiology, and energy requirements can affect digestive processes. Carnivorous fish, herbivorous fish, omnivorous fish, and detritivorous species may consequently exhibit different patterns of bile acid production and utilization.

Modern analytical techniques allow researchers to detect and quantify a wide range of bile acid molecules in biological samples. Liquid chromatography coupled with mass spectrometry is particularly useful because it can separate compounds according to their chemical properties and detect molecules according to their mass-to-charge ratios. High-resolution mass spectrometry can provide additional information regarding molecular composition. Nuclear magnetic resonance spectroscopy can also contribute to metabolite profiling, although its sensitivity and analytical requirements differ from those of mass spectrometric methods.

The interaction between bile acids and intestinal microorganisms has become an important research area in fish nutrition. Certain intestinal bacteria can transform primary bile acids into secondary or modified forms through enzymatic reactions. These microbial transformations can alter the biological properties of bile acids and influence their movement through the digestive tract. Conversely, bile acids can affect microbial community composition because some microorganisms are more tolerant of bile-associated conditions than others. This two-way interaction means that bile acid measurements can be considered alongside intestinal microbiome analysis when studying digestive health.

Feed composition is another major factor affecting bile acid profiles. Diets containing different levels or sources of protein, lipid, fiber, and plant-derived ingredients may produce distinct biliary responses. Replacement of marine-derived ingredients with plant proteins can alter digestive physiology because plant materials contain different nutrient profiles and various non-nutritional compounds. Changes in dietary oils can also affect lipid digestion and the demand for bile-mediated emulsification. Monitoring bile acids during feeding trials may therefore provide an additional biochemical measure for evaluating new aquafeed formulations.

Bile acid profiling may also be relevant to fish exposed to environmental stressors. Temperature changes, variations in dissolved oxygen, salinity shifts, crowding, and poor water

Correspondence to: Yubexi Xie, Department of Aquaculture, National Taiwan Ocean University, Keelung, Taiwan, E-mail: yubexixie@mail.ntou.edu.tw

Received: 03-Aug-2026, Manuscript No. JARD-26-32966; **Editor assigned:** 06-Aug-2026, PreQC No. JARD-26-32966 (PQ); **Reviewed:** 20-Aug-2026, QC No. JARD-26-32966; **Revised:** 27-Aug-2026, Manuscript No. JARD-26-32966 (R); **Published:** 03-Sep-2026, DOI: 10.35248/2155-9546.26.17.1139

Citation: Xie Y (2026). Fish Bile Acid Profiling: A Molecular Approach to Digestive Function and Aquaculture Nutrition. J Aquac Res Dev. 17.1139.

Copyright: © 2026 Xie Y. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

quality can modify metabolic activity. Because the liver is involved in both nutrient processing and bile acid synthesis, physiological stress may influence the concentration and composition of circulating or biliary compounds. Repeated profiling during controlled experiments could help researchers determine whether observed metabolic changes are associated with environmental conditions, dietary treatment, or both.

The relationship between bile acids and energy metabolism is also gaining attention. Certain bile acids can interact with cellular receptors and signaling pathways involved in glucose regulation, lipid metabolism, and energy expenditure. Although many of these mechanisms have been investigated extensively in mammals, fish research is still developing. Differences among fish species mean that mammalian findings cannot simply be transferred to aquatic animals without experimental validation. Species-specific studies are therefore needed to establish appropriate interpretations of bile acid changes.

CONCLUSION

Research combining metabolomics, microbiome sequencing, digestive physiology, and gene-expression measurements could provide a more complete understanding of biliary regulation in cultured fish. Fish bile acid profiling represents a valuable biochemical approach for examining digestion, lipid utilization, intestinal microbial activity, and metabolic responses. Its usefulness is particularly relevant to aquaculture nutrition, where efficient nutrient utilization has direct importance for fish growth and feed efficiency. As analytical methods become more accessible and species-specific datasets increase, bile acid measurements can become an informative component of nutritional and physiological studies involving cultured aquatic animals.