OPEN ACCESS Freely available online

ISSN: 2155-9546
Journal of Aquaculture Research & Development

Short Communication

# Fish Osmoregulatory Physiology: Ionic Balance, Water Regulation, and Environmental Adaptation

**Antoine Perol***

*Department of Aquatic Systems, University of Montpellier, Montpellier, France*

## DESCRIPTION

Fish live in aquatic environments where the concentration of dissolved salts can differ greatly from the internal conditions of their bodies. Maintaining a suitable balance of water and dissolved ions is therefore an essential physiological function. Fish osmoregulatory physiology describes the processes through which aquatic animals control body fluids despite continuous movement of water and ions across the skin, gills, digestive tract, and other tissues [1]. Sodium, chloride, potassium, calcium, magnesium, and other ions are required for nerve activity, muscle contraction, cellular signaling, enzyme activity, and maintenance of normal cellular volume. Water balance is equally important because excessive water entry or loss can disturb cell function and reduce physiological performance [2].

The osmoregulatory strategy of a fish is strongly influenced by the environment in which it lives. Freshwater fish generally have body fluids that contain higher concentrations of dissolved ions than the surrounding water. Water therefore tends to enter their bodies through permeable surfaces, while ions can move outward [3]. Marine teleosts experience the opposite general situation because seawater contains a much greater concentration of dissolved salts than their internal fluids. These fish tend to lose water and gain ions from the environment. Their physiological systems respond differently according to habitat, life stage, species, and environmental conditions [4].

The gills are major organs involved in ionic and acid-base regulation. Gill tissues contain specialized cells commonly called ionocytes, which transport ions between the blood and the surrounding water. These cells contain membrane proteins, ion channels, exchangers, and transport enzymes that regulate the movement of sodium, chloride, calcium, and other substances. Their activity can change when environmental salinity changes. Such flexibility allows many fish species to maintain relatively stable internal fluid conditions even when external water chemistry varies [5].

Freshwater fish generally produce large quantities of dilute urine because water continuously enters the body through osmosis. Their kidneys recover valuable ions from the filtrate while removing excess water. At the gills, ion uptake mechanisms assist in maintaining concentrations of sodium and chloride in the blood. Dietary intake also contributes to mineral balance, although the importance of food varies among species and environmental conditions. The combined activity of the gills, kidneys, digestive tract, and endocrine system allows freshwater fish to maintain internal stability [6].

Marine teleosts face a different physiological challenge. Because seawater is more concentrated than their internal body fluids, these fish lose water across permeable surfaces. They compensate by drinking seawater and absorbing water through the digestive tract. During this process, substantial quantities of salts enter the body. Specialized gill cells actively remove excess sodium and chloride back into the surrounding seawater. The kidneys also contribute by controlling the elimination of divalent ions such as magnesium and sulfate. Urine production in many marine teleosts is relatively limited compared with that of freshwater fish because conserving water is important [7].

The endocrine system provides an important regulatory network for these processes. Hormones influence ion transport, kidney activity, drinking behavior, and changes in gill cell function. Cortisol, prolactin, growth hormone, and several other signaling molecules have been associated with salinity adaptation in different fish groups. Their effects vary according to species and environmental conditions. Hormonal signals can modify the number and activity of transport proteins, alter cellular metabolism, and change the behavior of organs involved in water and ion regulation [8].

Euryhaline fish are particularly useful for studying osmoregulation because they can tolerate substantial changes in environmental salinity. Species such as salmon, mullet, tilapia, and several estuarine fishes may move between freshwater and seawater during different stages of their life cycle. When

**Correspondence to:** Antoine Perol, Department of Aquatic Systems, University of Montpellier, Montpellier, France, E-mail: antoine.perol@umontpellier.fr

**Received:** 27-May-2026, Manuscript No. JARD-26-32735; **Editor assigned:** 29-May-2026, PreQC No. JARD-26-32735 (PQ); **Reviewed:** 12-Jun-2026, QC No. JARD-26-32735; **Revised:** 19-Jun-2026, Manuscript No. JARD-26-32735 (R); **Published:** 26-Jun-2026, DOI: 10.35248/2155-9546.26.17.1117

**Citation:** Perol A (2026). Fish Osmoregulatory Physiology: Ionic Balance, Water Regulation, and Environmental Adaptation. J Aquac Res Dev. 17:1117.

**Copyright:** © 2026 Perol A. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

environmental salinity changes, their gill, kidney, intestine, and endocrine systems undergo coordinated adjustments. Ion transport pathways can be modified, drinking behavior can change, and energy allocation can shift toward physiological processes required for maintaining internal balance [9].

Salinity adaptation requires energy because active ion transport depends on cellular metabolism. Mitochondria provide energy for many transport processes occurring in gill ionocytes and other tissues. Changes in salinity can therefore influence oxygen consumption and metabolic demand. When fish experience an abrupt environmental change, energy must be allocated between osmoregulation, growth, movement, feeding, immune activity, and reproduction [10].

## CONCLUSION

Fish osmoregulation is therefore a coordinated physiological process involving the gills, kidneys, intestine, endocrine system, nervous system, blood, and cellular transport mechanisms. These systems work together to maintain water content, ion concentrations, and acid-base conditions within suitable limits. Differences between freshwater and marine environments have produced distinct regulatory strategies, while euryhaline species demonstrate substantial flexibility when moving across salinity gradients. Continued study of ionic transport, hormonal regulation, metabolism, and environmental interactions can improve understanding of fish physiology and contribute to more effective management of aquatic production systems.

## REFERENCES

1. Faleiros RO, Furriel RP, McNamara JC. Transcriptional, translational and systemic alterations during the time course of Osmoregulatory acclimation in two palaemonid shrimps from distinct osmotic niches. Comp Biochem Physiol A Mol Integr Physiol. 2017;212:97-106.
2. Principe SC, Augusto A, Costa TM. Differential effects of water loss and temperature increase on the physiology of fiddler crabs from distinct habitats. J Therm Biol. 2018;73:14-23.
3. Shrivastava J, Ndugwa M, Caneos W, De Boeck G. Physiological trade-offs, acid-base balance and ion-osmoregulatory plasticity in European sea bass (*Dicentrarchus labrax*) juveniles under complex scenarios of salinity variation, ocean acidification and high ammonia challenge. Aquat Toxicol. 2019;212:54-69.
4. Vargas-Chacoff L, Arjona FJ, Ruiz-Jarabo I, García-Lopez A, Flik G, Mancera JM. Water temperature affects osmoregulatory responses in gilthead sea bream (*Sparus aurata L.*). J Therm Biol. 2020;88:102526.
5. Cuenca AL, Souza MM, Freire CA. Osmoregulatory power influences tissue ionic composition after salinity acclimation in aquatic decapods. Comp Biochem Physiol A Mol Integr Physiol. 2021;259:111001.
6. Seale AP, Breves JP. Endocrine and Osmoregulatory responses to tidally-changing salinities in fishes. Gen Comp Endocrinol. 2022;326:114071.
7. Yulianto B, Soegianto A, Affandi M, Payus CM. The impact of various periods of mercury exposure on the osmoregulatory and blood gas parameters of tilapia (*Oreochromis niloticus*). Emerg. Contam. 2023;9(3):100244.
8. Zhao H, Zuo Z, Yang L, Zhang L, Lv T, Yu D, et al. Similarities and differences in the physiological adaptation to water salinity between two life forms of aquatic plants in alpine and arid wetlands. Sci Total Environ. 2024;908:168449.
9. Vieira CE, De Jesus AC, Branco GS, Fadini PS, Dos Santos DM, Moreira RG. Sublethal effects of diclofenac and ibuprofen, individually and in mixture, in a Neotropical fish: a multi-biomarker assessment under environmentally realistic concentrations. Aquat Toxicol. 2025:107609.
10. Wei Y, Shi C, Wang F, Lai Q, Gao P, Zhou K, et al. Single-cell transcriptomics identifies a candidate Osmoregulatory cell population in the mid-intestine of the saline-alkaline fish *Gymnocypris przewalskii*. Comp Biochem Physiol A Mol Integr Physiol. 2026:112050.