



Irrigation Systems for Efficient Water Management and Improved Performance

Félix Vecilla*

Department of Agro technology, Polytechnic University of Madrid, Madrid, Spain

DESCRIPTION

Sensor-based irrigation has become an important approach to improving water management in agricultural production. Conventional irrigation often follows fixed schedules based on calendar dates or general observations of crop appearance. Such practices may result in excessive watering during periods when soil contains sufficient moisture or insufficient watering when plants require additional water. Sensor-based systems provide continuous information about soil and environmental conditions, allowing irrigation decisions to be based on measurements collected directly from the growing area. This approach can improve water-use efficiency, support healthy crop development, and reduce unnecessary irrigation costs.

Agricultural water demand varies according to crop type, soil characteristics, weather conditions, growth stage, and field location. A crop growing in sandy soil may require more frequent irrigation than the same crop growing in soil with greater water-holding capacity. Similarly, plants exposed to high temperatures and dry winds may lose water more rapidly than plants growing under cooler conditions. Sensors help identify these changes by collecting measurements at regular intervals. When combined with an irrigation controller, the collected information can determine when irrigation should begin and how much water should be supplied.

Soil moisture sensors are among the most widely used devices in sensor-based irrigation systems. These sensors measure the amount of water present within the soil and transmit information to a controller or digital platform. Different sensor technologies measure soil moisture through electrical, dielectric, or other physical properties. Installation depth is selected according to crop rooting patterns and soil characteristics. For shallow-rooted crops, sensors may be positioned closer to the surface, while deeper placement can be useful for crops with extensive root systems. Multiple sensors at different depths can provide a more complete picture of water availability within the root zone.

Temperature and humidity sensors can provide additional information for irrigation management. Air temperature affects evaporation and plant water loss, while relative humidity influences the movement of water from leaves into the atmosphere. When temperature is high and humidity is low, crop water demand can increase. Combining atmospheric measurements with soil moisture readings allows irrigation systems to respond to both soil and weather conditions. This can provide better irrigation timing than relying on soil moisture information alone.

Weather stations can further improve irrigation decisions by recording rainfall, wind speed, solar radiation, temperature, and humidity. Rainfall data are particularly useful because natural precipitation can reduce or eliminate the need for scheduled irrigation. If an adequate amount of rain falls, an automated system can postpone irrigation until soil moisture declines below the selected threshold. This prevents water from being applied unnecessarily and reduces the possibility of saturated soil conditions.

Crop growth stage is another factor that influences irrigation requirements. Young plants generally have smaller root systems and may require different irrigation intervals than mature plants. During flowering, fruit formation, or grain filling, some crops have increased water requirements. Sensor-based systems can combine crop growth information with soil and weather data to adjust irrigation schedules throughout the growing season. This helps maintain suitable moisture conditions during periods when water availability strongly influences crop development.

Wireless communication has expanded the usefulness of agricultural sensors. Modern systems can transmit field measurements through wireless networks to computers, tablets, or mobile phones. Farmers can monitor soil moisture and environmental conditions without visiting every section of a field. Alerts can be generated when moisture levels become too low, equipment stops functioning, or water use exceeds a selected range. Remote monitoring is particularly useful for large farms where frequent physical inspection of every irrigation zone may be difficult.

Correspondence to: Félix Vecilla, Department of Agro technology, Polytechnic University of Madrid, Madrid, Spain, E-mail: felix.vecilla@upm.es

Received: 29-Aug-2025, Manuscript No. AGT-25-32493; **Editor assigned:** 01-Sep-2025, Pre QC No. AGT-25-32493 (PQ); **Reviewed:** 15-Sep-2025, QC No. AGT-25-32493; **Revised:** 22-Sep-2025, Manuscript No. AGT-25-32493 (R); **Published:** 29-Sep-2025, DOI:10.35248/2168-9891.25.14.419

Citation: Vecilla F (2025) Irrigation Systems for Efficient Water Management and Improved Performance. Agrotechnology. 14:419.

Copyright: © 2025 Vecilla F. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

CONCLUSION

Sensor-based irrigation provides a practical method for managing agricultural water according to measured field conditions. By combining soil moisture information with weather observations, crop growth data, automated controls, and efficient water-delivery systems, farmers can make irrigation

decisions with greater accuracy. Proper sensor placement, regular equipment maintenance, reliable data collection, and suitable user training are necessary for effective operation. When these elements are integrated into farm management, sensor-based irrigation can reduce unnecessary water use, support crop growth, lower energy consumption, and contribute to more efficient agricultural production.