OPEN ACCESS Freely available online

ISSN: 2155-9546
Journal of Aquaculture Research & Development

Opinion Article

Kinetic Profiling of Aquafeed Breakdown and Time-Dependent Nutrient Assimilation

Linh Taher*

Department of Water, Atmosphere and Environment, University of Natural Resources and Life Sciences, Vienna, Austria

DESCRIPTION

Aquafeed digestibility kinetics describes the rate and extent at which nutrients and feed components are broken down, absorbed, and utilized by cultured aquatic animals. Feed digestibility is often reported as a single percentage, but digestion is a dynamic process that changes with feeding rate, ingredient composition, animal size, water temperature, gastrointestinal transit, enzyme activity, and physiological condition. Studying the timing of nutrient release provides additional information about how efficiently a formulated diet supports growth and metabolic activity. This approach is particularly useful in aquaculture because fish, shrimp, and other cultured species differ substantially in digestive anatomy, enzyme production, feeding behavior, and nutrient requirements.

The digestion of an aquafeed begins soon after ingestion. Physical disruption of feed particles increases their contact with digestive fluids, while enzymes act on proteins, lipids, carbohydrates, and other organic components. Proteins are converted into peptides and amino acids, lipids are separated into smaller components and fatty acids, and carbohydrates are transformed into simpler molecules that can be absorbed through the intestinal surface. The speed of these processes is influenced by feed structure and ingredient characteristics. A diet containing highly digestible protein sources may release amino acids rapidly, whereas ingredients containing substantial structural carbohydrates or resistant compounds may remain in the digestive tract for longer periods.

Digestibility kinetics can therefore be considered in terms of both extent and time. Apparent digestibility coefficients indicate the proportion of a nutrient that appears to be absorbed from the gastrointestinal tract, while kinetic measurements examine how nutrient availability changes after feeding. Sampling at different intervals can reveal when digestion is most active and when nutrients begin to leave the digestive system. Such information can help researchers compare ingredients that produce similar final digestibility values but have different digestion rates.

Protein digestion is especially important in aquafeed evaluation because amino acids are required for tissue formation, enzyme production, immune functions, and other biological activities. Fish species with strong gastric digestion may begin substantial protein hydrolysis in the stomach, while species without a conventional stomach depend more heavily on intestinal and pancreatic processes. Shrimp also possess a distinctive digestive system involving the hepatopancreas, which contributes digestive enzymes and nutrient processing. These anatomical differences mean that the same feed ingredient can produce different digestion profiles among cultured species.

Lipid digestion follows another temporal pattern. Dietary lipids provide concentrated energy and supply essential fatty acids required for cellular membranes, growth, reproduction, and physiological regulation. Lipid digestion depends on emulsification, lipase activity, bile-associated processes, and intestinal absorption. The physical form of dietary oil can affect its accessibility to digestive enzymes. Excessive lipid inclusion may also alter gastric or intestinal passage and influence the timing of nutrient absorption. Measuring lipid disappearance at several post-feeding intervals can therefore provide useful information beyond a final digestibility value.

Carbohydrate digestion is equally influenced by ingredient structure. Starch from cereal-based ingredients can be rapidly hydrolyzed when properly processed, while complex polysaccharides may resist enzymatic breakdown. Extrusion, cooking, grinding, and other feed-processing methods can modify starch structure and particle characteristics. These changes may affect how quickly carbohydrates become available after ingestion. In carnivorous fish, which generally have lower tolerance for high carbohydrate inclusion than many omnivorous species, understanding this timing can assist in evaluating alternative formulations.

Correspondence to: Linh Taher, Department of Water, Atmosphere and Environment, University of Natural Resources and Life Sciences, Vienna, Austria, E-mail: linh.taher@boku.ac.at

Received: 17-Aug-2026, Manuscript No. JARD-26-32982; **Editor assigned:** 20-Aug-2026, PreQC No. JARD-26-32982 (PQ); **Reviewed:** 03-Sep-2026, QC No. JARD-26-32982; Revised: 10-Sep-2026, Manuscript No. JARD-26-32982 (R); **Published:** 17-Sep-2026, DOI: 10.35248/2155-9546.26.17.1154

Citation: Taher L (2026). Kinetic Profiling of Aquafeed Breakdown and Time-Dependent Nutrient Assimilation. J Aquac Res Dev. 17.1154.

Copyright: © 2026 Taher L. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Feed particle size has a direct relationship with digestive kinetics. Smaller particles generally provide a larger surface area for enzymatic action, although excessively fine feed can create practical problems such as nutrient loss in water and poor feeding behavior. Pellet density, water stability, texture, and expansion also influence how feed changes after immersion. Once a pellet enters the digestive tract, hydration can alter its physical structure and affect the accessibility of nutrients. Consequently, feed manufacturing conditions should be considered when interpreting digestibility measurements.

CONCLUSION

Aquafeed digestibility kinetics provides a quantitative framework for examining how nutrients move through the digestive process after feeding. Instead of treating digestibility as a single endpoint, kinetic analysis considers the sequence and timing of nutrient release and absorption. This perspective can support ingredient selection, feed formulation, feeding management, and waste reduction in aquaculture. Careful measurement of digestion rates, combined with information on animal physiology and feed characteristics, can contribute to more efficient use of dietary nutrients and better matching between feed composition and the biological needs of cultured aquatic species.