

## Magnitude, Characteristics, Maternal and Feto-neonatal Outcomes of Obstetric Emergencies in Western Ethiopia, Nekemte, Ethiopia, 2017

Ashenafi Habte Woyessa<sup>1\*</sup>, Jote Markos Cafo<sup>2</sup> and Thanasekaran P<sup>3</sup>

<sup>1</sup>Department of Emergency and Critical Care Nursing, Wollega University, Institute of Health Sciences, school of Nursing and Midwifery, Nekemte, Ethiopia; <sup>2</sup>Department of Nursing, Wollega University, Institute of Health Sciences, School of Nursing and Midwifery, Nekemte, Ethiopia; <sup>3</sup>Department of Psychiatric Nursing, Wollega University, Institute of Health Sciences, School of Nursing, Nekemte, Ethiopia

### ABSTRACT

**Aim:** In Ethiopia very little or probably nothing is known about significance of obstetric emergencies. This study was therefore aimed at assessing magnitude, characteristics, and outcomes of obstetric emergencies in western Ethiopia.

**Methods:** Institution based prospective study was employed from January to June, 2017. To select the hospitals, area sampling technique was used. Total of 567 pregnant women with obstetric emergencies presented and treated in respective hospitals during the study periods and meet the inclusion criteria were consecutively included.

**Result:** Majority (91.7%) of the identified obstetric emergencies have led to termination of pregnancy. Significant proportions (11%) of pregnant women who reached health facility were died of obstetric emergencies. Pregnant women with obstetric emergencies travelled to facility carried by people r were found to die about 8 times more likely as compared to those who transported by ambulance. While 29.21% were normal life births, still birth and neonatal death were 8.02% and 7.4% respectively. Higher number of neonatal death was also observed among mothers in whom final mode of delivery was cesarean section (AOR: 0.19(0.05, 0.62) compared to spontaneous vaginal delivery.

**Conclusion:** This study revealed that obstetric emergencies are responsible for significant number of maternal and perinatal death. If the women have been accessed early and received optimum emergency care, many cases of the occurred death would have been prevented. Better outcome can be achieved through maximum utilization of quality and comprehensive antenatal care and organized pre-hospital obstetric emergency services.

Keywords: Obstetric emergencies; Maternal; Feto-neonatal outcome; Western Ethiopia

### INTRODUCTION

Obstetric emergencies are one of the leading causes of maternal mortality in our globe. World Health Organization estimates that maternal mortality varies up to 100-fold, from 10 in wealthier countries to about 1,000 in under developed nations. Despite tremendous efforts made to reduce death that could arise in relation to pregnancy and childbirth, maternal death still remains a significant problem in many developing countries [1].

Sub Saharan African countries share the highest maternal death rate in the world. Annually about 500 maternal deaths per 100,000 live births representing over half of the total world maternal

deaths occur [2]. In Ethiopia maternal and neonatal mortality rates are extremely in the highest in the world. The 2011 Ethiopian Demographic Health Survey revealed that the maternal death rate was 676 deaths per 100,000 live births and neonatal mortality was 37 deaths per 1,000 live births. In other words, for every 1,000 live births, about seven women died during pregnancy, child birth, or within two months of child birth [3,4].

Lack of organized pre hospital services, inadequate health infrastructures, financial constraints, illiteracy, and low health seeking behaviors worsen the problem in developing countries [5]. Evidence indicate that the high maternal, neonatal and child

\*Correspondence to: Ashenafi Habte Woyessa, Department of Emergency and Critical Care Nursing, Wollega University, Institute of Health Sciences, school of Nursing and Midwifery, Nekemte, Ethiopia, Tel: +251 922183085; FAX: +251576617980; E-mail: ashexman2015@gmail.com

Received: March 21, 2019, Accepted: March 25, 2019, Published: April 02, 2019

Citation: Woyessa AH, Cafo JM, Thanasekaran P. Magnitude, characteristics, maternal and feto-neonatal outcomes of obstetric emergencies in western Ethiopia, Nekemte, Ethiopia, 2017. Clinics Mother Child Health. 2019;16:319. doi:10.24105/2090-7214.1000319

Copyright: © 2019 Woyessa, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

mortality rates are associated with inadequate and poor quality maternal health care [6]. A great deal of studies indicate that an estimated 74% of maternal deaths due to obstetric emergencies could have been averted if all women had access to the interventions for preventing and/ or treating all pregnancy related emergencies [7-10].

Primary prevention where possible and/or effective management of obstetric emergencies plays key role in the effort made to reduce an ever increasing maternal mortality [5]. In many third world countries obstetric emergencies like bleeding, prolonged labor, sepsis, toxemia of pregnancy, ruptured uterus, abortion and its complications are the commonly encountering obstetric emergencies [6,11]. Information of obstetric emergencies in terms of types, magnitude, and characteristic are crucial especially in countries like Ethiopia where the MMR is not yet satisfactorily reduced. Nevertheless, in Ethiopia, very little or probably nothing is known about significance of obstetric emergencies in causing maternal and neonatal morbidity and mortality.

This study was therefore undertaken to indicate the significance of obstetric emergencies in causing maternal and neonatal mortality as end clinical outcomes. The findings of this study are very much helpful for all agents including policy makers and other stakeholders who mainly work on maternal health care.

## MATERIAL AND METHODS

### Study setting

This study was conducted in three administrative zones of Wollega, Western Ethiopia. Five hospitals were selected (one from each zone). Namely, Nekemte referral hospital from East Wollega Zone, Nedjo, Mendi and Gimbi hospitals from West Wollega Zone, and Shambu Hospital from Horo Guduru Wollega Zone. The study was carried out from January 1st to June 30, 2017 among pregnant women who visited the selected hospitals for obstetric emergencies.

### Study design, Sampling procedure and techniques

Hospital based prospective cross sectional study was conducted the study among pregnant women presented to selected

hospitals in western Ethiopia, from January to June 2017. To select the hospitals, area sampling technique was used. All pregnant women with obstetric emergencies presented and treated in respective hospitals during the study periods and meet the inclusion criteria were consecutively included.

### Research tools, data processing and analysis

Data was collected by using structured and pre tested interviewer administered tool for the study purpose. The questionnaire was comprised of four parts (socio demographic characteristics, obstetric history, Types of obstetric emergencies, and maternal and perinatal outcomes). Data was initially entered in to Epi-data version 7 and later the exported to SPSS program Version 20 for data analysis. Descriptive statistics were generated and possible associations among independent variables and outcome variables were examined in bivariate analysis. Odds ratio and 95% CI was used as a measure of association.

## RESULTS

The study explored the patterns and clinical outcomes of obstetric emergencies on mothers, fetus and newborn among pregnant women who visited four hospitals in western Ethiopia the last six month duration. A total of 567 pregnant women with various obstetric emergencies were included in the study. Of which, 530 were successfully interviewed and the analyzed yielding a response rate of 93.47 percent.

### Socio demographic characteristics the respondents

This section looks at the socio demographic characteristics distribution including age, ethnicity, level of education, occupation, and decision making authority on reproductive issues. The mean age of the women in this study was 25.83 and more than half (52.6 %) were between the age of 25-34 years. As depicted in table below, majority (86.2%) of the respondents were Oromo in their ethnicity followed by Amhara (6.8%). Of prime concern, this study identified that the decision making authority on fertility issues was 20.8%, 44.0% and 34.3% herself, her husband and both of the couples respectively (Table 1).

**Table 1:** Distribution of respondents by their socio demographic characteristics, western Ethiopia, Nekemte, 2017.

No	Variables	Category	Frequency	Percent
1	Age of the mother	15-24	213	40.2
		25-34	279	52.6
		>35	38	7.2
2	Ethnicity	Oromo	457	86.2
		Amhara	36	6.8

		Tigre	11	2.1
		Gurage	8	1.5
		Other	18	3.4
3	Marital status	Single	71	13.4
		Married	441	83.2
		Divorced	9	1.7
		Widowed	9	1.7
4	Level of education attained	No schooling	160	30.2
		Primary school	185	34.9
		Secondary school	120	22.6
		College diploma	38	7.2
		Degree and above	27	5.1
5	Occupation of mother	Government employee	47	8.9
		Private business	85	16
		Servant	14	2.6
		House wife	247	46.6
		Farmer	94	17.7
		Other	43	8.1
6	Occupation of husband/partner	Government employee	61	11.5
		Private business	165	29.5
		Farmer	259	48.9
		other	54	10.2
7	Residence	Town	229	43.2
		Rural	301	56.8

### Findings related to base line information and obstetric history

The decision making authority on fertility issues was found to be 20.8%, 44.0% and 34.3% respectively for the mother,

husband and both of the couples. Most importantly, this study revealed that more than half (55.1%) pregnant mothers who visited the hospitals for obstetric emergencies had no emergency fund (Table 2).

**Table 2:** Distribution of respondents by their base line information in western Ethiopia, Nekemte, Ethiopia, 2017.

No	Base line information	Category	Frequency	Percent
1	Mode of arrival	Being carried by people	10	1.9

		Public car	114	21.5
		Ambulance	345	65.1
		On foot	56	10.6
		Other	5	0.9
2	Treatment given in ambulance	Yes	276	52.1
		No	254	47.9
3	Forms of treatment given in ambulance(N=276)	Positioning	161	58.3
		Resuscitation withy fluid given	68	24.6
		Supplemental oxygen	20	7.2
		Other	27	9.7
4	Decision making on fertility issues	Her self	110	20.8
		Her husband/partner	233	44
		Both couples	182	34.3
		His/her family	5	0.9
5	Presence of emergency fund	Yes	238	44.9
		No	292	55.1
6	Arrival time	Day time	290	54.7
		Night time	146	27.5
		On weekend	91	17.7
7	Time elapsed between arrival and initiation of treatment in minutes	<15	456	86
		15-30	38	7.2
		30-60	21	4
		>60	15	2.8
8	Mothers' habit	Alcohol intake	64	12.1
		Smoking	8	1.5
		Fasting	37	7
		Chew chat	5	0.9
		Have no history of habits	416	78.5
9	Distance from home to nearest health facility in kilometer	<5	114	21.5
		May-15	120	22.6
		15-30	205	38.7

	>30	91	17.2	
	knows some of obstetric danger signs	182	34.33	
10	Women's knowledge and readiness for emergencies	Knows nearby ambulance Number	242	45.7
		Knows selection of place of delivery	443	83.6

This study has explored pregnant mothers' relevant obstetric history. About 21.9% have not attended ANC. Among the women who have been attended ANC, only 29.9% of them attended four times. Common types of preexisting medical conditions identified during ANC were chronic hypertension (55%) followed by 'other' (16%) problems like anemia and respiratory tract infections. Even though ambulance was means

of arrival for majority (65.1%) of respondents, nearly half (47.1%) of them did not receive any form of treatment before arrival. The study also revealed that more than half (55.1%) had no emergency fund. Time elapsed between arrival and initiation of treatment in minutes for majority (86.0%) of them was <15 mins (Table 3).

**Table 3:** Distribution of respondents by their obstetric history in western Ethiopia, Nekemte, Ethiopia, 2017.

No	Obstetric history	Category	Frequency	Percent
1	Type of pregnancy	Wanted/planned	450	84.9
		Unwanted/unplanned	80	15.1
2	Parity	Primi	181	34.2
		Para 2 - 4	276	52.1
		Para 5 and above	73	13.8
3	ANC follow up	Yes	414	78.1
		No	116	21.9
4	Time of the first ANC visit	First trimester	206	38.9
		Second trimester	196	37
		Third trimester	21	4
5	Number of ANC visit(N= 414)	Once	38	9.1
		Twice	111	26.8
		Three times	152	36.7
		Four times and above	124	29.9
6	History of violence	Beating	24	4.5
		Stabbing	4	0.8
		Throwing	10	1.9
		Kicking	7	1.5
		Rape	33	6.2
		Control of finance	33	6.2

	Other	7	1.5
	No history of abuse	412	77.7
	Yes	41	7.7
7	Presence of preexisting diseases	No	471
		Unknown	18
			88.9
			3.4

Chronic hypertension (55%) followed by 'other' (16%) problems like anemia and respiratory tract infections were among the identified preexisting medical or surgical problems during ANC visit. Out of all respondents, 88 (16.6%) of respondents have found to have some disorders during ANC visit (Table 4).

**Table 4:** Types of newly diagnosed disorders during ANC visit in western Ethiopia, Nekemte, Ethiopia, 2017.

Preexisting disease	Frequency(N=88)	Percent
Hypertension	44	50
Anemia	28	31.88
Bleeding	7	7.95

Hyperemesis gravidrum	1	1.14
Weight loss	3	3.41
others	5	5.68
Total	88	100

### Findings related to diagnosed obstetric emergencies

One of the aims of this study was identifying magnitude and commonly encountering obstetric emergencies in the study area. The most common diagnosed obstetric emergencies were obstructed labor (17.9%) followed by abortion (12.5%). Antepartum hemorrhage (11.3%) and preterm labor (7.9 %) were also common diagnoses (Table 5).

**Table 5:** Distribution of diagnosed of obstetric emergency by age of respondents in western Ethiopia, Nekemte, Ethiopia, 2017.

Patterns of obstetric emergency	Age of respondents			Total	
	15-24	25-34	>35	Frequency	percent
Retained placenta	4	23	0	27	5.1
Preterm labor	15	27	0	42	7.9
Premature rupture of membrane	15	23	0	38	7.2
Ante partum hemorrhage	25	33	2	60	11.3
Postpartum hemorrhage	4	17	2	23	4.3
Severe anemia	1	6	0	7	1.3
Retained second baby	3	1	0	4	0.8
Ectopic pregnancy	0	2	1	3	0.6
Disseminated intravascular coagulation	0	0	5	5	0.9
Cord prolapse	0	7	1	8	1.5
Puerperal sepsis	0	1	1	5	0.6
Obstructed labor	40	51	4	95	17.9

Mal presentation	10	19	3	32	6
Abortion	49	8	9	66	12.5
Severe pre eclampsia	15	17	2	34	6.4
Eclampsia	13	2	3	18	3.4
Uterine rupture	1	14	3	18	3.4
Fetal distress	7	18	0	25	4.7
Intra uterine fetal death	7	3	2	12	2.3
Uterine inversion	0	2	0	2	0.4
Others	3	5	0	8	1.5
Total	213	279	38	530	100

Majority (91.7 %) of identified cases of obstetric emergencies in this study have led to termination of pregnancy. As to the place of delivery, out of hospital delivery i.e. home delivery and on street delivery were 9 (1.79%) and 16 (3.03%) respectively. Delivery conducted outside of health facility, the delivery was assisted by untrained birth attendants (8%), health extension workers (56%), Emergency Medical and technicians (20%) and family members (16%). Among pregnant women in whom obstetric emergencies have led to termination of pregnancy, the final mode of delivery was identified. While emergency caesarian section was final mode of delivery form most mothers (36.62%), about 3.49% of the women have undergone laparotomy for ectopic pregnancy and ruptured uterus (Figure 1).




Figure 1: Final mode of delivery due to obstetric emergencies in western Ethiopia, Nekemte, Ethiopia, 2017.

### Maternal outcomes of obstetric emergencies

The study identified the consequences of obstetric emergencies on mothers' health and life. Nearly two third (57.07%) of the women were peacefully discharged after termination of pregnancy. A significant proportion (11%) of maternal death due to obstetric emergencies was recorded (Figure 2).




Figure 2: Distribution of respondents by their maternal outcomes of obstetric emergencies in western Ethiopia, Nekemte, Ethiopia, 2017.




Figure 3: Underlying cause of maternal death in obstetric emergencies in western Ethiopia, Nekemte, Ethiopia, 2017.

The study also explored the underlying causes of maternal death. The main contributing factors for death were obstetric hemorrhage (31.9%) followed pregnancy related infections

(21.27%). In about 15%, the underlying cause of death was unknown (Figure 3).

The main contributing factors for maternal death were obstetric hemorrhage (31.9%) followed pregnancy related infections (21.27%). Mothers who did not attend ANC were more likely to die of obstetrics emergencies as compared to mothers who received ANC service (P-value: 0.00). Mothers who travel longer

than 15 kilometer to reach health facility were about 10 times more likely to die of obstetrics emergencies as compared to mothers who travel less than 5 kilometer (P-value: 0.00). Pregnant women with obstetric emergencies travelled to facility carried by people should were found to die about 8 times more likely as compared to those who transported by ambulance (Table 6).

**Table 6:** Binary logistic regression of maternal obstetric emergency outcomes with selected study variable, Nekemte, Ethiopia, 2017.

Variables	Category	Crude		Adjusted	
		OR (95%CI)	P-value	OR (95%CI)	P-value
Age of the Women	15-24	4.89(1.82, 13.15)	0.00	3.68(0.76,17.87))	0.10
	25-34	2.39( 0.99, 5.71)	0.05	1.21(0.28,5.14)	0.78
	35 and above	R	R	R	R
Mode of facility arrival	By people/foot	1.42(0.53,3.76)	0.48	7.92(1.76, 35.70)	0.00*
	Public car	2.19(0.90, 5.34)	0.08	5.30(1.18,23.81)	0.02*
	Ambulance	R	R	R	R
Presence of emergency fund	Yes	R	R	R	R
	No	0.67(0.36, 1.25)	0.21	1.29(.400, 4.19)	0.666
Marital status	Single	0.93(0.37, 2.30)	0.88	0.28(.05,1.57)	0.15
	Divorced	0.04(0.01, 0.18)	0.00	0.008(0.00,0.11)	0.00*
	Married	R	R	R	R
Distance from facility in kilometer	<5	1.05(0.46, 2.39)	0.89	1.24(.0.40,3.83 )	0.70
	6-15	1.52(0.70, 3.28)	0.28	10.08( 2.90,34.97)	0.00*
	16-30	2.21(0.75, 6.45)	0.14	16.79(3.46,81.28 )	0.00*
	>31	R	R	R	R
Time of treatment initiation	<15 minute	R	R	R	R
	16- 30 minutes	4.05(0.54, 30.23)	0.173	2.613(.243, 28.137)	0.428
	>30 minutes	1.53(0.19, 11.9)	0.68	.274(.022,3.473)	0.318
Pre-existing illness	No	R	R	R	R
	Yes	0.24(0.109, 0.53)	0.00	0.24(.08,0.72)	0.01*
	Unknown	0.36(0.09,1.32)	0.12	0.52(0.10, 2.65)	0.43
Type of pregnancy	Wanted/planned	R	R	R	R
	Unwanted	0.45(0.22, 0.92)	0.03	2.27(.204, 25.33)	0.50
Arrival times	Day time	R	R	R	R



	Night time	0.51(0.27,0.97)	0.043	.918(.377, 2.237 )	0.85
	On weekend	1.43(0.52,3.91)	0.47	2.60(.60, 11.31)	0.20
ANC attendance	Yes	R	R	R	R
	No	0.41(0.21,0.76)	0.00	0.15(0.05, 0.45)	0.00*
Decision making role on sexuality	Both Couples	R	R	R	R
	Husband alone	0.37(0.20, 0.709)	0.002	0.09(0.02, 0.31)	0.00*

Duration of hospital stay following the diagnosed obstetric emergency or related problem was considered as possible outcome. Among admitted cases, while majority (83%) of them stayed in hospital less than of 10 days, the hospital duration of stay for significant proportion (10%) of women was 15 days and above. Among commonly identified underlying causes of ward admission, pregnancy related infection accounted contributed 28.23% followed by obstetric hemorrhage (25.8%) (Table 7).


**Table 7:** Binary logistic regression of maternal obstetric emergency outcomes with selected study variable, Nekemte, Ethiopia, 2017.

Cause of ward/ICU admission	Frequency(N=85)	Percent
Complications of Abortion	6	7.05
Obstetric hemorrhage	22	25.88
Hypertensive disorders	12	14.11
Pregnancy related infection	24	28.23
Medical, surgical/mental disease complications	8	9.41
Suboptimal management	11	12.94
coincidental conditions	2	2.35
Total	85	100

**Feto-neonatal outcomes of obstetric emergencies**

This research has looked the fate of infants and neonates with a mother diagnosed of obstetric emergency. About 29.21% of

women gave normal life birth. Still birth and neonatal death were 8.02% and 7.4% respectively (Figure 4).



**Figure 4:** Feto- neonatal outcomes of obstetric emergencies in western Ethiopia, in western Ethiopia, Nekemte, Ethiopia, 2017.

The most common underlying cause of neonatal death was noticed to be distress/asphyxia (39%) followed by neonatal sepsis (33%).

The remaining death was contributed by hypothermia (11%) and other causes (17%).

Infants born from mothers arriving hospitals on weekend days carried higher chance of death (P- vale: 0.01).

Higher number of neonatal death was also observed among mothers in whom final mode of delivery was cesarean section (AOR: 0.19(0.05, 0.62) compared to spontaneous vaginal deliver (Table 8).

**Table 8:** Underlying cause of ward admission after obstetric emergencies in western.

Variables	Category	Crude OR(95%CI)	P-vale	Adjusted OR (95%CI)	P-value
Mode of facility arrival	People /on foot	1.20(0.45,3.23)	0.7	0.53(0.14,1.98)	0.35
	Public car	3.76(1.12,12.55)	0.03	3.62(0.85,15.31)	0.08
	Ambulance	R	R	R	R

Residence	Urban	R	R	R	R
	Rural	0.63(0.31, 1.25)	0.18	0.50(0.18,1.40)	0.19
Marital status	Divorced	0.32(0.06,1.61)	0.16	0.02(0.00,0.28)	0.01*
	Married	R	R	R	R
Distance from facility in kilometer	<5	R	R	R	
	Jun-15	0.42(0.15,1.13)	0.08	0.46(0.14,1.55)	0.21
	16-30	1.54(0.50, 4.72)	0.44	2.05(0.48,8.80)	0.33
	>31	0.36(0.13,1.01)	0.05	0.47(0.10,2.11)	0.32
Time of treatment initiation	<15 minute	R	R	R	R
	16- 30 minutes	3.38(0.45,25.3)	0.23	2.80(0.30,26.26)	0.36
Type of pregnancy	Wanted	R	R	R	R
	Unwanted	3.38(0.79,14.34)	0.09	0.50(0.02,8.87)	0.63
Arrival times	Day time	R	R	R	R
	Night time	0.44(0.21,0.92)	0.03	0.63(0.26,1.52)	0.3
	On weekend	0.57(0.23,1.40)	0.22	0.20(0.05,0.71)	0.01*
ANC attendance	Yes	R	R	R	R
	No	11.71(1.59,86.27)	0.01	3.60(0.39,33.07)	0.25
Decision making role on sexuality matter	Both Couples	R	R	R	R
	Husband alone	0.65(0.34,1.25)	0.2	0.96(0.39,2.35)	0.94
Mother's substance use	No	R	R	R	R
	Yes	2.11(0.63,7.03)	0.22	25.19(1.89,334.3)	0.01*
Final mode of delivery	SVD	R	R	R	R
	MVA	2.31(0.26,20.22)	0.44	0.07(0.00,1.12)	0.06
	Cesarean section	0.19(0.07,0.51)	0	0.19(0.05,0.62)	0.00*
	Instrumental Delivery	2.53(0.29,22.15)	0.4	4.00(0.38,41.30)	0.24

## DISCUSSION

Globally, obstetric emergencies take the top rank in causing of maternal mortality. Literacy, poverty, lacks of quality antenatal care, low transportation facilities and insufficient equipment and providers amplify the problem in underdeveloped nations [3-8]. Ethiopia is one of the countries blamed to have the highest maternal death in the world. It has been reported that the country is one of the six countries that contribute about over 50% of the deaths [12,13]. Poor health care seeking habit lesser coverage of health services, unavailability of sufficient transport

facilities, deep rooted poverty and wrong beliefs are among the few factors known to worsen the end outcomes obstetric emergency in developing countries [14,15].

Very socio-economic status a community certainly contributes to negative outcomes of obstetric emergency. In present study, while in higher proportion (44.0%) husband or partners were mandated for decision making on fertility issues, only 110 (20.8%) women have authority for decision making. Women of higher socio-economic status usually receive better pregnancy outcomes. Because they are relatively better informed, likely to

develop better choices, more likely to develop and implement birth readiness and are more empowered to make decisions in case of they come across with obstetric emergencies [16-18].

In this study, 414(78.1%) of pregnant women who have visited the selected hospitals for obstetric emergencies had received ANC at least once. Study done in Kenya to assess fetal outcome of obstetric emergencies reported that 93.4% of women attended an antenatal clinic at least once. The difference could be due to better awareness and service of ANC as a result of the difference of socio economic status between the two countries [19].

When faced with obstetric emergency that might need referral, arranging a private car consume time and. Delays to reach facilities and misdiagnosis of cases are directly related to near miss mortality. In this study, 345 (65.1%) patients have used ambulance to reach the hospital which is higher as compared with the finding from India study [20-22].

Nearly two third (57.07%) of the women were peacefully discharged from hospitals after termination of pregnancy. Significant number, 47(11%) of pregnant women who reached health facility were died of their obstetric emergencies. This is in line with a study done in south west Nigeria which reported were 17 maternal deaths out of 262 obstetric emergencies and greater than 4% of Indian study [19,23]. It is also noticed that fetal and neonatal death were noticed to be 8.02% and 7.4% respectively As reported, in this study there were 11.52% low birth weight and 7.8% pre term babies while the remaining 7.2% were admitted to the neonatal intensive care units.

In conclusion, this study has shown that obstetric emergencies do occur and found to be responsible for significant number of the maternal and perinatal mortality. If the women have been accessed early and received optimum emergency help both during transportation and in health facilities, many of the occurred death would have been avoided. Better outcome can therefore be achieved through maximum utilization of quality and comprehensive antenatal care and organized pre-hospital and in hospital obstetric emergency services.

## DECLARATION

**Ethics approval and consent to participate:** The research was approved by Wollega University (Reference Number / 100, 617/1-26). Formal letter of permission and support was written to the respective health facilities. Prior to collecting the required data both verbal consent and written consent were taken from all of the study subjects.

**Availability of data and materials:** the raw data supporting our findings are available from authors on a reasonable request.

**Consent for publication:** Consent to publish this work has been taken from the approving committee

## AUTHOR'S CONTRIBUTION

AH designed and developed the research proposal, involved in supervision of data collection process, statistical analysis and preparation of the final document. JM helped with research tool

preparations, supervising data collection, and involved in revision of the final manuscript.

## ACKNOWLEDGEMENT

We would like to acknowledge Wollega University for funding this research. We would also like to extend our heartfelt gratitude to our study participants for their cooperation in responding to the questions whilst they were in physical and emotional sufferings.

## REFERENCES

1. Buekens P. Is estimating maternal mortality useful? Bull world health organ. 2001;79:179.
2. UN and African union commission: Report on progress in achieving the millennium development goals in Africa.2013
3. Chukwudebelu WO, Okonofua A, Odunsi K. Preventing maternal mortality in developing countries. In contemporary obstetrics and gynaecology for developing countries. 2013;644.
4. Federal democratic republic of ethiopia central statistical agency om 2015: edhs 2015. Addis Ababa, Ethiopia.
5. Haines A, Cassels A. Can the millennium development goals be attained. British Medical J. 2004;329:394-397.
6. The millennium development goals 2012.
7. Tarekegn SM, Lieberman LS, Giedraitis V. Determinants of maternal health service utilization in Ethiopia: Analysis of the 2011 Ethiopian demographic and health survey. BMC Pregnancy and Child birth.2014;14:161.
8. World Health Organization: WHO maternal and child health fact sheet,2012.
9. Federal democratic republic of Ethiopia central statistical agency OM: EDHS. Addis Ababa, Ethiopia.2015.
10. WHO, UNFPA, UNICEF and World bank 2012: Trends in maternal mortality: 1990-2010.
11. Drife J, Lueslay DM, Baker PN (2004) "Maternal mortality" in obstetrics and gynecology and evidence-based text for MRCOG, pp: 196-204.
12. Birmeta K, Dibaba Y, Woldeyohannes D (2013) Determinants of maternal health care utilization in Holeta town, central Ethiopia, birmeta et al. BMC health services research 13:256.
13. Hogan CH, Foreman KJ, Naghavi M, Ahn SY, Wang M, et al. (2010) Maternal mortality for 181 countries. 1980 - 2008: A systematic analysis of progress towards millennium development goal 5. Lancet 376: 1389-1390.
14. Najam R, Gupta S, Chowdhury H (2015) Pattern of obstetrical emergencies and fetal outcomes in a tertiary care center. Acta Med Int 2: 105-110.
15. Catherin N, Anushila SR, Ramakrishna B (2014) Goud 3 obstetric emergencies presenting to a rural community maternity hospital 2: 264-269.
16. Bhandari TR, Dangal G (2014) Emergency obstetric care: strategy for reducing maternal mortality in developing countries. NJOG 17: 8-16.
17. Nwobodo EL (2006) Obstetric emergencies as seen in a tertiary health institution in North-Western Nigeria: Maternal and fetal outcome. Nigerian Medical Practitioner 49: 54-55.
18. Drife J, Lueslay DM, Baker PN (2004) Maternal mortality in obstetrics and gynecology and evidence-based text for MRCOG. pp: 196-204.

19. Najam R, Gupta S, Chowdhury H (2015) Pattern of obstetrical emergencies and fetal outcomes in a tertiary care center. *Acta Med Int* 2: 105-110.
20. Maryland R (2016) Ethiopia sy. Central statistical agency (CSA).
21. Elizabeth W (2011) Maternal and fetal outcomes among women with obstetric emergencies referred to Kenyatta national hospital, Kenya. pp: 32- 38.
22. Shivani B (2017) Pattern of term obstetric emergencies referred from health centers to Rims. *Emir* 4: 376-382.
23. Zhang WH, Alexander S, Bouvier-Colle MH, Macfarlane A (2015) Incidence of severe pre-eclampsia, postpartum hemorrhage and sepsis as surrogate marker for severe maternal morbidity in a European-based study: MOMS survey. *BJOG* 112: 89-96.