



Maternal Nutritional Status and Its Influence on Maternal and Child Health

Daniel Foster*

Department of Pediatrics, Faculty of Medicine, University of Helsinki, Helsinki, Finland

DESCRIPTION

Maternal nutrition is a fundamental aspect of reproductive and child health that influences the well-being of women during pregnancy and the growth and development of their children. Adequate intake of essential nutrients before conception, throughout pregnancy, and during lactation supports maternal health, fetal development, and successful early childhood outcomes. Pregnancy increases the body's nutritional requirements due to physiological changes, expansion of maternal tissues, and the need to support fetal growth [1]. Nutritional deficiencies or imbalances during this period may affect pregnancy outcomes and increase the risk of complications for both mothers and infants. Therefore, assessment and improvement of maternal nutritional status remain important components of comprehensive maternal healthcare. A woman's nutritional status before pregnancy has a significant influence on fetal growth and pregnancy progression. Preconception nutrition helps establish adequate nutrient reserves required for the increased demands of pregnancy. Factors such as undernutrition, obesity, micronutrient deficiencies, and inadequate dietary diversity before conception may influence maternal health and fetal development [2].

During pregnancy, nutrients including iron, folate, calcium, iodine, vitamin D, protein, and essential fatty acids contribute to the formation of fetal organs, development of the nervous system, maintenance of maternal tissues, and preparation for breastfeeding. Balanced nutrition combined with appropriate antenatal care supports healthy pregnancy outcomes and reduces the likelihood of nutrition-related complications. Iron deficiency is one of the most common nutritional concerns among pregnant women worldwide [3]. Increased blood volume during pregnancy raises the requirement for iron, which is necessary for the production of hemoglobin and oxygen transport to maternal and fetal tissues. Insufficient iron intake may result in maternal anemia, which can contribute to fatigue, reduced physical capacity, and increased risk of adverse birth outcomes. Folic acid is another important nutrient required during early pregnancy

because it supports fetal neural tube development and reduces the risk of neural tube defects. Adequate intake of these micronutrients through diet and supplementation when recommended forms an important part of antenatal nutritional care [4].

Maternal nutrition also affects fetal growth and birth outcomes. Inadequate maternal weight gain, insufficient protein intake, or micronutrient deficiencies may contribute to restricted fetal growth and low birth weight. Infants with poor growth during pregnancy may face increased risks of developmental challenges and health concerns during childhood. On the other hand, excessive maternal weight gain and high intake of energy-dense foods may increase the likelihood of gestational diabetes, hypertension, and complications during delivery. Maintaining appropriate nutritional balance helps support healthy fetal development while reducing pregnancy-related risks [4].

Breastfeeding further emphasizes the importance of maternal nutrition after childbirth. Lactating mothers require additional energy and nutrients to support breast milk production and maintain their own health. A varied diet containing adequate amounts of proteins, fruits, vegetables, whole grains, and healthy fats contributes to the nutritional quality of breast milk and supports infant growth [5]. Maternal dietary counseling during the postpartum period helps women understand their nutritional needs, recover after childbirth, and continue providing appropriate nourishment for their infants. Supportive healthcare practices encourage successful breastfeeding and contribute to improved maternal and child health outcomes [6].

Maternal nutrition is influenced by social, economic, cultural, and environmental factors. Limited access to nutritious foods, food insecurity, lack of nutrition knowledge, and healthcare inequalities may affect dietary quality among pregnant women. Community-based nutrition programs, antenatal counseling, and public health initiatives play an important role in addressing these challenges [7]. Healthcare providers can support mothers by assessing nutritional status, providing individualized dietary advice, monitoring weight changes, and recommending appropriate

Correspondence to: Daniel Foster, Department of Pediatrics, Faculty of Medicine, University of Helsinki, Helsinki, Finland, E-mail: daniel.foster.dan@outlook.com

Received: 31-Mar-2026, Manuscript No. CMCH-26-32330; **Editor assigned:** 02-Apr-2026, Pre QC No. CMCH-26-32330; **Reviewed:** 16-Apr-2026, QC No. CMCH-26-32330; **Revised:** 23-Apr-2026, Manuscript No. CMCH-26-32330; **Published:** 30-Apr-2026, DOI: 10.35248/2090-7214.26.23.577

Citation: Foster D (2026). Maternal Nutritional Status and Its Influence on Maternal and Child Health. Clinics Mother Child Health. 23:577.

Copyright: ©2026 Foster D. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

supplementation when needed. Involving families and communities in nutrition education can further improve maternal dietary practices and promote healthier pregnancies [8].

Nutritional assessment during pregnancy should include evaluation of dietary patterns, body weight, medical history, and possible nutrient deficiencies. Regular antenatal visits provide opportunities to identify nutritional concerns and provide guidance tailored to individual needs [9]. Modern approaches, including digital health tools and nutrition monitoring systems, have created additional opportunities for improving access to maternal nutrition information and follow-up care. Integrating nutrition services into maternal healthcare programs strengthens preventive care and supports healthier outcomes for mothers and children [10].

CONCLUSION

In conclusion, maternal nutrition plays an important role in ensuring healthy pregnancy, supporting fetal development, and improving early childhood health outcomes. Adequate intake of essential nutrients, proper nutritional assessment, and continuous dietary guidance help prevent deficiencies and promote maternal well-being throughout pregnancy and lactation. Strengthening maternal nutrition services within healthcare systems and communities contributes to healthier mothers, improved birth outcomes, and better developmental prospects for children.

REFERENCES

1. Mimica B, Pribisalic A, Knezovic Z, Sutlovic D. Essential Elements (Fe, Cu, Mn, Zn) in Meconium, and Newborn Length and Weight, in Relation to Maternal Lifestyle and Diet. *Nutrients*. 2025;17:2700.
2. Ichikawa R, Shibata M, Nakura Y, Iizuka K, Uenishi K, Sekiya T, Suzuki A, Nishizawa H. Inadequate Calcium and Vitamin D Intake Among Japanese Women During the Perinatal Period: A Cross-Sectional Study with Bone Health Assessment. *Nutrients*. 2025; 17:1075.
3. Catalano PM, Presley L, Minium J, Mouzon SH De. Fetuses of obese mothers develop insulin resistance in utero. *Diabetes Care*. 2009;32:1076-1080.
4. Rockaway M.M, Daniel AI, Reyes SM, Gauglitz JM, Granger M, McDermid JM, et al. Human milk bioactive components and child growth and body composition in the first 2 years: a systematic review. *Adv Nutr*. 2024;15(1).
5. Dror DK, Allen LH. Overview of nutrients in human milk. *Adv Nutr*. 2018;9:278-294.
6. Allen L H, Hampel D, Shahab-Ferdows S, Andersson M, Barros E, Doel A.M, et al. The Mothers, Infants, and Lactation Quality (MILQ) Study: a multi-center collaboration. *Curr Dev Nutr*. 2021;5(10).
7. Ackerson K, Subramanian SV. Domestic violence and chronic malnutrition among women and children in India. *Am J Epidemiol*. 2008;167(10):1188-1196.
8. Allen LH, Dror DK. Introduction to current knowledge on micronutrients in human milk: adequacy, analysis, and need for research. *Adv Nutr*. 2018;9:275-277.
9. Keats EC, Das JK, Salam RA. Effective interventions to address maternal and child malnutrition: an update of the evidence. *Lancet Child Adolesc*. 2021;5(5):367-384.
10. Senarath U, Agho KE, Akram DeS. Comparisons of complementary feeding indicators and associated factors in children aged 6-23 months across five South Asian countries. *Matern Child Nutr*. 2012;8:89-106.