OPEN ACCESS Freely available online

ISSN: 2155-9546
Journal of Aquaculture Research & Development

Opinion Article

Metabolic Adjustment and Energy Allocation in Aquatic Animals

Hongbin Calabrese*

Department of Aquatic Systems, University of Tasmania, Hobart, Australia

DESCRIPTION

Osmoacclimation energetics describes the changes in energy use that occur when aquatic animals adjust to differences in environmental salinity. Fish, shrimp, crabs, mollusks, and other aquatic organisms constantly regulate the concentration of salts and water within their bodies. When an animal moves between environments with different salinities, its internal fluid balance can be disturbed. Maintaining suitable concentrations of ions and water requires physiological regulation, and this regulation consumes energy. The amount of energy available for growth, movement, reproduction, tissue repair, immunity, and other biological activities can therefore change during salinity adjustment. Understanding these energetic changes is particularly useful in aquaculture because cultured animals may experience changes in salinity caused by water exchange, rainfall, evaporation, seasonal variation, management practices, or movement between production systems.

Aquatic animals living in freshwater, brackish water, and seawater face different osmotic conditions. Freshwater animals generally experience an external environment with a lower concentration of dissolved salts than their body fluids. Water tends to enter the body, while ions can move outward. Marine animals commonly experience the opposite condition, where the surrounding water contains a higher concentration of dissolved salts than their internal fluids. Their physiological systems must limit excessive water loss and manage the uptake and removal of ions. Euryhaline species can tolerate a relatively broad salinity range because their physiological systems can modify their activity according to environmental conditions. These changes require energy, and the energetic cost can vary according to species, developmental stage, body size, temperature, salinity level, rate of environmental change, and nutritional condition.

During the initial phase of salinity change, an aquatic animal may experience a temporary increase in metabolic activity. The animal must detect the new environmental condition and modify ion transport, water balance, endocrine signaling, membrane properties, and tissue function. This adjustment period may require more energy than maintenance under a stable salinity condition. Once physiological regulation becomes more efficient, metabolic expenditure may decline. The length and intensity of this adjustment period differ among species. Some organisms can adapt relatively quickly, whereas others require several days or longer to establish a new physiological balance.

Energy allocation during osmoacclimation is influenced by the availability of food. An animal receiving sufficient dietary energy can allocate nutrients toward both maintenance and physiological adjustment. When food availability is limited, energy may be redirected from growth or reproduction toward ion regulation and cellular maintenance. This can result in slower growth even when the animal survives the salinity change. In aquaculture systems, this relationship is important because feed intake and nutrient utilization can change during environmental transitions. Reduced feeding activity during a salinity change may further limit the energy available for growth.

Temperature also influences osmoacclimation energetics. Warmer water generally increases metabolic activity, although the exact response depends on the species and the temperature range involved. When a salinity change occurs under elevated temperature, the combined physiological demands may increase oxygen consumption and energy expenditure. Conversely, low temperature can reduce metabolic rate and alter the speed of physiological adjustment. Aquaculture managers therefore need to consider salinity and temperature together rather than treating them as completely independent environmental variables.

Body size and developmental stage can also affect energetic responses. Juvenile fish and crustaceans often have different metabolic requirements from adults. Smaller animals may have relatively high metabolic rates per unit of body mass, while larger individuals may show different patterns of energy allocation. Larval stages can be especially sensitive because their osmoregulatory structures and physiological systems are still developing. A sudden salinity change during early development may therefore require greater physiological adjustment than a gradual change applied to an established adult population.

Correspondence to: Hongbin Calabrese, Department of Aquatic Systems, University of Tasmania, Hobart, Australia, E-mail: hongbin.calabrese@utas.edu.au

Received: 29-Jun-2026, Manuscript No. JARD-26-32764; **Editor assigned:** 01-Jul-2026, PreQC No. JARD-26-32764 (PQ); **Reviewed:** 15-Jul-2026, QC No. JARD-26-32764; **Revised:** 22-Jul-2026, Manuscript No. JARD-26-32764 (R); **Published:** 29-Jul-2026, DOI: 10.35248/2155-9546.26.17.1135

Citation: Calabrese H (2026). Metabolic Adjustment and Energy Allocation in Aquatic Animals. J Aquac Res Dev. 17.1135.

Copyright: © 2026 Calabrese H. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Energy expenditure during osmoacclimation is not limited to ion transport. Protein synthesis, membrane remodeling, antioxidant activity, endocrine regulation, and cellular repair can also contribute to metabolic costs. Changes in salinity can alter the function of cellular membranes and influence the movement of water and ions. Animals may respond by changing the abundance or activity of transport proteins and modifying membrane composition. These processes require biochemical resources and therefore contribute to the total energetic cost of adaptation. These adjustments help maintain cellular homeostasis and ensure normal physiological functioning under changing salinity conditions.

CONCLUSION

Osmoacclimation energetics therefore provides an important framework for understanding how aquatic animals balance environmental regulation with biological production. Salinity adjustment requires energy, and the amount required depends on species characteristics, environmental conditions, nutritional status, developmental stage, and the speed of environmental change. When energetic costs become high, fewer resources may remain available for growth and reproduction. Careful management of salinity transitions, temperature, feeding, and water quality can help cultured animals maintain physiological stability while supporting productive performance.