



Microbial Biocatalysis: Harnessing Cellular Chemistry for Sustainable Biochemical Production

Elena Marquez*

Department of Microbial Biotechnology, Federal University of São Paulo, São Paulo, Brazil

DESCRIPTION

Microbial biocatalysis uses microorganisms and their enzymes to convert raw materials into valuable chemical products under relatively mild biological conditions. Bacteria, yeasts, fungi and other microorganisms possess extensive biochemical capabilities that allow them to transform carbohydrates, lipids, proteins, aromatic compounds and numerous organic molecules. These biological catalysts can support the production of pharmaceuticals, food ingredients, biofuels, specialty chemicals, enzymes and industrial intermediates. Compared with several conventional chemical processes, microbial systems can operate at moderate temperatures and pressures while offering considerable control over reaction selectivity [1].

The catalytic capacity of microorganisms originates from their diverse enzyme systems. Enzymes such as oxidoreductases, hydrolases, transferases, lyases and isomerases participate in reactions involving oxidation, reduction, hydrolysis, molecular rearrangement and functional-group modification. A microorganism may contain numerous enzymes that act individually or sequentially within a metabolic pathway. This feature allows microbial cells to perform complex conversions that may require multiple processing stages when carried out through conventional chemical synthesis. Microbial biocatalysis can be performed using whole cells or isolated enzymes. Whole-cell systems are useful when several enzymes are required for a conversion or when intracellular cofactors must be regenerated continuously. Nicotinamide adenine dinucleotide and its phosphate-containing form, Nicotinamide Adenine Dinucleotide (NADH) and Nicotinamide Adenine Dinucleotide Phosphate (NADPH), participate in many oxidation-reduction reactions. Microbial cells can regenerate these cofactors through their normal metabolic activities, reducing the need for repeated external addition. Isolated enzymes, in contrast, provide greater control over reaction conditions and can be incorporated into defined catalytic systems [2,3].

Fermentation technology provides an important platform for producing microbial catalysts. Selected bacterial or fungal strains can be cultivated in controlled nutrient media containing carbon, nitrogen, minerals and other essential components. Parameters such as temperature, pH, dissolved oxygen, agitation and nutrient concentration influence microbial growth and enzyme production. Optimizing these factors can improve catalytic output and increase the efficiency of downstream processing. Modern fermentation systems can operate at laboratory, pilot and industrial scales while maintaining controlled environmental conditions. Microbial biocatalysts are especially valuable for reactions requiring high chemical selectivity. Enzymes can distinguish between structurally related molecules and may selectively act on particular functional groups or stereochemical configurations. This property is important in pharmaceutical manufacturing because different molecular forms of a compound can display different biological activities. Enzymatic transformations can therefore contribute to the preparation of specific intermediates and active pharmaceutical ingredients with reduced formation of unwanted products [4-6].

Hydrolases represent one of the widely utilized groups of microbial enzymes. Lipases, proteases, amylases, cellulases and esterases participate in numerous industrial processes. Lipases can catalyze esterification and transesterification reactions and have applications in food processing, detergent formulation, biodiesel production and organic synthesis. Proteases are used in food processing, leather treatment, detergents and biotechnology. Cellulases assist in the conversion of cellulose-rich materials into soluble sugars, supporting processes associated with biofuel and bioproduct generation. Oxidoreductases also have significant value in microbial biocatalysis. Alcohol dehydrogenases, ketoreductases, oxidases, peroxidases and related enzymes can modify alcohols, ketones, aldehydes and other substrates. These reactions are useful for producing fine chemicals and pharmaceutical intermediates. Microbial sources offer substantial biochemical diversity,

Correspondence to: Elena Marquez, Department of Microbial Biotechnology, Federal University of São Paulo, São Paulo, Brazil, E-mail: elena.marquez@fubio.br

Received: 29-Aug-2025, Manuscript No. JMBT-25-32547; **Editor assigned:** 01-Sept-2025, PreQC No. JMBT-25-32547 (PQ); **Reviewed:** 15-Sept-2025, QC No. JMBT-25-32547; **Revised:** 22-Sept-2025, Manuscript No. JMBT-25-32547 (R); **Published:** 29-Sept-2025, DOI: 10.35248/1948-5948.25.17.654

Citation: Marquez E (2025). Microbial Biocatalysis: Harnessing Cellular Chemistry for Sustainable Biochemical Production. J Microb Biochem Technol. 17:654.

Copyright: © 2025 Marquez E. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

allowing enzymes with different substrate preferences and operational characteristics to be identified [7,8].

Genetic engineering has expanded the ability to improve microbial production systems. Genes encoding useful enzymes can be introduced into suitable host microorganisms, allowing increased enzyme production or modification of catalytic characteristics. Alterations in enzyme structure may influence activity, substrate specificity, temperature tolerance, pH stability and resistance to inhibitors. Metabolic engineering can also redirect cellular resources toward the synthesis of a desired compound by modifying selected biochemical pathways. Immobilization provides another approach for improving enzyme utilization. Enzymes can be attached to or trapped within suitable supporting materials, allowing them to be separated from reaction mixtures and reused. Immobilized microbial enzymes may offer improved operational stability and easier recovery. Such systems are particularly useful in continuous processing, where substrates and products move through a reactor while the catalytic material remains within the processing unit [9,10].

CONCLUSION

Microbial biocatalysis also contributes to environmental applications. Microorganisms can transform certain pollutants, agricultural residues and industrial waste compounds into less harmful or more useful substances. Enzymatic degradation of organic contaminants can support wastewater treatment and bioremediation processes. At the same time, microbial conversion of agricultural residues into sugars, organic acids, enzymes and other valuable products can improve resource utilization. Despite its advantages, microbial biocatalysis requires careful control of biological and processing conditions. Enzyme activity can decrease outside suitable temperature or pH ranges, while substrate toxicity may restrict microbial growth. Product

accumulation can also interfere with cellular metabolism. Process design therefore involves selecting suitable microorganisms, optimizing cultivation conditions, controlling reaction parameters and developing effective product recovery methods

REFERENCES

1. Sheldon RA, Brady D. Broadening the scope of biocatalysis in sustainable organic synthesis. *Chem Sus Chem*. 2019;12:2859-2881.
2. Rossino G, Robescu MS, Licastro E, Tedesco C, Martello I, Maffei L, et al. Biocatalysis: A smart and green tool for the preparation of chiral drugs. *Chirality*. 2022;34:1403-1418.
3. Wu S, Snajdrova R, Moore JC, Baldenius K, Bornscheuer UT. Biocatalysis: Enzymatic synthesis for industrial applications. *Angew Chem Int Ed*. 2021;60:88-119.
4. Bilal M, Iqbal MN. Tailoring multipurpose biocatalysts via protein engineering approaches: A review. *Catal Lett*. 2019;149:2204-2217.
5. Redeker ES, Ta DT, Cortens D, Billen B, Guedens W, Adriaenssens P. Protein engineering for directed immobilization. *Bioconj Chem*. 2013;24:1761-1777.
6. Bolivar JM, Woodley JM, Lafuente FR. Is enzyme immobilization a mature discipline? some critical considerations to capitalize on the benefits of immobilization. *Chem Soc Rev*. 2022;51:6251-6290.
7. Sheldon RA. Enzyme immobilization: The quest for optimum performance. *Adv Synth Catal*. 2007;349:1289-1307.
8. Kumari A, Kaur B, Srivastava R, Sangwan RS. Isolation and immobilization of alkaline protease on mesoporous silica and mesoporous ZSM-5 zeolite materials for improved catalytic properties. *Biochem Biophys Rep*. 2015;2:108-114.
9. Sondhi S, Kaur R, Kaur S, Kaur PS. Immobilization of laccase-ABTS system for the development of a continuous flow packed bed bioreactor for decolorization of textile effluent. *Int J Biol Macromol*. 2018;117:1093-1100.
10. Cen YK, Liu YX, Xue YP, Zheng YG. Immobilization of enzymes on membranes and their applications. *Adv Synth Catal*. 2019;361:5500-5515.