OPEN ACCESS Freely available online

ISSN: 1948-5948
Journal of Microbial & Biochemical Technology

Opinion Article

Microbial Bioremediation: Biological Approaches for Restoring Contaminated Soil and Water

Daniel Pereira*

Department of Environmental Biotechnology, University of Campinas, Campinas, Brazil

DESCRIPTION

Microbial bioremediation is an environmental management approach that uses microorganisms to reduce, transform, immobilize, or remove contaminants from soil, water, sediments and industrial environments. Bacteria, fungi, archaea and selected algae possess diverse metabolic abilities that allow them to interact with petroleum compounds, pesticides, solvents, dyes, heavy metals and other pollutants. Their natural biochemical activities can be used to support the recovery of environments affected by industrial activities, agricultural chemicals, accidental releases and improper waste disposal. Contaminated environments often contain complex mixtures of chemical substances. The behavior of each contaminant depends on its chemical structure, concentration, solubility, temperature, pH, moisture and interaction with soil particles. Microorganisms respond to these conditions through different metabolic pathways. Some pollutants can serve as sources of carbon or energy, while others are transformed as secondary reactions occur during microbial metabolism. The effectiveness of microbial bioremediation therefore depends on both the properties of the contaminant and the biological characteristics of the microorganisms present.

Petroleum contamination is one of the important areas where microbial activity can be useful. Crude oil contains hydrocarbons with different molecular structures, including alkanes, aromatic compounds and larger hydrocarbon molecules. Certain bacteria and fungi can use some of these compounds as carbon and energy sources. During microbial degradation, hydrocarbons may be converted into smaller molecules and, under suitable conditions, eventually into carbon dioxide, water and microbial biomass. The rate of degradation varies according to oxygen availability, nutrient supply, temperature, moisture and the structure of the petroleum compounds. Soil microorganisms are naturally involved in the breakdown of organic pollutants. When a contaminant enters soil, native microbial communities may respond to its presence. Some microorganisms already possess enzymes capable of transformingthe chemical, while others may become active after environmental conditions change. This natural process can be supported through biostimulation, in which nutrients, oxygen, moisture, or other suitable conditions are provided to encourage microbial activity. The objective is to improve the conditions required for naturally occurring microorganisms to function effectively.

Bioaugmentation represents another approach in which selected microorganisms are introduced into a contaminated environment. These organisms may possess specific metabolic abilities that are useful for transforming a particular pollutant. Their successful establishment depends on environmental conditions and interactions with existing microbial communities. Introduced microorganisms may face competition for nutrients, changes in temperature, unsuitable pH, predation, or other factors that influence their survival. Therefore, selecting compatible microorganisms and maintaining suitable environmental conditions are important considerations. Fungi also contribute significantly to bioremediation. Certain filamentous fungi produce extracellular enzymes that can act on complex organic compounds. Enzymes such as laccases, peroxidases and other oxidative enzymes can transform various pollutants, including some dyes, aromatic compounds and industrial chemicals. Fungal hyphae can spread through soil and organic materials, increasing contact with contaminated particles. Agricultural residues and other organic materials may sometimes be used as substrates to support fungal growth during treatment processes.

Industrial wastewater can contain dyes, phenolic compounds, hydrocarbons, solvents and other chemical substances. Microbial treatment systems can reduce selected contaminants before treated water is released or reused. Aerobic microorganisms can metabolize biodegradable organic matter when sufficient oxygen is available. Anaerobic microbial communities can function in oxygen-limited environments and transform organic substances into simpler compounds. Different treatment systems can therefore be selected according to wastewater composition and

Correspondence to: Daniel Pereira, Department of Environmental Biotechnology, University of Campinas, Campinas, Brazil, E-mail: daniel.pereira@microbio.br

Received: 01-Dec-2025, Manuscript No. JMBT-25-32563; **Editor assigned:** 03-Dec-2025, PreQC No. JMBT-25-32563 (PQ); **Reviewed:** 17-Dec-2025, QC No. JMBT-25-32563; **Revised:** 24-Dec-2025, Manuscript No. JMBT-25-32563 (R); **Published:** 31-Dec-2025, DOI: 10.35248/1948-5948.25.17.669

Citation: Pereira D (2025). Microbial Bioremediation: Biological Approaches for Restoring Contaminated Soil and Water. J Microb Biochem Technol. 17:669.

Copyright: © 2025 Pereira D. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

treatment objectives. Heavy metals require a different approach because microorganisms cannot destroy metallic elements. However, microbial cells can interact with metals through adsorption, precipitation, transformation and accumulation. Cell surfaces may contain chemical groups capable of binding metal ions. Some microorganisms can also alter the oxidation state of metals, affecting their mobility and availability. Fungal biomass, bacterial cells and microbial products have therefore been examined as biological materials for removing or immobilizing selected metals from contaminated environments.

Pesticide contamination is another area where microbial processes can contribute to environmental management. Agricultural soils may receive repeated inputs of herbicides, insecticides, fungicides and other crop-protection chemicals. Certain microorganisms possess enzymes capable of transforming pesticide molecules into less complex compounds. The rate and extent of degradation depend on pesticide chemistry, soil characteristics, microbial composition, temperature, moisture and previous exposure to the chemical. Microbial bioremediation can be performed directly at contaminated locations or after contaminated material has been transferred to controlled treatment systems. In situ treatment involves biological activity within the affected soil or groundwater, reducing the need for excavation. Ex situ approaches involve removing contaminated material and treating it under controlled conditions. The selection between these approaches depends on contamination depth, pollutant concentration, site accessibility, soil properties, groundwater movement and treatment objectives.