

## Micronutrient Deficiency in Women Living in Industrialized Countries During the Reproductive Years: Is there a Basis for Supplementation with Multiple Micronutrients?

Ella Schaefer\*

Bayer Consumer Care AG, Basel, Switzerland

### Abstract

Maternal diet is widely recognized to be one of the major environmental factors influencing the development of the embryo and fetus. It is important that women of childbearing age maintain good nutrition throughout their reproductive years, including before conception, as there is a strong relationship between fertility and a successful pregnancy, and a sufficient intake of micronutrients. Yet even in 'high-income', industrialized countries, where dietary resources are more readily available, micronutrient levels in such women may be inadequate.

This review looked at the micronutrient status of women of childbearing age in industrialized countries, as well as those who were pregnant, to determine whether there are any gaps in micronutrient levels. A second objective was to assess whether the evidence indicates a role for multiple micronutrients other than folate and iron during these periods. Results indicated that although some women might have a sufficient intake of micronutrients (although not necessarily all of them), there are those who have lower than the currently-recommended daily intake of micronutrients, particularly folate, vitamin B12, vitamin D, calcium, iodine, iron and selenium. The evidence suggests that multiple micronutrient supplementation during the periconceptional period (i.e. before conception until the end of the first trimester) and throughout pregnancy could help to address inadequate dietary intake of micronutrients, improve maternal status prior to and during pregnancy, and thereby help to minimize reproductive risks.

**Keywords:** Maternal diet; Micronutrient deficiency; Multiple micronutrient supplementation; Periconceptional period; Reproductive health

**Abbreviations:** MMN: Multiple Micronutrients; MTHF: L-5-Methyl-Tetrahydrofolate; NTD: Neural Tube Defects; RBC: Red Blood Cell

### Introduction

It is well established that good nutrition is important and that a balanced diet helps to maintain health in each of us. This is particularly true in women of childbearing age, as there is always the potential of becoming pregnant - and normal development of the embryo and fetus depends on the availability and supply of nutrients from the mother, and thus on her nutritional state [1-4]. In theory, women who are planning to have a child can accommodate the additional requirements through their diet, and a sufficient supply of micronutrients during the periconceptional period (i.e. the period before conception until the end of the first trimester) may support normal fertility, conception, placentation and embryogenesis and ultimately lead to better pregnancy outcomes [2]. However, the diet cannot always supply the micronutrients in the sufficient quantities required. In addition, not all pregnancies are planned; it has been estimated that up to 41% of pregnancies worldwide are unintended [5]. Thus, maintenance of optimal nutritional status throughout the reproductive years is essential.

Yet many women are at risk of insufficient micronutrient intake, and not just in developing countries [6,7]. This can be problematic, as deficiencies in micronutrients can affect fertility and can lead to adverse pregnancy outcomes such as neural tube defects (NTD), low birth weight, preterm deliveries, and pregnancy loss [1,2]. The health of the mother can also be affected by micronutrient deficiency, which contributes to anemia [8] and pre-eclampsia [1] for example. It is very important that women everywhere are made aware of the reality of suboptimal nutrition and the potential problems that can arise-and that

inadequate intake prior to pregnancy usually cannot be rectified once pregnancy has been recognized [3]. Education as to the steps that can be taken to prevent micronutrient deficiency during periconception is vital. The simplest remedy is to consume a varied and healthy diet, which theoretically should be easy to do in more developed countries where women have greater access to good food. However, in reality, the micronutrient needs of many women are not always satisfied. In addition, it should be noted that deficiencies in vitamins and minerals can occur in both undernutrition and overnutrition [9].

Thus, micronutrient supplementation is necessary to address suboptimal nutrition before conception and during pregnancy. To date, periconceptional supplementation has focused on iron and folate, as much evidence supports the increased need for both during pregnancy [10]. There is a growing interest in supplementation with multiple micronutrients (MMN)-but is there a widespread need to supplement with other micronutrients, such as the B vitamins, calcium, vitamin D or iodine, for example? Is there any evidence to suggest that women of reproductive age in more developed countries have a suboptimal intake of such micronutrients? This review aims to offer a perspective of the micronutrient status of women of childbearing potential and pregnant women, to ascertain whether there any benefits associated

\*Corresponding author: Dr. Ella Schaefer, Bayer Consumer Care AG, Basel, Switzerland, Tel: +41 58 272 7522; E-mail: [ella.schaefer@bayer.com](mailto:ella.schaefer@bayer.com)

Received August 18, 2016; Accepted October 17, 2016; Published October 24, 2016

**Citation:** Schaefer E (2016) Micronutrient Deficiency in Women Living in Industrialized Countries During the Reproductive Years: Is there a Basis for Supplementation with Multiple Micronutrients? J Nutr Disorders Ther 6: 199. doi: [10.4172/2161-0509.1000199](https://doi.org/10.4172/2161-0509.1000199)

**Copyright:** © 2016 Schaefer E. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

with supplementation with micronutrients other than folate and iron before conception and during pregnancy.

### Impact of insufficient micronutrients during periconception

The protective effect of folate during the first 28 days of pregnancy is a familiar and accepted concept, and supplementation is proven to significantly reduce the risk of congenital anomalies such as NTD [11-13]. When folate levels are inadequate, the risk of NTD increases [14,15]. Similarly, it has long been recognized that iron deficiency is the major cause of anemia in pregnancy worldwide [8], occurring in more than half of pregnant women in developing countries and almost one-fifth of women in developed countries [8,16]. Even women who enter pregnancy with reasonable iron stores can develop anemia in the later stages of pregnancy [4]. However, less is known about other micronutrients in periconception [1], such as the B vitamins, vitamin D, calcium, selenium and iodine. Table 1 provides a summary of the key adverse effects that are associated with an inadequate intake of these micronutrients during the periconceptual period.

What is increasingly being understood is that adequate nutrition has a role to play at all stages of periconception, not just during pregnancy [2,17]. Various nutrients may alter both maternal and fetal metabolism and thus influence pregnancy outcomes [1,2]. Inadequate

nutrition before conception can negatively affect fertility and may have an adverse effect on implantation [2]. After conception, abnormal placentation is a risk if nutrition is poor [2], which can affect the supply of nutrients to the fetus throughout gestation [1]. Certain nutrients are involved in several biochemical pathways that ultimately affect cell replication and differentiation [18], and thus play a critical role in the development of the brain and nervous system during embryogenesis, while others influence oxidative pathways [1,2]. Abnormalities in placental function and oxidative pathways are associated with adverse effects such as pre-eclampsia, pre-term delivery, and fetal growth restriction [1,2]. Low folate and vitamin B12 levels increase the risk of NTD [2] as well as pregnancy loss [1]. The evidence supports the fact that there is a strong relationship between a sufficient micronutrient intake, good maternal health, and a successful pregnancy outcome [2].

### Recommended micronutrient levels in females

It is clear that periconceptual nutrition is crucial for the successful onset and optimal development of pregnancy, delivery of a healthy child, and maternal health. However, the levels of micronutrients that are necessary during pregnancy are not always apparent. There are sometimes rather wide variations in the levels of micronutrients that are recommended by healthcare agencies and government bodies at different stages of life (Table 2) although there is

Micronutrient	Potential adverse effects of inadequate micronutrient intake	
	Effects on the offspring	Maternal effects
Vitamin A	Premature birth [74] Malformed offspring [2]	Vitamin A-deficiency anemia [2,74] Xerophthalmia [2] Night blindness [2]
Vitamin B6	Neurologic disease of infants [75] Impaired neuromotor development [3] Teratogenicity (omphalocele, exencephaly, cleft palate, micrognathia, digital defects, splenic hyperplasia) [3] Decreased body weight [3] Convulsions [3]	Pre-eclampsia [3,75] Anemia [2] Gestational carbohydrate intolerance [75] Hyperemesis gravidarum [75] Thrombosis [2]
Vitamin B12	Neural tube defects [3] Teratogenicity [3] Reduced fetal growth [75] Neurological abnormalities (cognitive disturbances, dementia, paresthesias, ataxia) [3,75] Fetal demyelination [3] Myelopathy	Anemia Megaloblastic anemia [75]
Folate	Congenital malformations (neural tube defects, orofacial clefts, cardiac anomalies) [3,75] Intrauterine growth retardation Premature birth [3]	Anemia [75] Megaloblastic anemia [3] Spontaneous abortions [75] Pre-eclampsia [75] Placental abruption [75] Thrombosis [75]
Vitamin D	Rickets [4] Impaired dental enamel formation [4] Reduced bone mineral accumulation	Osteomalacia [4,75]
Calcium	Intrauterine growth retardation [75] Low birth weight [75]	Pre-eclampsia [75]
Iron	Oxidative damage to fetal erythrocytes [75] Intrauterine growth retardation [75] Neonates small-for-gestational age/low birth weight [4,75] Premature delivery [4] Cognitive and behavioral problems [4]	Anemia [2,4] Increased corticotropin-releasing hormone and cortisol production [75]
Zinc	Intrauterine growth retardation [4] Teratogenicity [75] Embryonic and fetal death [4] Small-for-gestational age infants [4] Preterm delivery [4]	Gestational hypertension [4] Pre-eclampsia [4] Intrapartum hemorrhage [4]
Iodine	Spontaneous abortions [4] Increased perinatal mortality [4] Mental retardation, cretinism [4]	Thyroid enlargement [4]
Selenium	Intrauterine growth retardation [75]	Recurrent abortions [75] Pre-eclampsia [75]

**Table 1:** Impact of inadequate micronutrient status during preconception or pregnancy.

a general consensus that micronutrient requirements increase during pregnancy. Furthermore, apart from folic acid the importance of other micronutrients is rarely disseminated to women planning a pregnancy, beyond the need for a 'varied and healthy' diet [17]. Even in the US, which offers a comprehensive preconception healthcare plan, the clinical need for a sufficient intake of micronutrients other than folic acid seems not to be overtly communicated directly to women [19]. This is something that should be addressed, and research should enable guidelines to become more consistent as preconception care begins to emerge and develop worldwide.

### An adequate micronutrient intake is not always possible

In industrialized countries, it could be assumed that most women would have easy access to fresh and nutritious foods to maintain a healthy and balanced diet. However, the reality is that women in society are from a diverse range of social, economic, educational, ethnic and cultural backgrounds, all of which have an impact on what they consume. For example, women from low-income families might not always have the money to buy fresh foods, or the means to travel to places that sell them; instead, they may live in areas populated by convenience stores (that often stock only energy-dense but micronutrient-poor processed foods) and fast-food restaurants. Vegetarians and vegans

are often lacking in certain essential micronutrients such as vitamin B12, which is primarily derived from animal products. Certain genetic disorders may prevent digestion or absorption of specific foods, leading to a deficiency in vital micronutrients; for example, lactose intolerance precludes consumption of dairy products, leading to low calcium intake and impaired vitamin D status [20]. An increasingly hectic and stressful lifestyle can result in poor eating habits such as skipping meals and increased consumption of fast foods. Weight loss is ever a concern to many women and the endeavor to reach the perfect weight may result in an adequate consumption of micronutrients. There are numerous factors that may adversely influence a woman's micronutrient status, some of which are listed in Table 3.

### Micronutrient status of women in the general population

Despite the recommendations provided by healthcare and government bodies regarding the importance of consuming a varied and healthy diet in the periconceptional period, micronutrient intake can be inadequate in women of childbearing age [2] and in pregnant women [6]. For example, a high proportion of women of childbearing potential have been shown to have folate levels that are suboptimal (i.e. around 500-600 nmol/l, whereas that thought to be protective of NTD is 906 nmol/l) [21-23]. Similarly, mean iron intake in women aged 25-

Micronutrient	WHO 2004 [10]	IOM 2011 [76]	D-A-CH 2000 [7,77] *	Australia and NZ 2005 [78]	UK Gov 2014 [79]	Russia 2008 [80]	China 2013 [81]	Japan [82]
Vitamin B2 (riboflavin), mg/d	14-18 y: 1.0 19-50 y: 1.1 Pregnancy: 1.4	14-18 y: 1.0 19-50 y: 1.1 Pregnancy: 1.4	19-24 y: 1.5 25-50 y: 1.4	14-50 y: 1.1 Pregnancy: 1.4	15-50 y: 1.1 Pregnancy: 1.4	1.8 Pregnancy (2nd half): 2.0	14-50 y: 1.2 Pregnancy: 1st trimester: 1.2 2nd trimester: 1.4 3rd trimester: 1.5	15-17 y: 1.4 18-49 y: 1.2 Pregnancy: 1.5-1.7†
Vitamin B6 (pyridoxine), mg/d	14-18 y: 1.2 19-50 y: 1.3 Pregnancy: 1.9	14-18 y: 1.2 19-50 y: 1.3 Pregnancy: 1.9	1.5	14-18 y: 1.2 19-50 y: 1.3 Pregnancy: 1.9	15-50 y: 1.2 Pregnancy: 1.2	2.0 Pregnancy (2nd half): 2.3	14-49 y: 1.4 Pregnancy: 2.2	15-17 y: 1.3 18-49 y: 1.2 Pregnancy: 1.4-1.5†
Vitamin B12 (cobalamin), µg/d	14-50 y: 2.4 Pregnancy: 2.6	14-50 y: 2.4 Pregnancy: 2.6	3	14-50 y: 2.4 Pregnancy: 2.6	15-50 y: 1.5 Pregnancy: 1.5	3.0 Pregnancy (2nd half): 3.5	14-50 y: 2.4 Pregnancy: 2.9	15-17 y: 2.5 18-49 y: 2.4 Pregnancy: 2.8-2.9†
Folate, µg/d	14-50 y: 400 Pregnancy: 600	14-50 y: 400 Pregnancy: 600	15-19 y: 300 [83] 20+ y: 400 Pregnancy: 550 [7]	14-50 y: 400 Pregnancy: 600	15-50 y: 200 Pregnancy: 300	400 Pregnancy (2nd half): 600	14-50 y: 400 Pregnancy: 600	15-17 y: 250 18-49 y: 240 Pregnancy: 480-490†
Vitamin D, µg/d	14-50 y: 5 Pregnancy: 5	14-50 y: 15 Pregnancy: 15	5	14-50 y: 5 Pregnancy: 5	Pregnancy: 10[84]	10 Pregnancy (2nd half): 12.5	14-50 y: 10 Pregnancy: 10	15-17 y: 6.0 18-49 y: 5.5 Pregnancy: 12.5-13.0†
Calcium, mg/d	14-18 y: 1,300 19-50 y: 1,000 Pregnancy, 3rd trimester: 1,200	14-18 y: 1,300 19-50 y: 1,000 Pregnancy: 14-18 y: 1,300 19-50 y: 1,000	1,000	14-18 y: 1,300 19-50 y: 1,000 Pregnancy: 14-18 y: 1,300 19-50 y: 1,000	15-18 y: 800 19-50 y: 700 Pregnancy: 700	1000 Pregnancy (2nd half): 1300	14-17 y: 1000 18-50 y: 800 Pregnancy: 1st trimester: 800 2/3 trimester: 1000	15-49 y: 650 Pregnancy: NS
Iron, mg/d	14-17 y: 19.6-65.4 18-50 y: 20.7-58.8 Pregnancy: 7.5-22.6	14-18 y: 15 19-50 y: 18 Pregnancy: 27	10	14-18 y: 15 19-50 y: 18 Pregnancy: 27	15-50 y: 14.8 Pregnancy: 14.8	18 Pregnancy (2nd half): 33	14-17 y: 18 18-49 y: 20 Pregnancy: 1st trimester: 20 2nd trimester: 24 3rd trimester: 29	15-49 y: 10.5‡ Pregnancy: Early stage: +2.5 Mid to late stage: +15.0
Iodine, µg/d	14-50 y: 150 Pregnancy: 200	14-50 y: 150 Pregnancy: 220	19-50 y: 200	14-50 y: 150 Pregnancy: 220	15-50 y: 140 [85] Pregnancy: 140 [85]	150 Pregnancy (2nd half): 220	14-50 y: 120 Pregnancy: 230	15-17 y: 140 18-49 y: 130 Pregnancy: 240-250†
Selenium, µg/d	14-50 y: 26 Pregnancy: 2nd trimester: 28 3rd trimester: 30	14-50 y: 55 Pregnancy: 60	30-70	14-50 y: 60 Pregnancy: 65	15-50 y: 60 [86] Pregnancy: 60 [86]	55 Pregnancy (2nd half): 65	14-50 y: 60 Pregnancy: 65	15-49 y: 25 Pregnancy: 30

\*Values provided as listed in Elmadfa 2009, for non-pregnant women [7]; † depending on age; ‡ menstruating women. NS: Not Stated; RE: Retinol Equivalent

Table 2: Example recommended nutrient intakes of certain micronutrients in females.

Factors		Comments
Social and economic	Poverty/low income [87-89]	May limit access to nutritious foods [90], little dietary diversity [89] May have poor knowledge of nutrition [89] Associated with most of the major risk factors for adverse pregnancy outcomes [87]
	The need to work [87]	Inadequate time for shopping and food preparation [87] Work-related travel alters eating habits (e.g. missed meals) [87] Increased consumption of low-calorie (often nutritionally-poor) diets [87]
	Occupation [90]	Certain jobs are associated with low socioeconomic status (e.g. manual labor, driving), food choice may be limited or irregular [90]
Lifestyle	Alcohol consumption [87]	Excretion of high levels of zinc [87] Altered transport of amino acids across the placenta [87] Overall decrease in maternal nutritional status [87]
	Smoking [87]	Associated with poor eating habits (e.g. skipping breakfast) [87] May alter the metabolism of certain micronutrients [87] Can result in decreased antioxidant status [87]
	Illicit drug use [87]	Can lead to poor eating habits [87]
	Vegetarianism/veganism [91]	Can restrict intake of micronutrients mainly derived from animal products (e.g. vitamin B12) [91]
Health-related	Dieting [87]	May lead to substantially lower than recommended levels of calories, protein, vitamins and minerals [87]
	Health conditions/medication use [87]	For example, chronic digestive diseases can reduce micronutrient absorption, while HIV infection is associated with several nutritional deficiencies [87] Medication use may reduce absorption of micronutrients (e.g. oral contraceptive use is associated with reduced folic acid, B6 and B12 levels, and certain rheumatoid arthritis drugs are folic acid antagonists or interfere with iron absorption) [87]
	Genetic disorders	May prevent digestion or absorption of specific foods, leading to a deficiency in vital micronutrients (e.g. lactose intolerance reduces tolerance of dairy products, leading to low calcium intake and impaired vitamin D status [20])
	Micronutrient/food interactions	Certain foods or micronutrients can inhibit the absorption of other micronutrients; for example, zinc absorption is inhibited by phytate (in staple foods such as cereals, corn and rice), while increasing amounts of zinc in a meal can itself decrease the absorption of zinc [92]
Geographical	'Food deserts' [89]	People in some urban environments (e.g. low-income areas) and rural settings might need to travel to shops that sell more nutritious food, and instead may rely on local convenience stores (generally a poor selection of foods) [89,90] Those with a lack of mobility (e.g. lack of transport) may not have access to nutritious food sources [89]
	Home-grown or farmed produce [90]	Food choice is season dependent, which can limit choice [90] In some regions, the soil may lack certain micronutrients (e.g. selenium in China, [93] iodine deficiency in mountainous or flood-risk regions [94])

**Table 3:** Factors that may adversely affect the nutritional status of women.

49 years has been shown to be one-third lower than the recommended level [24]. Thus, we decided to explore more closely whether such women receive an adequate intake of the micronutrients required during each stage for a healthy pregnancy-whether that pregnancy was planned or not. In general, micronutrient status and deficiency have been assessed in pregnant women from developed countries [25]. But what about the micronutrient status of women in the general population of 'high-income' countries, without pre-existing medical conditions and who presumably have the resources to consume an adequate diet to meet their nutritional needs? Does their micronutrient status reach the recommended levels suggested in the guidelines? If not, how large is the gap? Is there any change in status from those of childbearing potential (including young girls) to those who are already pregnant? To this end, we searched the literature for studies published within the last 10 years that would provide evidence of the micronutrient status of women of childbearing age or those who were pregnant in the general population in high-income countries. Although different methods of data collection (e.g. 24 h dietary recalls, Food Frequency Questionnaires, blood samples) and different age classifications were used in the studies found, the data nevertheless provide an overview of the nutritional status.

In general, as outlined in Supplementary Table 1, it was found that for each micronutrient there are women who at least have an adequate daily intake (within the countries for which data is available)-and those who in fact have a surfeit. It is apparent that these women already consume a nutritionally-sufficient diet and/or may be using micronutrient supplementation (particularly during pregnancy) and/or live in a country where food fortification (with folate and/or iron) is

mandatory. However, more often than not the nutrient intake or serum concentration of most micronutrients is inadequate, as outlined in the summary of Supplementary Table 1.

**Folate:** There was insufficient folate intake across all countries for which data was found, in all of the populations included and across all ages. Where folate was insufficient, intakes ranged from 23% to 98% of the recommended nutrient intake in non-pregnant women [7,26-28] (depending on the reference value used), and between 30% and 95% in pregnant women [7,29-37]. Intake was regularly less than half of the daily amount required [7,27,28,31,33,36,37]. The exception was in non-pregnant women in Japan [34], whose folate levels were apparently sufficient (Table 2); however, the recommended level of folate for such women in Japan is lower than for other countries, apart from the UK (Table 2). In the studies that looked at folate status during several stages of pregnancy, inadequate levels could be seen at each stage (between 33% and 78% of the recommended levels) [31-36].

**Vitamin B2 (riboflavin):** There was some evidence of inadequate intake in non-pregnant [7] and pregnant [7,29] younger women in European countries (where insufficient intakes were between 74% and 92% of the recommended daily intake) and in older pregnant women in Japan [34] (76% and 82% of the recommended intake in the first and second trimester, respectively); otherwise, there was an overconsumption of riboflavin in the women included [7,29,34,38].

**Vitamin B6 (pyridoxine):** On the whole, pyridoxine consumption was sufficient in non-pregnant women (younger and older) [7,38]. However, there was mild insufficiency in non-pregnant women in Japan [34] (73% of the recommended intake) and in some pregnant

women in Austria [7], Canada [32] and Japan [34,35], where inadequate intakes between 65% to 95% of the recommended daily intake were observed.

**Vitamin B12 (cobalamin):** There was some deficiency in cobalamin intake in non-pregnant women (42% [7] and 55% [28] of the recommended intake in Europe), as well as in some pregnant women in Belgium [30], Spain [31] and Australia [33]. In both Spain [31] and Australia [33], intake decreased over the course of the pregnancy, while in Australia the intake increased again after giving birth [33]. There appears to be a surfeit of vitamin B12 in pregnant women in Japan [34,35], where vitamin B12 levels were double the recommended levels, as well as in non-pregnant adolescent girls in Europe [7], non-pregnant women in the USA [38], pregnant adolescent girls in the UK [29], and pregnant women in Canada [32].

**Vitamin D:** Insufficient vitamin D intake was widespread across all women, often to a large extent. In non-pregnant women, insufficient intake ranged between 22% to 88% of the recommended intake [7,38,39], and most intakes accounted for less than one-quarter of the daily amount required [38]. The situation was similar in pregnant women, where insufficient intakes ranged from 21% to 97% of the recommended amount [7,40-78]. However, intakes were more often closer to the recommended nutrient intake (e.g. >75% of the amount required). The greatest insufficiency was observed in China [44-46], Japan [47] and the USA [38,48], but was less apparent in Denmark [40], Sweden [39], New Zealand [42] and Australia [43]. Adolescent girls seemed to be at a high risk of vitamin D deficiency, whether pregnant (UK [29], Austria [7]) or not (Europe [7]). Vitamin D intake appeared to decrease slightly during the course of a pregnancy [40,41].

**Calcium:** Calcium intakes were insufficient in many cases, ranging from 50% to 82% of recommended levels in non-pregnant women [7,38] and 61% to 92% in pregnant women [7,29,49]. Non-pregnant younger girls appeared to be at greatest risk of inadequate intake in Europe [7].

**Iron:** Suboptimal levels of iron were evident in both non-pregnant [7,26,38] and pregnant women [7,29,50,51], according to the varied recommendations set by each country or region. In non-pregnant women, the intakes were particularly low across Europe [7,26], making up just 13% to 68% of the recommended levels with most inadequate intakes below 50%. Even in the US, where flour fortification with iron is mandatory [52], the intake was only three-quarters of the required amount [38]. In pregnant women, iron intake was slightly higher but was still insufficient in many cases [7,29,50,51].

**Iodine:** Inadequate iodine intake could be seen in all women, across all countries. Intakes that were insufficient ranged from 15% to 77% of the required amount in non-pregnant women [7,53,54], and were mostly at the higher end in pregnant women (ranging from 36% to 96%) [7,55-57].

**Selenium:** Selenium intake was mostly sufficient in non-pregnant women [7,38], although a slightly low intake was observed in Australia [26] (around 80% of the recommended intake). In pregnant women in the UK [58], selenium intake was generally adequate although some women had a lower than recommended intake.

Overall, these studies suggest that apart from riboflavin and possibly pyroxidine, there are inadequate nutrient intakes or blood concentrations of most micronutrients in women of childbearing potential, as well as those who are pregnant. In all of these women, a healthy, balanced diet rich in the micronutrients essential for a healthy

pregnancy could ideally prevent such inadequacies. However, as already discussed nutritional intake is highly dependent on a woman's financial situation, social and cultural background, and personal habits [59], which often lead to poor dietary habits and a suboptimal diet. Even women who seemingly consume a sufficient amount of most micronutrients may have a suboptimal intake of one or two essential vitamins or minerals.

### Role of supplementation

Based on the available evidence, it is clear that it is not only deficiencies in iron and folate that need to be addressed in women of childbearing potential in industrialized countries, but that there is also an inadequate intake or blood concentration of multiple micronutrients. Undoubtedly, there is a need for greater education regarding the importance of good nutrition in women of all ages, but particularly during pregnancy. It is vital that women of childbearing potential are fully aware of the importance of a good supply of fresh food, for their own wellbeing as well that of the developing child. Nevertheless, there is a rationale to supplement with multiple micronutrients before conception and during pregnancy, based on the adverse effects that an inadequate nutrient status can produce during this time, both in the developing embryo (e.g. the development of NTD and other congenital abnormalities, decreased birth weight, increased incidence of stillbirth) and in the mother (e.g. increased anemia and pre-eclampsia) [1,2]. Supplementation with multiple micronutrients could improve micronutrient status and thereby allow normal fertility, improve maternal health, and ultimately lead to the healthy development of a baby. However, is there any evidence that MMN supplementation before conception and during pregnancy will provide real clinical advantages?

From a pharmacokinetic standpoint, an MMN supplement that includes folic acid at a daily dose of 800 µg has been evaluated in healthy women of childbearing potential [21]. The study aimed to ascertain at what point red blood cell (RBC) folate levels considered preventive of NTD (906 nmol/l [14]) could be reached. In this study, it was observed that the RBC folate concentration, as a surrogate marker for NTD, increased to a protective level within 4 weeks in more than two-thirds of women, whereas at least 8-12 weeks of daily intake was required at a dose of 400 µg/day as observed in other studies [60-62]. In women who took longer than 4 weeks, a significantly lower baseline folate status compared to the other participants was observed. Furthermore, compared with placebo at 4 weeks and up to 16 weeks, MMN supplementation resulted in significantly higher levels of vitamins B12, B6 and B2, and a significantly lower concentration of homocysteine (where high levels are a risk factor for adverse pregnancy outcomes, including NTD [63]). These results were confirmed in a later study using an MMN supplement containing 400 µg folic acid plus 451 µg L-5-methyl-tetrahydrofolate (MTHF), which also determined its efficacy in achieving NTD-protective RBC folate levels [23]. It was observed that within 4 weeks compared with placebo, the MMN supplement was effective at replenishing RBC folate concentrations to a level considered to prevent NTD. In addition, there was a significant increase in the levels of folate and vitamin B6, while homocysteine concentrations significantly decreased.

Do these increases in RBC folate levels translate to a reduction in NTD and other congenital malformations? A randomized controlled trial using a similar MMN supplement (also containing 800 µg folic acid) in a Hungarian cohort determined that approximately 92% of NTD may be prevented using the MMN supplement, while the total prevalence of congenital abnormalities decreased by around

50% compared to no supplement [64]. Other studies have also demonstrated that MMN supplementation has beneficial effects on pregnancy outcomes. In an analysis of pooled outcomes from 16 high quality trials, there was a significant reduction in the incidence of low birth weight and small-for-gestational age (SGA) births with MMN supplementation, as well as an increase in mean birth weight [65]. This time there was an increased risk of neonatal death with MMN supplementation-but only in the subgroup of studies that began the intervention after the first trimester, again emphasizing the importance of the timing of initiation of MMN supplementation. A recent meta-analysis of MMN supplementation showed that the use of multivitamins rather than folic acid alone during the periconceptional period resulted in a modest but persistent reduction in limb reduction defects and congenital urinary tract abnormalities [66]. In addition, the risk of pre-eclampsia was reduced by 27%, multiple congenital abnormalities by 43%, and occurrent NTD in high-income countries by 49%.

In China, in a study of almost 61,000 women from socioeconomically disadvantaged areas, the incidence of stillbirth, malformation and low birth weight babies was significantly lower with MMN supplementation taken from at least 3 months before pregnancy compared with no nutrient interventions [67]. A meta-analysis of 17 randomized controlled trials in low- to middle-income countries indicated that SGA births were significantly reduced with MMN supplementation compared with standard iron-folate supplements [68]. Overall, there was no significant increase in the risk of neonatal mortality. It was observed that participants remained anemic in the included studies, despite seemingly inadequate amounts of supplemental iron; it was noted that starting the supplementation during pregnancy may be too late for many women, especially those with pre-existing anemia [68]. This highlights the fact that timing of initiation of MMN supplementation is an important aspect that should be considered; supplementation should begin before conception.

There is some concern that excessive periconceptional supplementation with folic acid may modulate DNA methylation, with potential epigenetic effects and long-term health consequences in the child [69-73]. Nevertheless, the number of studies is currently limited and the clinical significance of chronic or high folic acid intake is not well established [72,73]. There is a need for future studies to thoroughly investigate any potential adverse effects, but in the meantime the data is insufficient to impact on the proven benefits of folate supplementation [72,95-98].

## Summary

The evidence suggests that a proportion of women in high-income countries-of childbearing potential or who are already pregnant-have lower either than currently-recommended nutrient intakes or blood concentrations of micronutrients, particularly folate, vitamin B12, vitamin D, calcium, iodine, iron and selenium. All of these women would benefit from supplementation with multiple micronutrients, rather than just folate and iron. Studies have confirmed that MMN supplementation can redress insufficient micronutrient levels and help to improve the health of both the developing embryo (e.g. by reducing the risk of NTD and other congenital abnormalities, increasing birth weight, and reducing the incidence of stillbirth) and that of the mother (e.g. by tackling anemia and pre-eclampsia). Timing of the intervention is critical, as supplementation during pregnancy is often too late to prevent adverse outcomes. To support the optimal reproductive health of women of childbearing potential and pregnant women from

a nutritional perspective, MMN supplements should be taken at least one month before conception and throughout pregnancy.

## Acknowledgements

The draft manuscript was prepared by a professional medical writer (Deborah Nock, Medical WriteAway, Norwich, UK), with full review and approval by the author.

## Author Contributions

ES was responsible for the concept of the review, fully reviewed the manuscript at every stage, and gave final approval.

## Conflicts of Interest

The author is currently a full-time employee of Bayer Consumer Health.

## Declaration of Funding

Funding for the study was provided by Bayer, as were the services of the medical writer, Deborah Nock (Medical WriteAway, Norwich, UK).

## References

1. Ramakrishnan U, Grant F, Goldenberg T, Zongrone A, Martorell R (2012) Effect of women's nutrition before and during early pregnancy on maternal and infant outcomes: a systematic review. *Paediatr Perinat Epidemiol* 26: 285-301.
2. Cetin I, Berti C, Calabrese S (2010) Role of micronutrients in the periconceptional period. *Hum Reprod Update* 16: 80-95.
3. Simpson JL, Bailey LB, Pietrzik K, Shane B, Holzgreve W (2010) Micronutrients and women of reproductive potential: required dietary intake and consequences of dietary deficiency or excess. Part I--Folate, Vitamin B12, Vitamin B6. *J Matern Fetal Neonatal Med* 23: 1323-1343.
4. Simpson JL, Bailey LB, Pietrzik K, Shane B, Holzgreve W (2011) Micronutrients and women of reproductive potential: required dietary intake and consequences of dietary deficiency or excess. Part II--vitamin D, vitamin A, iron, zinc, iodine, essential fatty acids. *J Matern Fetal Neonatal Med* 24: 1-24.
5. Singh S, Sedgh G, Hussain R (2010) Unintended pregnancy: worldwide levels, trends, and outcomes. *Stud Fam Plann* 41: 241-250.
6. Blumfield ML, Hure AJ, Macdonald-Wicks L, Smith R, Collins CE (2013) A systematic review and meta-analysis of micronutrient intakes during pregnancy in developed countries. *Nutr Rev* 71: 118-132.
7. Elmadfa I, Meyer A, Nowak V, Hasenegger V, Putz P, et al. (2009) European Nutrition and Health Report. *Forum Nutr* 62: 1-405.
8. World Health Organization and Food and Agricultural Organization of the United Nations (1998, 2004). Iron. In: Vitamin and mineral requirements in human nutrition 2nd. Report of a joint FAO/WHO expert consultation, Bangkok, Thailand, 21-30 September
9. Young H, Borrel A, Holland D, Salama P (2004) Public nutrition in complex emergencies. *Lancet* 364: 1899-1909.
10. World Health Organization and Food and Agricultural Organization of the United Nations (2004) Vitamin and mineral requirements in human nutrition.
11. Lumley J, Watson L, Watson M, Bower C (2001) Periconceptional supplementation with folate and/or multivitamins for preventing neural tube defects. *Cochrane Database of Systematic Reviews* 4: CD001056.
12. Persad VL, Van den Hof MC, Dubé JM, Zimmer P (2002) Incidence of open neural tube defects in Nova Scotia after folic acid fortification. *CMAJ* 167: 241-245.
13. Pitkin RM (2007) Folate and neural tube defects. *Am J Clin Nutr* 85: 285S-288S.
14. Daly LE, Kirke PN, Molloy A, Weir DG, Scott JM (1995) Folate levels and neural tube defects. Implications for prevention. *JAMA* 274: 1698-1702.
15. Cuskelly GJ, McNulty H, Scott JM (1996) Effect of increasing dietary folate on red-cell folate: implications for prevention of neural tube defects. *Lancet* 347: 657-659.
16. World Health Organization (2001) Iron deficiency anaemia: assessment, prevention, and control. Geneva: WHO/NHD/01.3.
17. Shawe J, Delbaere I, Ekstrand M, Hegaard HK, Larsson M, et al. (2015) Preconception care policy, guidelines, recommendations and services across six European countries: Belgium (Flanders), Denmark, Italy, the Netherlands,

- Sweden and the United Kingdom. *Eur J Contracept Reprod Health Care* 20: 77-87.
18. Cetin I, Alvino G (2009) Intrauterine growth restriction: implications for placental metabolism and transport. A review. *Placenta* 30: S77-82.
19. Advisory Committee on Immunization Practices (2015) Advisory committee on immunization practices recommended immunization schedules for persons aged 0 through 18 years—United States, 2015. *MMWR Morb Mortal Wkly Rep* 64: 93-94.
20. Segal E, Dvorkin L, Lavy A, Rozen GS, Yaniv I, et al. (2003) Bone density in axial and appendicular skeleton in patients with lactose intolerance: influence of calcium intake and vitamin D status. *J Am Coll Nutr* 22: 201-207.
21. Brämswig S, Prinz-Langenohl R, Lamers Y, Tobolski O, Wintergerst E, et al. (2009) Supplementation with a multivitamin containing 800 microg of folic acid shortens the time to reach the preventive red blood cell folate concentration in healthy women. *Int J Vitam Nutr Res* 79: 61-70.
22. Thamm M, Mensink GB, Thierfelder W (1999) Folic acid intake of women in childbearing age. *Gesundheitswesen* 61: S207-212.
23. Schaefer E, Bieri G, Sancak O, Barella L, Maggini S (2016) A randomized, placebo-controlled trial in women of childbearing age to assess the effect of folic acid and methyl-tetrahydrofolate on erythrocyte folate levels. *Vitam Miner* 5: 134.
24. Public Health England and Food Standards Agency (2014) National Diet and Nutrition Survey. Results from Years 1, 2, 3 and 4 (combined) of the Rolling Programme p: 23.
25. Tulchinsky TH (2010) Micronutrient deficiency conditions: global health issues. *Public Health Rev* 32: 243-255.
26. Fayet-Moore F, Petocz P, Samman S (2014) Micronutrient status in female university students: iron, zinc, copper, selenium, vitamin B12 and folate. *Nutrients* 6: 5103-5116.
27. Gall S, Seal J, Taylor R, Dwyer T, Venn A (2012) Folate status and socio-demographic predictors of folate status, among a national cohort of women aged 26-36 in Australia, 2004-2006. *Aust N Z J Public Health* 36: 421-426.
28. Zappacosta B, Persichilli S, Iacoviello L, Di Castelnuovo A, Graziano M, et al. (2013) Folate, vitamin B12 and homocysteine status in an Italian blood donor population. *Nutr Metab Cardiovasc Dis* 23: 473-480.
29. Baker PN, Wheeler SJ, Sanders TA, Thomas JE, Hutchinson CJ, et al. (2009) A prospective study of micronutrient status in adolescent pregnancy. *Am J Clin Nutr* 89: 1114-1124.
30. Jacquemyn Y, Ajaji M, Karepouan N, Jacquemyn N, Van Sande H (2014) Vitamin B12 and folic acid status of term pregnant women and newborns in the Antwerp region, Belgium. *Clin Exp Obstet Gynecol* 41: 141-143.
31. Ubeda N, Reyes L, Gonzalez-Medina A, Alonso-Aperte E, Varela-Moreiras G (2011) Physiologic changes in homocysteine metabolism in pregnancy: A longitudinal study in Spain. *Nutrition* 27: 925-930.
32. Fayyaz F, Wang F, Jacobs RL, O'Connor DL, Bell RC, et al. (2014) Folate, vitamin B12, and vitamin B6 status of a group of high socioeconomic status women in the Alberta Pregnancy Outcomes and Nutrition (APRON) cohort. *Appl Physiol Nutr Metab* 39: 1402-1408.
33. Hure AJ, Collins CE, Smith R (2012) A longitudinal study of maternal folate and vitamin B12 status in pregnancy and postpartum, with the same infant markers at 6 months of age. *Matern Child Health* 16: 792-801.
34. Shibata K, Fukuwatari T, Sasaki S, Sano M, Suzuki K, et al. (2013) Urinary excretion levels of water-soluble vitamins in pregnant and lactating women in Japan. *J Nutr Sci Vitaminol (Tokyo)* 59: 178-186.
35. Takimoto H, Mito N, Umegaki K, Ishiwaki A, Kusama K, et al. (2007) Relationship between dietary folate intakes, maternal plasma total homocysteine and B-vitamins during pregnancy and fetal growth in Japan. *Eur J Nutr* 4: 300-306.
36. Matsuzaki M, Haruna M, Ota E, Sasaki S, Nagai Y, et al. (2008) Dietary folate intake, use of folate supplements, lifestyle factors, and serum folate levels among pregnant women in Tokyo, Japan. *J Obstet Gynaecol Res* 34: 971-979.
37. Yamada T, Morikawa M, Yamada T, Kishi R, Sengoku K, et al. (2013) First-trimester serum folate levels and subsequent risk of abortion and preterm birth among Japanese women with singleton pregnancies. *Arch Gynecol Obstet* 287: 9-14.
38. Rai D, Bird JK, McBurney MI, Chapman-Novakofski KM (2014) Nutritional status as assessed by nutrient intakes and biomarkers among women of childbearing age – is the burden of nutrient inadequacies growing in America?. *Public Health Nutr* 18: 1658-1669.
39. Hedlund L, Brembeck P, Olausson H (2013) Determinants of vitamin D status in fair-skinned women of childbearing age at northern latitudes. *PLoS One* 8: e60864.
40. Milman N, Hvas AM, Bergholt T (2011) Vitamin D status during normal pregnancy and postpartum. A longitudinal study in 141 Danish women. *J Perinat Med* 40: 57-61.
41. Moon RJ, Crozier SR, Dennison EM, Davies JH, Robinson SM, et al. (2015) Tracking of 25-hydroxyvitamin D status during pregnancy: the importance of vitamin D supplementation. *Am J Clin Nutr* 102: 1081-1087.
42. Ekeroma AJ, Camargo CA Jr, Scragg R, Wall C, Stewart A, et al. (2015) Predictors of vitamin D status in pregnant women in New Zealand. *N Z Med J* 128: 24-34.
43. Perampalam S, Ganda K, Chow KA, Opie N, Hickman PE, et al. (2011) Vitamin D status and its predictive factors in pregnancy in 2 Australian populations. *Aust N Z J Obstet Gynaecol* 51: 353-359.
44. Song SJ, Si S, Liu J, Chen X, Zhou L, et al. (2013) Vitamin D status in Chinese pregnant women and their newborns in Beijing and their relationships to birth size. *Public Health Nutr* 16: 687-692.
45. Tao M, Shao H, Gu J, Zhen Z (2012) Vitamin D status of pregnant women in Shanghai, China. *J Matern Fetal Neonatal Med* 25: 237-239.
46. Xiao JP, Zang J, Pei JJ, Xu F, Zhu Y, et al. (2015) Low maternal vitamin D status during the second trimester of pregnancy: a cross-sectional study in Wuxi, China. *PLoS One* 10: e0117748.
47. Shiraiishi M, Haruna M, Matsuzaki M, Murayama R (2014) Demographic and lifestyle factors associated with vitamin D status in pregnant Japanese women. *J Nutr Sci Vitaminol (Tokyo)* 60: 420-428.
48. Eckhardt CL, Gernand AD, Roth DE, Bodnar LM (2015) Maternal vitamin D status and infant anthropometry in a US multi-centre cohort study. *Ann Hum Biol* 42: 215-222.
49. Rocha VS, Lavanda I, Nakano EY, Ruano R, Zugaib M, et al. (2012) Calcium and magnesium status is not impaired in pregnant women. *Nutr Res* 32: 542-546.
50. Vandevijvere S, Amsalkhir S, Van Oyen H, Moreno-Reyes R (2012) Determinants of folate status in pregnant women: results from a national cross-sectional survey in Belgium. *Eur J Clin Nutr* 66: 1172-1177.
51. Mei Z, Cogswell ME, Looker AC, Pfeiffer CM, Cusick SE, et al. (2011) Assessment of iron status in US pregnant women from the National Health and Nutrition Examination Survey (NHANES), 1999-2006. *Am J Clin Nutr* 93: 1312-1320.
52. U.S. Department of Health and Human Services, Food and Drug Administration and Center for Food Safety and Applied Nutrition (November 2015) Questions and Answers on FDA's Fortification Policy. Guidance for Industry.
53. Bath SC, Sleeth ML, McKenna M, Walter A, Taylor A, et al. (2014) Iodine intake and status of UK women of childbearing age recruited at the University of Surrey in the winter. *Br J Nutr* 112: 1715-1723.
54. Watutantrige Fernando S, Cavedon E, Nacamulli D, Pozza D, Ermolao A, et al. (2016) Iodine status from childhood to adulthood in females living in North-East Italy: Iodine deficiency is still an issue. *Eur J Nutr* 55: 335-340.
55. Charlton K, Yeatman H, Lucas C, Axford S, Gemming L, et al. (2012) Poor knowledge and practices related to iodine nutrition during pregnancy and lactation in Australian women: pre- and post-iodine fortification. *Nutrients* 4: 1317-1327.
56. Katz PM, Leung AM, Braverman LE, Pearce EN, Tomlinson G, et al. (2013) Iodine nutrition during pregnancy in Toronto, Canada. *Endocr Pract* 19: 206-211.
57. Mao G, Ding G, Lou X, Zhang R, Zheng P, et al. (2015) Survey of iodine nutritional status in 2011, Zhejiang, China. *Asia Pac J Clin Nutr* 24: 234-244.
58. Rayman MP, Bath SC, Westaway J, Williams P, Mao J, et al. (2015) Selenium status in U.K. pregnant women and its relationship with hypertensive conditions of pregnancy. *Br J Nutr* 113: 249-258.

59. Kotic-Vucinic O, Sulovic N, Radunovic N (2006) Micronutrients in women's reproductive health: II. Minerals and trace elements. *Int J Fertil Womens Med* 51: 116-124.
60. Lamers Y, Prinz-Langenohl R, Brämwig S, Pietrzik K (2006) Red blood cell folate concentrations increase more after supplementation with [6S]-5-methyltetrahydrofolate than with folic acid in women of childbearing age. *Am J Clin Nutr* 84: 156-161.
61. Norsworthy B, Skeaff CM, Adank C, Green TJ (2004) Effects of once-a-week or daily folic acid supplementation on red blood cell folate concentrations in women. *Eur J Clin Nutr* 58: 548-554.
62. Adank C, Green TJ, Skeaff CM, Briars B (2003) Weekly high-dose folic acid supplementation is effective in lowering serum homocysteine concentrations in women. *Ann Nutr Metab* 47: 55-59.
63. Mills JL, Scott JM, Kirke PN, McPartlin JM, Conley MR, et al. (1996) Homocysteine and neural tube defects. *J Nutr* 126: 756S-760S.
64. Czeizel AE (2004) The primary prevention of birth defects: Multivitamins or folic acid?. *Int J Med Sci* 1: 50-61.
65. Ramakrishnan U, Grant FK, Goldenberg T, Bui V, Imdad A, et al. (2012) Effect of multiple micronutrient supplementation on pregnancy and infant outcomes: A systematic review. *Paediatr Perinat Epidemiol* 26: 153-167.
66. Dean SV, Lassi ZS, Imam AM, Bhutta ZA (2014) Preconception care: nutritional risks and interventions. *Reprod Health* 11: S3.
67. Wang YF, Pei LJ, Song XM, Chen G, Zheng XY (2013) Impact of periconceptional multi-micronutrient supplementation on gestation: a population-based study. *Biomed Environ Sci* 26: 23-31.
68. Haider BA, Bhutta ZA (2015) Multiple-micronutrient supplementation for women during pregnancy. *Cochrane Database of Systematic Reviews* 11(Art. No.: CD004905).
69. Sie KK, Li J, Ly A, Sohn KJ, Croxford R, et al. (2013) Effect of maternal and postweaning folic acid supplementation on global and gene-specific DNA methylation in the liver of the rat offspring. *Mol Nutr Food Res* 57: 677-685.
70. Penailillo R, Guajardo A, Llanos M, Hirsch S, Ronco AM (2015) Folic acid supplementation during pregnancy induces sex-specific changes in methylation and expression of placental 11 $\beta$ -hydroxysteroid dehydrogenase 2 in rats. *PLoS ONE* 10: e0121098.
71. Steegers-Theunissen RP, Obermann-Borst SA, Kremer D, Lindemans J, Siebel C, et al. (2009) Periconceptional maternal folic acid use of 400 microg per day is related to increased methylation of the IGF2 gene in the very young child. *PLoS ONE* 4(11): e7845.
72. Burdge GC, Lillycrop KA (2012) Folic acid supplementation in pregnancy: Are there devils in the detail? *Br J Nutr* 108: 1924-1930.
73. Barua S, Kuizon S, Junaid MA (2014) Folic acid supplementation in pregnancy and implications in health and disease. *J Biomed Sci* 21: 77.
74. Radhika MS, Bhaskaram P, Balakrishna N, Ramalakshmi BA, Devi S, et al. (2002) Effects of vitamin A deficiency during pregnancy on maternal and child health. *BJOG* 109: 689-693.
75. Hovdenak N, Haram K (2012) Influence of mineral and vitamin supplements on pregnancy outcome. *Eur J Obstet Gynecol Reprod Biol* 164: 127-132.
76. IOM (Institute of Medicine) (2011) Dietary reference intakes for calcium and vitamin D. The National Academies Press, Washington DC.
77. D-A-CH: German Nutrition Society (DGE), Austrian Nutrition Society (OGE), Swiss Society for Nutrition Research (SGE) and Swiss Nutrition Association (SVE) (2000) Reference values for nutrient intake. Frankfurt/Main, Umschau Braus.
78. Australian Government Ministry of Health, Department of Health and Ageing and National Health and Medical Research Council (2005) Nutrient Reference Values for Australia and New Zealand Including Recommended Dietary Intakes.
79. UK Government and Department for Environment Food & Rural Affairs (2014) Reference nutrient intakes. Guidance: Family Food Methodology.
80. Guidelines 2.3.1.2432-08 (2008) Norms of physiological needs of energy and nutrients for different groups of the population in the Russian Federation.
81. Chinese Nutrition Society (2013) Chinese DRIs handbook.
82. Japan Ministry of Health Labour and Welfare (2015) Overview of dietary reference intakes for Japanese (2015).
83. Krawinkel MB, Strohm D, Weissenborn A, Watzl B, Eichholzer M, et al. (2014) Revised D-A-CH intake recommendations for folate: how much is needed?. *Eur J Clin Nutr* 68: 719-723.
84. Wood CL, Cheetham TD (2016) Vitamin D: increasing supplement use among at-risk groups (NICE guideline PH56). *Arch Dis Child Educ Pract Ed* 101: 43-45.
85. SACN (Scientific Advisory Committee on Nutrition) (2014) SACN statement on iodine and health.
86. SACN (Scientific Advisory Committee on Nutrition) (2013) SACN position statement on selenium and health.
87. Bendich A (1993) Lifestyle and environmental factors that can adversely affect maternal nutritional status and pregnancy outcomes. *Ann N Y Acad Sci* 678: 255-265.
88. Attree P (2005) Low-income mothers, nutrition and health: a systematic review of qualitative evidence. *Matern Child Nutr* 1: 227-240.
89. Biesalski HK (2013) Hidden hunger in the developed world. The road to good nutrition. A global perspective. Karger, Basel, Switzerland.
90. Dindyal S, Dindyal S (2003) How personal factors, including culture and ethnicity, affect the choices and selection of food we make. *Internet J Third World Med* 1: 1-4.
91. World Health Organization and Food and Agricultural Organization of the United Nations (2006) Guidelines on food fortification with micronutrients.
92. Lönnerdal B (2000) Dietary factors influencing zinc absorption. *J Nutr* 130: 1378S-83S.
93. Haug A, Graham RD, Christophersen OA, Lyons GH (2007) How to use the world's scarce selenium resources efficiently to increase the selenium concentration in food. *Microb Ecol Health Dis* 19: 209-228.
94. Biesalski HK (2013) Hidden Hunger. Hidden Hunger. Springer Berlin Heidelberg.
95. World Health Organization (2012) Serum and red blood cell folate concentrations for assessing folate status in populations.
96. Herbert V (1996) Present knowledge in nutrition. 7th, International Life Sciences Institute Press, Washington DC.
97. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, et al. (2011) Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab* 96: 1911-1930.
98. World Health Organization (2013) Urinary iodine concentrations for determining iodine status in populations.