



Multi-Omics Pharmacology: Integrating Genomic, Proteomic, Transcriptomic and Metabolic Information for Drug Response

Sophia Tanaka*

Department of Pharmacological Sciences, Kyoto International University, Kyoto, Japan

DESCRIPTION

Multi-omics pharmacology combines information from several biological measurement platforms to examine how medicines influence cells, tissues and whole organisms. Genomics evaluates Deoxyribonucleic Acid (DNA) variation, transcriptomics examines Ribonucleic Acid (RNA) expression, proteomics measures proteins and their modifications and metabolomics evaluates small molecules produced through cellular activity. When these data are considered together, they can provide a broader description of drug action than any single molecular layer. Pharmacological responses differ among individuals because biological systems vary at multiple levels. A genetic variant may influence the production of an enzyme, while changes in RNA expression can modify the amount of that enzyme. Protein abundance and activity may then affect drug metabolism and metabolic changes can alter the final physiological response. Multi-omics pharmacology connects these stages and helps explain how molecular events contribute to medication effects.

Genomics provides information about inherited and acquired DNA variation. Single-nucleotide variants, insertions, deletions, copy-number changes and structural alterations can affect genes involved in drug metabolism, transport, receptors and signaling pathways. For example, variations in *Cytochrome P450 2D6*, *CYP2C19* and *CYP2C9* can influence the metabolism of selected medicines. Genetic information can therefore provide an initial indication of differences in drug exposure among individuals. Transcriptomics adds information about gene activity. Messenger RNA levels can change after exposure to a medicine, reflecting activation or suppression of specific cellular pathways. Non-coding RNAs, including microRNAs and long non-coding RNAs, can also influence gene regulation. Measuring RNA profiles before and after treatment can help identify biological pathways affected by a drug and provide information about cellular responses.

Proteomics examines the proteins produced within cells or present in biological samples. Proteins are directly involved in enzymatic reactions, receptors, transport, cellular communication, immune responses and structural functions. Drug exposure can change protein abundance, protein localization, or post-translational modifications such as phosphorylation and acetylation. Proteomic measurements can therefore provide information about functional changes occurring after medication exposure. Metabolomics examines small molecules such as amino acids, lipids, sugars, organic acids and other metabolic compounds. These molecules reflect ongoing biochemical activity and can change in response to medicines, disease conditions, diet, or environmental factors. Metabolic profiles can provide information about how a treatment affects energy production, lipid metabolism, amino acid pathways and other cellular processes.

The integration of these molecular layers can provide a more complete picture of drug response. Consider a medicine that acts on a cellular receptor. Genomic analysis may identify a variant affecting the receptor, transcriptomic analysis may indicate altered receptor RNA expression, proteomic analysis may show changes in receptor abundance and metabolomic analysis may reveal downstream biochemical effects. Together, these observations can connect genetic variation with molecular activity and physiological response. Multi-omics pharmacology has important applications in oncology. Tumors contain complex molecular alterations that can differ between patients and even between regions of the same tumor. Genomic information can identify mutations, while transcriptomic data can indicate active genes and signaling pathways. Proteomic measurements can show which proteins are present and modified, while metabolomic information can describe changes in cellular metabolism. Combining these data can assist in characterizing tumor biology and treatment response.

Drug resistance can also be examined through multi-omics approaches. A resistant cell may acquire a genetic alteration,

Correspondence to: Sophia Tanaka, Department of Pharmacological Sciences, Kyoto International University, Kyoto, Japan, E-mail: sophia.tanaka@multiomics.jp

Received: 30-May-2025, Manuscript No. JPP-25-32596; **Editor assigned:** 02-Jun-2025, PreQC No. JPP-25-32596 (PQ); **Reviewed:** 16-Jun-2025, QC No. JPP-25-32596; **Revised:** 23-Jun-2025, Manuscript No. JPP-25-32596 (R); **Published:** 30-Jun-2025, DOI: 10.35248/2153-0645.25.16.141

Citation: Tanaka S (2025). Multi-Omics Pharmacology: Integrating Genomic, Proteomic, Transcriptomic and Metabolic Information for Drug Response. J Pharmacogenom Pharmacoproteomics. 16:141.

Copyright: © 2025 Tanaka S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

change RNA expression, increase production of a drug-efflux protein and modify its metabolic pathways. Examining only one molecular layer may miss important parts of this process. Integrated analysis can connect these changes and provide a broader description of mechanisms associated with reduced drug sensitivity. Multi-omics methods are also useful during pharmaceutical development. Candidate medicines can produce effects across several molecular pathways. Genomic and transcriptomic information can indicate changes in gene activity, while proteomic and metabolomic measurements can provide information about downstream biological consequences. Such data can assist in evaluating drug targets, pharmacological activity, toxicity signals and molecular responses.

Computational analysis is essential for integrating multi-omics datasets. Each molecular platform generates information in a

different format and at a different scale. Bioinformatics tools can normalize datasets, identify relationships between molecular features and organize results into biological pathways. Statistical models and machine-learning methods can also identify combinations of molecular variables associated with treatment outcomes. Data quality is an important consideration in multi-omics pharmacology. Differences in sample collection, storage, laboratory procedures, sequencing platforms and analytical methods can influence results. Appropriate quality control and standardized processing can reduce unwanted technical variation. Biological factors such as age, diet, disease status, medication history and environmental exposure should also be considered when interpreting molecular profiles.