

## Multiplex PCR-based detection of *Salmonella* Typhimurium and *Salmonella* Enteritidis in Specific Pathogen Free (SPF) and Commercial Eggs

Jakee El Jakee<sup>1</sup>, Daa El Din Gad Khelfa<sup>2</sup>, Monier Mohammed El-Safty<sup>3</sup>, Ahmed Adel Seida<sup>1\*</sup>, Sherif Marouf<sup>1</sup>, Jens Hahne<sup>4</sup>, Zafar Mahmood<sup>5</sup> and Sarah Sobhy Nagy<sup>3</sup>

<sup>1</sup>Department of Microbiology and Immunology, Faculty of Veterinary Medicine, Cairo University, Egypt

<sup>2</sup>Department of Poultry and Rabbit Diseases, Faculty of Veterinary Medicine, Cairo University, Egypt

<sup>3</sup>Quality Control of SPF Eggs Department in Central Laboratory for Evaluation of Veterinary Biologics (CLEVB) Abassia, Cairo, Egypt

<sup>4</sup>Institute of Cancer Research, London, England

<sup>5</sup>Graduate School of Life Sciences, University of Würzburg, Würzburg, Germany

\*Corresponding author: Ahmed Adel Seida, Microbiology and Immunology Department, Faculty of Veterinary Medicine, Cairo University, Giza, P.O. 12211, Egypt, Tel: 0020237175004; E-mail: [Seida\\_AA@cu.edu.eg](mailto:Seida_AA@cu.edu.eg)

Received date: February 23, 2016; Accepted date: March 28, 2016; Published date: April 04, 2016

Copyright: © 2016 El Jakee J, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Abstract

*Salmonella* serovars are one of the major bacterial causes of food borne diseases. Eggs are commonly identified as food sources responsible for salmonellosis outbreaks. This study aimed to isolate *Salmonella* Typhimurium and *Salmonella* Enteritidis from 1750 hens' eggs, and use of multiplex polymerase chain reaction (Multiplex PCR) in the identification of different *Salmonella* serovars from eggs. The incidence of salmonellae among the Balady eggs yolk was 1.3%, while the incidence was 1.2% among white and brown eggs samples (each). *S. Typhimurium* and *S. Enteritidis* were identified (0.6 and 0.5% respectively). The isolates were confirmed using *fliC*, *sefA* genes and gene specific for genus *Salmonella*. All albumen samples negative for isolation of salmonellae by culture method were retested by PCR.

From the retested albumen samples 3%, 8.4% and 6% collected from Balady, white and brown eggs respectively were positive for *Salmonella* serovars using Multiplex PCR. No salmonellae could be detected from specific pathogen free (SPF) eggs using both PCR and conventional methods.

**Keywords:** Eggs; SPF; Multiplex PCR; *S. Typhimurium*; *S. Enteritidis*

### Materials and Methods

### Introduction

Contamination of eggs has been identified as one of the major causes of food borne *Salmonella* [1]. In the United States all cases of *Salmonella* contamination of eggs were reported to the Centre for Disease Control and Prevention [2]. There are two pathways for eggs to become internally contaminated with *Salmonella*, direct contamination occurs during the formation of an egg in the ovary and oviduct of hens; whereas, indirect contamination occurs after penetration of salmonellae the egg shell membrane [3].

*Salmonella Pullorum* or *Salmonella Gallinarum* in the ovules before ovulation likely and probably constitutes the chief mode of vertical transmission [4]. The majority of human illnesses caused by *Salmonella* Enteritidis are attributed to the consumption of contaminated eggs [5].

The aim of this study was to determine *S. Typhimurium* and *S. Enteritidis* in eggs of Balady, white, brown and SPF layer breed collected from different governorates using conventional microbiology detection compared to that detected using Multiplex-PCR technique.

### Samples

Total 1750 eggs were collected from Balady (n=1000), brown (n=250), white (n=250) and SPF (n=250) eggs from Kafr El sheikh, Elqalubia, El Monofia, Al Fayoum, EL Menia farms and from the SPF egg producing project (Koom Ousheem-Al Fayoum), Egypt. Egg yolk and egg albumins were collected from each egg and these samples were cultured within 24 hrs from collection.

### Identification of Salmonellae

Under complete sterile condition each egg was cleaned by cotton swab soaked in alcohol. The egg was broken in a Petri dish plate then egg yolk and egg albumin were collected by two separate syringes. Detection of *Salmonella* was carried out according to ISO 6579: (2002) [6]. The samples were cultured on xylose lysine deoxycholate (XLD) and brilliant green agar (BGA) plates. The suspected colonies on XYD and BGA plates were picked up for microscopically examination by Gram's stain before being transferred into semisolid and slope agar for preservation and further identification. *Salmonella* isolates were identified biochemically and serologically as reported in previous literatures [7,8].

### Detection of the genus *Salmonella* using the multiplex PCR-based assay

DNA was extracted from the examined samples using QIAamp® DNA Mini kit (catalog no. 51304, QIAGEN GmbH, Germany) according to manufactures recommendations. The PCR was conducted

according to modified Oliveira protocol using specific primers as shown in table 1 [9]. The primers were prepared by Sigma Company in Germany according to Soumet et al., [10] and amplified PCR products were analyzed gel electrophoresis in 1% agarose gel.

Target sequence	Primer sets	Primer sequence 5'→3'	Amplification region (bp)
Random sequence	ST11	GCCAACCATTGCTAAATTGGCGCA	429
	ST15	GGTAGAAATTCACGCGGTACTGG	
<i>fliC</i> gene	Fli15	CGGTGTTGCCAGGTTGGTAAT	559
	Tym	ACTCTTGCTGGCGGTGCGACTT	
<i>sefA</i> gene	Sef167	AGGTTTCAGGCAGCGGTTACT	312
	Sef478	GGGACATTTAGCGTTTCTTG	

**Table 1:** Primers used for the detection of *Salmonella* species [10].

## Results

### Detection of the *Salmonella* serovars among the examined eggs

The highest number of isolates was recovered from the Balady eggs collected from Kafr El sheik (2.7%) and EL Monofia (3%). The incidence was 1.2% from both the white and brown eggs as shown in table 2. All isolates of salmonellae were recovered from yolk samples

only. The results revealed that 13 serovars (1.3%) were isolated from the yolk samples of Balady eggs and serotyped as 7 *S. Typhimurium* and 6 *S. Enteritidis*. Three serovars (1.2%) were isolated from the yolk of white eggs and serotyped as 2 *S. Typhimurium* and one *S. Enteritidis*. Also 3 serovars (1.2%) were isolated from the yolk of brown eggs and serotyped as 2 *S. Enteritidis* and one *S. Typhimurium* as shown in table 3. It is also clear that all SPF eggs were free from salmonellae infections.

Type of eggs	Number of the examined eggs	Number of <i>Salmonella</i> isolates	Percentage (%)	Governorates
White	250	3	1.20%	Kafr El sheik
	150	2	1.30%	
	100	1	1%	
Brown	250	3	1.20%	Elqalubia
	100	1	1%	
	150	2	1.30%	
Balady	1000	13	1.30%	EL Sharkia
	250	2	0.80%	
	250	2	0.80%	
	250	2	0.80%	
	150	4	2.70%	
	100	3	3%	
SPF	250	-	0%	Al Fayoum
<b>Total</b>	1750	19	1.09%	

**Table 2:** Prevalence of *Salmonella* serovars recovered from the examined eggs.


### Confirmation of the isolates using multiplex PCR

Using primers specific for Genus *Salmonella*, for *S. Enteritidis* (*sefA* gene) and *S. Typhimurium* (*filC* gene) serovars, Multiplex PCR was used to identify the specific isolates. All isolates were positive for

amplification of 429 bp specific for Genus *Salmonella*. Furthermore, *S. Enteritidis* and *S. Typhimurium* isolates were positive for amplification of 312 bp and 559 bp respectively (Figure 1).

Type of eggs	Salmonella serovars				Total	
	S. Enteritidis		S. Typhimurium		No.	%
	No.	%	No.	%		
Balady eggs(1000)	6	0.6	7	0.7	13	1.3
White eggs(250)	1	0.4	2	0.8	3	1.2
Brown eggs (250)	2	0.8	1	0.4	3	1.2
SPF eggs (250)	-	0	-	0	-	0
Total 1750	9	0.5	10	0.6	19	1.3


**Table 3:** *Salmonella* serovars isolated from the egg yolk of the Balady, brown and white eggs.



### Direct detection of the *Salmonella* from egg albumin using the multiplex PCR

All albumen samples collected from the examined eggs were retested by m PCR for detection of *S. Enteritidis* and *S. Typhimurium*. It is clear that 66 (4.4%) out of 1500 albumin samples were positive for salmonellae, 48 (3.2%) and 18 (1.2%) samples were positive for *S. Enteritidis* and *S. Typhimurium* respectively. Among the Balady eggs,

21 (2.1%) and 9 (0.9%) samples were positive for *S. Enteritidis* and *S. Typhimurium* respectively. From the white egg albumin samples, 17 (6.8%) and 4 (1.6%) samples were *S. Enteritidis* and *S. Typhimurium* respectively. The albumin samples of the brown eggs recorded 10 (4%) and 5 (2%) positive samples for *S. Enteritidis* and *S. Typhimurium* respectively as shown in table 4 and figure 2.



Types of egg	Number of examined albumin samples	Amplified PCR product		
		429 bp (Genus <i>Salmonella</i> ) n (%)	312 bp ( <i>S. Enteritidis</i> ) n (%)	559 bp ( <i>S. Typhimurium</i> ) n (%)
Balady	1000	30 (3%)	21 (2.1%)	9 (0.9%)
White	250	21 (8.4%)	17 (6.8%)	4 (1.6%)
Brown	250	15 (6%)	10 (4%)	5 (2%)
Total	1500	66 (4.4%)	48 (3.2%)	18 (1.2%)

**Table 4:** Direct detection of the *Salmonella* from egg albumin samples using the multiplex PCR.

## Discussion

*Salmonella* contamination of eggs has been identified as a public health concern worldwide. Globally, *Salmonella* is one of the most prevalent causes of food borne illness [3].

The present data revealed that 19 *Salmonella* isolates were isolated from 1500 examined Balady, white and brown eggs (1.3%). Earlier *Salmonella* was isolated by Jones et al., [11] from 72.0% of all samples collected from the laying house environment (flush water, ventilation fan, egg belt, and egg collector samples).

It is clear that 10 *S. Typhimurium* and 9 *S. Enteritidis* were identified serologically with incidence of 0.6 and 0.5% respectively. *Salmonella* Enteritidis and *S. Typhimurium* as well as other serotypes have been isolated from egg shells and egg content [12].

The two most commonly identified causative agents of food borne salmonellosis are *Salmonella enterica* serotypes Typhimurium and Enteritidis [13]. Both serotypes have the ability to colonize the reproductive organs of hens and are major causes of food borne illness [3].

*S. Enteritidis* is more commonly linked to contaminated eggs, except in Australia, where the majority of egg-related food borne salmonellosis is caused by *S. Typhimurium* [14-16]. It has been concluded that *S. Enteritidis* could penetrate the egg shell easier than other serotypes so they supposed that horizontal transmission of *Salmonella* in eggs is of less importance than the vertical transmission [17]. Hen's eggs are the most important vehicle of the *S. Enteritidis* infection in humans [18].

The most commonly used technique for *Salmonella* detection is the conventional culture technique. The polymerase chain reaction (PCR) method required only 2 days, compared to the 5 days required by conventional selective enrichment and serological tests for *Salmonella* serovars the culture method and the sensitivity of this assay was approximately less than 1 CFU/600 g of egg pool [19].

A polymerase chain reaction for the specific detection of the gene sequence, *sefA*, encoded by all isolates of *Salmonella* Enteritidis, was developed previously by Woodward and Kirwan [20].

The PCR assay proved by Seo et al., [19] to be a rapid and highly sensitive test for detection of low concentrations of *Salmonella* in egg samples. PCR represents a rapid procedure to detect *Salmonella* in a food sampled. In this study, *sefA* and *filC* genes were amplified to confirm the isolates as well as to detect *S. Enteritidis* and *S. Typhimurium* directly from egg albumen samples. While no albumen sample was detected by the microbiological method, 21 (2.1%) and 9 (0.9%) samples from Balady eggs, 17 (6.8%) and 4 (1.6%) samples from

white egg albumin samples and 10 (4%) and 5 (2%) from brown eggs were positive for *S. Enteritidis* and *S. Typhimurium* respectively. PCR is a sensitive method with a superior ability to detect *Salmonella* spp. in the presence of other competing bacteria [21,22].

All examined albumen samples were negative for isolation of salmonellae by culture method. Egg white proteins, such as lysozyme and ovotransferrin, are well known to play important roles in defense against bacterial invader Baron et al. [23]. Using PCR, 3%, 8.4% and 6% albumen samples collected from Balady, white and brown eggs respectively were positive for *Salmonella* serovars. *Salmonella* DNA could be detected from infertile eggs which incidence was higher than that by bacteria isolation [24]. *Salmonella* strains grow better in fresh egg white than in egg white of 2 or 3 weeks old [17]. In fresh eggs, only few *salmonellas* are present and as albumen is an iron-restricted environment, growth will only occur once storage-related changes to vitelline membrane permeability, which allows salmonellae to invade yolk contents, have taken place, when this happens high populations are achieved in both yolk contents and albumen [25]. In the present study 48 and 18 cases were positive to *S. Enteritidis* and *S. Typhimurium* respectively using m PCR. Baron et al., [23] reviewed critically assesses the available evidence on the antimicrobial components of egg white. In addition, mechanisms employed by *S. Enteritidis* to resist egg white exposure are also considered along with various genetic studies that have shed light upon egg white resistance systems. The egg-contamination capacity of *S. Enteritidis* includes its exceptional survival capability within the harsh conditions provided by egg white [23].

Multiplex PCR is a sensitive method with a superior ability to detect *Salmonella* spp. in the presence of other competing bacteria. Although *Salmonella* contamination of eggs is a complex issue that is influenced by many variables, making it difficult to implement appropriate management strategies. Further research is required to explore different protocols to ensure control of *Salmonella* through temperature and pH of food products. There is also a need to re-educate food handlers and consumers of the risk from raw eggs and cross contamination of food products and reduce the public health risk.

## Conflict of Interest

The authors declare that there is no conflict of interests.

## Acknowledgement

The authors gratefully acknowledge Dr. Zafar Mahmood from Germany and Dr. Jens Hahne from England for their scientific input,

experimental planning and revising the manuscript critically for important intellectual content.

## References

- Howard ZR, O'Bryan CA, Crandall PG, Ricke SC (2012) *Salmonella* Enteritidis in shell eggs: Current issues and prospects for control. Food Res Int 45: 755-764.
- Food and Drug Administration, HHS (2009) Prevention of *Salmonella* Enteritidis in shell eggs during production, storage, and transportation. Final rule. Fed Regist 74: 33029-33101.
- Whiley H, Ross K (2015) *Salmonella* and eggs: from production to plate. Int J Environ Res Public Health 12: 2543-2556.
- Shivaprasad HL (2008) Bacterial Diseases. Diseases of poultry. Black well publishing, USA.
- Gast RK, Guraya R, Jones DR, Anderson KE (2014) Contamination of eggs by *Salmonella* Enteritidis in experimentally infected laying hens housed in conventional or enriched cages. Poult Sci 93: 728-733.
- ISO 6579 (2002) Microbiology of food and animal feeding stuffs. Horizontal method for the detection of *Salmonella* spp. (4th edn.): International Organization for Standardization.
- Quinn PJ, Markey BK, Carter ME, Donnelly WJC, Leonard FC, et al. (2002) Veterinary microbiology and microbial disease. (1stedn.), Blackwell Science Ltd., USA.
- Popoff MY, Bockemühl J, Gheesling LL (2004) Supplement 2002 (no. 46) to the Kauffmann-White scheme. Res Microbiol 155: 568-570.
- Oliveira SD, Santos LR, Schuch DM, Silva AB, Salle CT, et al. (2002) Detection and identification of *salmonellas* from poultry-related samples by PCR. Vet Microbiol 87: 25-35.
- Soumet C, Ermel G, Rose N, Rose V, Drouin P, et al. (1999) Evaluation of a multiplex PCR assay for simultaneous identification of *Salmonella* sp., *Salmonella* Enteritidis and *Salmonella* Typhimurium from environmental swabs of poultry houses. Lett Appl Microbiol 28: 113-117.
- Jones FT, Rives DV, Carey JB (1995) *Salmonella* contamination in commercial eggs and an egg production facility. Poult Sci 74: 753-757.
- Loongyai W, Promphet K, Kangsukul N, Noppa R (2010) Detection of *Salmonella* in Egg Shell and Egg Content from Different Housing Systems for Laying Hens. International Journal of Biological, Biomolecular, Agricultural, Food and Biotechnological Engineering 4: 232-234.
- Galis AM, Marcq C, Marlier D, Portetelle D, Van I, et al. (2013) Control of *Salmonella* contamination of shell eggs-Preharvest and postharvest methods: A review. Compr Rev Food Sci Food Saf 12: 155-182.
- Gantois I, Eeckhaut V, Pasmans F, Haesebrouck F, Ducatelle R, et al. (2008) A comparative study on the pathogenesis of egg contamination by different serotypes of *Salmonella*. Avian Pathol 37: 399-406.
- Wales AD, Davies RH (2011) A critical review of *Salmonella* Typhimurium infection in laying hens. Avian Pathol 40: 429-436.
- Moffatt CRM, Musto J (2013) *Salmonella* and egg-related outbreaks. Microbiol Aust 34: 94-98.
- Dubocage L, Heyndrickx M, Grijspeerdt K, Herman L (2001) Growth of *Salmonella* in egg white. Meded Rijksuniv Gent Fak Landbouwkd Toegep Biol Wet 66: 531-534.
- Gantois I, Ducatelle R, Pasmans F, Haesebrouck F, Gast R, et al. (2009) Mechanisms of egg contamination by *Salmonella* Enteritidis. FEMS Microbiol Rev 33: 718-738.
- Seo KH, Valentin-Bon IE, Brackett RE, Holt PS (2004) Rapid, specific detection of *Salmonella* Enteritidis in pooled eggs by real-time PCR. J Food Prot 67: 864-869.
- Woodward MJ, Kirwan SE (1996) Detection of *Salmonella* Enteritidis in eggs by the polymerase chain reaction. Vet Rec 138: 411-413.
- Rozila A, Hartini AW, Tan DY, Wee SK (2007) Rapid Molecular Detection of *Salmonella* Isolated from Poultry Farm. The 19th Veterinary Association Malaysia Congress, Malaysia.
- Gallegos-Robles MA, Morales-Loredo A, Alvarez-Ojeda G, Osuna-García JA, Martínez IO, et al. (2009) PCR detection and microbiological isolation of *Salmonella* spp. from fresh beef and cantaloupes. J Food Sci 74: M37-40.
- Baron F, Nau F, Guérin-Dubiard C, Bonnassie S, Gautier M, et al. (2016) Egg white versus *Salmonella* Enteritidis! A harsh medium meets a resilient pathogen. Food Microbiol 53: 82-93.
- Tuchili LM, Kodama H, Sharma RN, Takatori I, Pandey GS, et al. (1996) Detection of *Salmonella* DNA in chicken embryos and environmental samples by polymerase chain reaction. J Vet Med Sci 58: 881-884.
- Humphrey TJ (1994) Contamination of egg shell and contents with *Salmonella* Enteritidis: a review. Int J Food Microbiol 21: 31-40.