OPEN ACCESS Freely available online

ISSN: 2155-9546
Journal of Aquaculture Research & Development

Opinion Article

Nutrient Allocation and Lipid Utilization in Cultured Aquatic Animals

Zlatko Greco*

Department of Aquatic Systems, University of Zagreb, Zagreb, Croatia

DESCRIPTION

Aquafeed phospholipid partitioning describes the way dietary phospholipids are digested, absorbed, transported, stored, transformed, and incorporated into tissues in cultured aquatic animals. Phospholipids are important components of cell membranes and participate in lipid transport, cellular signaling, digestive processes, and structural development. Their function in aquafeeds is therefore different from that of neutral lipids, which are mainly associated with energy storage. The proportion, molecular structure, and source of phospholipids in a formulated diet can influence how dietary fatty acids and other lipid-soluble nutrients are distributed throughout the animal.

In fish and crustaceans, dietary phospholipids are exposed to digestive enzymes after feed consumption. Their digestion commonly involves phospholipases that release fatty acids and lysophospholipid products. These products can then enter absorptive cells in the intestine or hepatopancreas, depending on the species. Following absorption, lipid molecules are assembled into transport particles or incorporated into intracellular membranes. The resulting lipid movement determines whether nutrients are directed toward energy production, membrane synthesis, growth-related processes, or longer-term storage.

Phospholipid partitioning is closely connected with the physical properties of the feed itself. Aquafeeds often contain mixtures of marine oils, vegetable oils, animal-derived ingredients, algal materials, and other lipid sources. Each source provides a different combination of phospholipid classes and fatty acids. Soybean lecithin, marine phospholipids, krill-derived ingredients, algal phospholipids, and other lipid-rich materials can therefore produce different metabolic responses even when their total lipid concentrations are similar. Feed formulation needs to consider both lipid quantity and lipid composition.

Phosphatidylcholine is one of the most widely studied phospholipids in aquatic nutrition. It contributes to cellular membranes and can participate in lipid transport from the intestine and liver or hepatopancreas. Phosphatidylethanolamine, phosphatidylinositol, phosphatidylserine, and sphingolipids also have distinct biological functions. The relative abundance of these classes can influence membrane characteristics, cellular communication, and the handling of dietary lipids. Differences among species are important because carnivorous fish, omnivorous fish, marine larvae, freshwater species, and crustaceans can process dietary lipids in different ways.

The digestive stage has a major influence on subsequent phospholipid allocation. Lipid digestion depends on feed particle size, intestinal conditions, bile components, enzyme activity, and the chemical structure of the dietary lipids. Some phospholipid molecules are hydrolyzed before absorption, while others may be absorbed as intact or partially modified molecules. After uptake, fatty acids can be re-esterified into phospholipids or neutral lipids. This creates a dynamic system in which dietary molecules may undergo chemical modification before reaching target tissues.

In crustacean production, the hepatopancreas has an important role in lipid metabolism. Shrimp and other cultured crustaceans can use dietary phospholipids in membrane formation, lipid transport, and developmental processes. During periods of rapid growth or molting, lipid requirements may change because tissue formation and physiological activity alter nutrient allocation. Dietary phospholipid composition can consequently affect lipid deposition in the hepatopancreas, muscle, reproductive tissues, and other organs.

Fish larvae present another important application of phospholipid partitioning. Early developmental stages require nutrients for rapid cellular division, organ formation, nervous-system development, and tissue differentiation. Larval digestive capacity can differ substantially from that of juvenile and adult animals. Phospholipid-rich feeds may therefore have a particular nutritional role during early development. The relationship between phospholipid supply and essential fatty acid delivery is especially relevant because membrane formation requires an appropriate balance of fatty acids within structural lipids.

Correspondence to: Zlatko Greco, Department of Aquatic Systems, University of Zagreb, Zagreb, Croatia, E-mail: zlatko.greco@unizg.hr

Received: 29-Jun-2026, Manuscript No. JARD-26-32765; **Editor assigned:** 01-Jul-2026, PreQC No. JARD-26-32765 (PQ); **Reviewed:** 15-Jul-2026, QC No. JARD-26-32765; **Revised:** 22-Jul-2026, Manuscript No. JARD-26-32765 (R); **Published:** 29-Jul-2026, DOI: 10.35248/2155-9546.26.17.1136

Citation: Greco Z (2026). Nutrient Allocation and Lipid Utilization in Cultured Aquatic Animals. J Aquac Res Dev. 17.1136.

Copyright: © 2026 Greco Z. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The interaction between phospholipids and long-chain polyunsaturated fatty acids is also significant. Eicosapentaenoic acid and docosahexaenoic acid can be incorporated into membrane phospholipids, where they influence membrane fluidity and cellular signaling. Dietary phospholipids may alter the delivery and retention of these fatty acids in tissues. When fish receive diets containing reduced marine ingredients, the form in which essential fatty acids are supplied may affect their incorporation into tissue lipids. Studying phospholipid partitioning can therefore help explain differences in tissue fatty acid profiles among alternative feed formulations.

Phospholipid partitioning is also affected by energy status. When dietary energy is adequate, structural lipids can be directed toward tissue development while excess fatty acids may be stored as triacylglycerols. Under energy limitation, stored lipids can be mobilized to support metabolism. The balance between structural lipid synthesis and energy storage changes according to feeding level, growth rate, temperature, reproductive condition, and environmental stress. Consequently, phospholipid metabolism cannot be considered independently from the animal's overall energy metabolism.

Temperature provides an additional factor in aquatic animals because many fish and crustaceans are ectothermic. Changes in environmental temperature can alter membrane fluidity and influence the fatty acid composition required to maintain suitable cellular function. Animals may adjust the proportion of saturated, monounsaturated, and polyunsaturated fatty acids in membrane phospholipids as environmental conditions change. Feed composition can interact with these physiological adjustments and influence the resulting tissue lipid profile.

Phospholipid partitioning can also be examined through lipidomic analysis. Modern lipid analysis can identify individual lipid classes and molecular species rather than measuring only total lipid concentration. Such measurements allow researchers to compare how specific phospholipid molecules change in the liver, hepatopancreas, muscle, intestine, gills, brain, and reproductive tissues. Combining lipidomic measurements with growth performance, feed utilization, enzyme activity, and gene-expression data can provide a more detailed picture of lipid allocation.

The source of dietary phospholipids is another important consideration. Marine-derived phospholipids may contain long-chain fatty acids and other compounds that differ from those supplied by terrestrial plant sources. Soy-derived phospholipids can provide substantial phosphatidylcholine, while algal ingredients can supply distinctive lipid molecules and fatty acid combinations. The nutritional value of each source depends on species, life stage, inclusion level, processing conditions, and the composition of the complete diet.

Feed processing can modify phospholipid characteristics as well. Extrusion, drying, heating, storage, and exposure to oxygen can affect lipid stability. Oxidized lipids may alter feed quality and can influence animal health when present at excessive levels. Proper ingredient handling and storage are therefore relevant to maintaining the nutritional characteristics of phospholipid-containing aquafeeds.

Understanding phospholipid partitioning also has practical value for reducing dependence on conventional marine ingredients. Alternative lipid sources are increasingly examined in aquaculture because feed manufacturers seek ingredients that can support production while maintaining nutritional quality. However, replacing one lipid source with another can modify the molecular composition of dietary phospholipids. Evaluating how these changes affect digestion, tissue deposition, growth, and physiological function is necessary before broad application.

Research in this area can benefit from controlled feeding experiments that compare phospholipid sources at equivalent lipid and energy levels. Measurements can include feed intake, growth rate, feed conversion, survival, whole-body composition, tissue lipid profiles, digestive enzyme activity, and selected metabolic markers. Sampling at multiple developmental stages can reveal whether phospholipid requirements change as animals mature.

CONCLUSION

Aquafeed phospholipid partitioning is therefore a useful framework for understanding how structural lipids move through the aquatic animal after feeding. It connects feed formulation with digestion, absorption, transport, membrane formation, energy metabolism, and tissue composition. Greater knowledge of these processes can support the design of diets that provide appropriate lipid classes and fatty acids for specific species and production stages. Continued investigation using lipidomics, nutritional physiology, molecular biology, and feeding trials can improve understanding of how dietary phospholipids are allocated within cultured aquatic animals and how feed composition influences this allocation.