



Phenomic Crop Modelling for Advanced Plant Assessment and Agricultural Decision Making

Matthew Debaeke*

Department of Precision Agriculture, Pacific Institute of Agricultural Technology University, Auckland, New Zealand

DESCRIPTION

Phenomic crop modeling is an emerging approach in agricultural research that combines large-scale plant measurements with computational analysis to understand how crops develop under different environmental and management conditions. Plant phenomics involves the systematic measurement of observable plant characteristics, including height, leaf area, biomass, color, architecture, growth rate, flowering time, and reproductive development. When these measurements are integrated into computational models, researchers can examine relationships between plant traits, genetic characteristics, environmental conditions, and agricultural practices. This provides detailed information that can support crop improvement, field management, and production planning.

Crop development is influenced by many interacting factors. Temperature, sunlight, water availability, soil properties, nutrient supply, planting density, and disease pressure can all affect plant growth. Traditional crop assessment methods often depend on visual observation and manual measurements collected at selected stages of development. Although these methods remain useful, they may not capture rapid changes occurring across large populations of plants. Phenomic technologies allow measurements to be collected more frequently and at a much larger scale, creating datasets that can be used for detailed crop modeling.

Digital imaging is an important component of phenomic crop analysis. Cameras mounted on field platforms, drones, robotic systems, or controlled greenhouse structures can capture images of plants at different growth stages. Image-processing software extracts measurable characteristics from these images. Leaf area, plant height, canopy structure, color variation, and growth patterns can be calculated from digital images. Repeated observations create time-series datasets that show how individual plants or groups of plants change throughout the growing season.

Remote sensing expands phenomic observations beyond individual plants. Unmanned aerial vehicles can collect high-resolution images across agricultural fields, while satellites provide repeated observations over much larger areas. Multispectral and hyperspectral sensors capture information across wavelengths that are not visible to the human eye. These measurements can indicate differences in vegetation condition, chlorophyll concentration, canopy development, and water status. When combined with field observations, remote sensing data can improve the spatial coverage of phenomic crop studies.

Phenomic crop modeling also depends on reliable data processing. Large datasets generated by imaging systems may contain thousands or millions of individual measurements. Computational methods are required to organize, clean, classify, and analyze these observations. Image segmentation separates plants from soil or background objects, while feature extraction converts images into numerical variables. Machine learning algorithms can then identify relationships between measured traits and crop performance. Careful data preparation is important because errors in image quality, lighting, sensor calibration, or plant identification can influence model results.

Time-series information is particularly valuable for crop modeling. A single measurement provides information about a plant at one moment, while repeated observations describe its development over time. Growth curves can be generated from measurements of plant height, biomass, canopy area, or vegetation indices. Researchers can compare these curves among cultivars, treatments, environments, and growing seasons. Differences in growth rate may help identify plants that respond efficiently to available resources or maintain development under unfavorable conditions.

Phenomic modeling has significant applications in crop breeding. Breeders evaluate large numbers of plant lines to identify individuals with desirable characteristics. Manual assessment of large populations can require substantial labor and may be affected by differences between observers. Automated

Correspondence to: Matthew Debaeke, Department of Agricultural Data Science, University of Agricultural Sciences, Melbourne, Australia, E-mail: matthew.debaeke@agtech.com

Received: 01-Dec-2025, Manuscript No. AGT-25-32507; **Editor assigned:** 03-Dec-2025, Pre QC No. AGT-25-32507 (PQ); **Reviewed:** 17-Dec-2025, QC No. AGT-25-32507; **Revised:** 23-Dec-2025, Manuscript No. AGT-25-32507 (R); **Published:** 31-Dec-2025, DOI:10.35248/2168-9891.25.14.433

Citation: Debaeke M (2025) Digital Decision Support Systems for Smarter Agricultural Planning and Crop Management. Agrotechnology. 14:433

Copyright: © 2025 Debaeke M. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

phenotyping provides standardized measurements that can be collected repeatedly.

CONCLUSION

The integration of phenomic crop modeling into agricultural research provides a quantitative approach to studying plant performance across different environments. By combining repeated plant measurements with environmental, genetic, and management information, researchers can obtain a more

detailed understanding of crop development. Applications include crop breeding, stress assessment, irrigation planning, nutrient management, disease detection, and yield analysis. Continued development of imaging systems, sensors, data-processing methods, and computational models can increase the efficiency and accuracy of plant assessment. These developments can support better crop research and more informed agricultural management while helping producers and researchers respond to changing production conditions.