OPEN ACCESS Freely available online

ISSN: 2155-9546
Journal of Aquaculture Research & Development

Opinion Article

Pond Ecosystem Modeling for Sustainable Aquaculture Production

Katie Rasgon*

Department of Aquatic Systems, University of Dschang, Dschang, Cameroon

DESCRIPTION

Pond aquaculture is a dynamic production system in which fish, microorganisms, plankton, nutrients, dissolved gases, organic matter, sediment, sunlight, and water movement interact continuously. Understanding these interactions is important for maintaining suitable conditions for aquatic animals and for improving production efficiency. Pond ecosystem modeling provides a structured method for representing these biological, chemical, and physical processes through mathematical relationships, computer simulations, and environmental datasets. Instead of examining individual water-quality measurements separately, a pond model can connect multiple variables and show how changes in one part of the system may influence other components.

A typical aquaculture pond receives inputs from feed, fertilizers, freshwater, rainfall, and atmospheric exchange. Fish and other cultured organisms remove nutrients and oxygen while releasing carbon dioxide, metabolic waste, and organic material. Uneaten feed and fecal material settle toward the bottom, where microorganisms decompose organic matter. During decomposition, oxygen is consumed and nutrients are released into the water and sediment. Phytoplankton use light and dissolved nutrients for photosynthesis, producing oxygen during daylight. At night, respiration by fish, plankton, bacteria, and other organisms continues without photosynthetic oxygen production. These daily cycles can produce substantial variation in dissolved oxygen and carbon dioxide concentrations.

Pond ecosystem models can represent these processes through interconnected variables. Water temperature influences metabolic activity, oxygen solubility, microbial decomposition, and plankton growth. Dissolved oxygen is affected by photosynthesis, respiration, aeration, diffusion from the atmosphere, and oxygen demand from organic material. Ammonia concentration is influenced by feed inputs, fish metabolism, microbial conversion, water temperature, pH, and removal processes. Phosphorus and nitrogen concentrations are also influenced by feed, fertilizers, sediment interactions, microbial activity, and biological uptake. A model can combine these factors to estimate how pond conditions may change over hours, days, or an entire production cycle.

One important application of pond ecosystem modeling is the prediction of dissolved oxygen. Low oxygen can reduce feeding activity, increase physiological stress, slow growth, and increase mortality when severe conditions persist. Oxygen concentration normally changes during the day, with higher values often occurring during periods of active photosynthesis and lower values during nighttime hours. A model can reproduce this daily pattern and estimate periods when oxygen may approach an unsuitable level. Such information can support decisions about aerator operation, feeding schedules, water exchange, and stocking density.

Water temperature is another major variable in pond simulations. Fish have metabolic rates that are strongly influenced by temperature, while microorganisms and phytoplankton also respond to thermal conditions. When temperature rises, oxygen solubility decreases while metabolic demand may increase. This combination can create additional pressure on pond oxygen resources. A temperature-sensitive model can therefore provide more realistic estimates of oxygen demand than a model using a constant temperature value. Seasonal simulations can also help compare expected pond conditions during different periods of the production year.

Phytoplankton dynamics are particularly important because plankton populations influence both oxygen production and water quality. Excessive phytoplankton growth can produce dense blooms, while rapid decline of a bloom may increase microbial decomposition and oxygen consumption. Pond models can represent phytoplankton biomass as a function of light availability, nutrient concentration, temperature, grazing, and loss processes.

CONCLUSION

Pond ecosystem modeling therefore offers a practical framework for studying the interconnected processes that determine aquaculture pond performance. By combining water quality, fish

Correspondence to: Katie Rasgon, Department of Aquatic Systems, University of Dschang, Dschang, Cameroon, E-mail: katie.rasgon@univ-dschang.org

Received: 27-May-2026, Manuscript No. JARD-26-32743; **Editor assigned:** 29-May-2026, PreQC No. JARD-26-32743 (PQ); **Reviewed:** 12-Jun-2026, QC No. JARD-26-32743; **Revised:** 19-Jun-2026, Manuscript No. JARD-26-32743 (R); **Published:** 26-Jun-2026, DOI: 10.35248/2155-9546.26.17.1125

Citation: Rasgon K (2026). Pond Ecosystem Modeling for Sustainable Aquaculture Production. J Aquac Res Dev. 17:1125.

Copyright: © 2026 Rasgon K. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

metabolism, plankton activity, microbial processes, sediment interactions, feed inputs, and environmental conditions, models can provide a broader understanding of pond behavior. Their value depends on appropriate data collection, sound model construction, careful validation, and responsible interpretation. When these requirements are addressed, ecosystem models can become useful tools for improving aquaculture management, resource efficiency, environmental performance, and production planning.