OPEN ACCESS Freely available online

ISSN: 2155-9546
Journal of Aquaculture Research & Development

Commentary

Population Genomics for Genetic Diversity and Sustainable Aquaculture Production

Gernot Classen*

Department of Aquatic Systems, University of Hamburg, Hamburg, Germany

DESCRIPTION

Aquaculture population genomics is an advanced area of aquatic science that examines genetic variation within and among cultured and wild populations of fish, shrimp, mollusks, and other aquatic organisms. Modern aquaculture depends on healthy stocks with suitable growth, survival, reproduction, disease resistance, and environmental tolerance. These characteristics are influenced by genetic variation as well as nutrition, water quality, management, stocking density, temperature, and other production conditions. Population genomics provides a detailed method for examining genetic differences across large numbers of genomic markers, allowing researchers to describe population structure, identify genetic diversity, monitor breeding populations, and support the responsible management of cultured stocks.

Traditional population genetics often relied on a limited number of molecular markers. Techniques such as single nucleotide polymorphism analysis, reduced-representation sequencing, whole-genome sequencing, and genotyping-by-sequencing can provide extensive information about genetic variation. This information can be used to compare hatchery populations, broodstock groups, breeding lines, and natural populations. The resulting genetic profiles can help researchers understand how breeding practices and stock movement influence the genetic composition of aquaculture populations.

Genetic diversity is particularly important for maintaining the long-term performance of cultured aquatic species. A population with sufficient genetic variation has a wider range of inherited characteristics. This variation can support adaptation to changing environmental conditions and reduce the effects associated with excessive inbreeding.

Inbreeding is a significant concern in breeding programs because closely related parents can produce offspring with increased levels of homozygosity. Such genetic conditions may influence growth, fertility, survival, immune performance, and resistance to environmental stress. Genomic data can be used to estimate relatedness among potential broodstock and assist hatchery managers in selecting mating combinations that maintain genetic variation. Instead of depending only on family records, breeders can use genome-wide information to obtain a more detailed assessment of relationships among individuals.

Population genomics also contributes to the identification of population structure. Aquatic species may contain geographically separated populations that have developed different genetic characteristics over long periods. Fish living in rivers, lakes, coastal waters, estuaries, or offshore environments can experience different temperature conditions, salinity levels, food availability, water chemistry, and seasonal patterns. These environmental differences can influence genetic composition. Genomic comparisons can reveal whether populations are closely related or genetically distinct. Such information is useful when selecting broodstock for aquaculture because mixing highly differentiated populations without adequate planning may produce unexpected genetic effects.

Another major application is the identification of genetic loci associated with economically valuable traits. Growth rate, feed utilization, maturation age, disease response, salinity tolerance, temperature tolerance, and flesh quality can have genetic components. Genome-wide association studies can compare genetic variants with measured traits in large groups of cultured animals. When particular genomic regions show consistent associations with desirable characteristics, they can become useful markers for selective breeding. Genomic selection can then use genome-wide marker information to estimate the breeding value of individuals before all production traits are fully expressed.

Disease resistance is another important area for population genomic research. Aquaculture species may encounter bacterial, viral, parasitic, and fungal pathogens during production. The response of individual animals to infection can differ because of genetic variation, immune condition, environmental exposure, and other factors. By comparing the genomes of animals with different disease outcomes, researchers can identify genetic

Correspondence to: Gernot Classen, Department of Aquatic Systems, University of Hamburg, Hamburg, Germany, E-mail: gernot.classen@uni-hamburg.de

Received: 27-May-2026, Manuscript No. JARD-26-32736; **Editor assigned:** 29-May-2026, PreQC No. JARD-26-32736 (PQ); **Reviewed:** 12-Jun-2026, QC No. JARD-26-32736; **Revised:** 19-Jun-2026, Manuscript No. JARD-26-32736 (R); **Published:** 26-Jun-2026, DOI: 10.35248/2155-9546.26.17.1118

Citation: Classen G (2026). Population Genomics for Genetic Diversity and Sustainable Aquaculture Production. J Aquac Res Dev. 17:1118.

Copyright: © 2026 Classen G. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

regions associated with resistance or tolerance. These findings can support breeding programs that select animals with improved health-related characteristics while maintaining sufficient genetic diversity.

CONCLUSION

Population genomics is becoming an increasingly useful component of modern aquaculture breeding and stock management. Its ability to examine genetic variation at a large genomic scale provides information that cannot always be obtained from conventional pedigree records or a small number of molecular markers. Applications range from estimating genetic diversity and relatedness to identifying trait-associated variants, evaluating hatchery effects, supporting disease-resistance breeding, and monitoring cultured and natural populations.