

ISSN: 2473-3350

Vol.23 No.5

# Possibilities to Increase Carbon Sequestration in Forest Ecosystems Through Biochar Soil Application

## Martyna Zywalewska and Maria Sandberg

Department of Engineering and Chemical Sciences, Karlstad University, Sweden


#### Abstract

 $\mathbf{I}_{n}$  the last years, interest of natural fertilizers has increased, their production, form and potential. At the same time from year to year growth harvested annually of forest, at a whole of world. Therefore, more and more research is being conducted which reports on the increasing potential of using biochar in agricultural production systems, as natural fertilizer. Growing problems of environmental protection related to progressive soil degradation, intensifying effects of climate change, energy production and waste management require the search for new, more effective and cheaper solutions. One of the proposed solutions to current problems in the area of environmental protection is biochar, i.e. carbonate obtained in the process of pyrolysis of plant biomass and organic waste. Biochar and its use is not a new solution - it has been used in agriculture for centuries. However, in recent years its properties and potential applications have been rediscovered and it can now be concluded that the traditionally known carbonate, in response to modern needs and applications in the field of environmental protection, has gained a new "brand" and functions as a biochar. Substrates for the production of biochar include a diverse group of materials, which include: energy crops, forest waste, agricultural biomass, sewage sludge, organic fraction of municipal waste or residues from agri-food processing. The choice of substrates depends, among others on physicochemical properties (e.g. water and organic matter content, particle size), potential applications (e.g. for energy production, agricultural purposes, for removing contaminants), logistics aspects and the pyrolysis process and its parameters. Biochar due to such physicochemical properties as high content of organic carbon in stable form and mineral substances, significantly developed porosity and specific surface, can be successfully used: in bioenergetics as a renewable fuel; for carbon sequestration in soil; in the composting process as a structural material or additive limiting ammonia emissions; in the production of organic fertilizers based on biochar; to improve the properties of agricultural land; for removing impurities from aqueous solutions, municipal and industrial wastewater, and process gases; in the remediation of soils contaminated with organic and inorganic compounds, and to reduce pollution of groundwater and surface waters through the retention of e.g. biogenic components in soil.

The use of biochar in environmental protection brings many benefits, including such as the possibility of replacing fossil fuels with a renewable fuel, improving soil properties, e.g. increasing the amount of carbon in the soil or water capacity of

ISSN: 2473-3350

the soil, reducing the consumption of organic and inorganic fertilizers and plant protection products, and thus the risk of



biochar to the environment, analysis of the cost of biochar

production and the availability of substrates useful for its

production, and the cost of using biochar, e.g. for energy

production, remediation of contaminated land, improvement of

soil properties or removal of pollution from municipal and

industrial wastewater.

Biochar is very interesting material which can provide for future forest more substitutes and help them grow faster, bigger and cover good conditions for growing. However, little is known about the impact of biochar, especially in form of hydrochar. This work paper aims to reviewing information and known facts about the effect of the use of biochar in breeding seedlings. Biochars in many paper have been proposed as novel materials for providing soilless growth media. However, is need to remember than much more knowledge is required before reliable advice can be given, to use these material as fertilizers in case of forests. In a production technology of biochar, depends on which process were applied, phytotoxicity and greenhouse gas emissions have been found for certain chars. I studied how these chars affected substrate characteristics, plant performance, water economy and respiratory CO2 emission. However, biochar is much recommended in agriculture production, to help provide higher yields.



ISSN: 2473-3350

Vol.23 No.5



### Biography:

Martyna Zywalewska, is a student of Chemical Sciences, department of Engineering, Karlstad University, Sweden. She worked on 2 projects. Entitled :Evaluation of nutritional pellets with enriched biocarbon from forest industrial residual products, NärSkog and Fosbe - Research Environment For Circular Forest-Based Bioeconomy

15<sup>th</sup> Global Summit and Expo on Biomass and Bioenergy; Rome, Italy- September 21-22, 2020

## **Abstract Citation:**

Martyna Zywalewska, Possibilities to increase carbon sequestration in forest ecosystems through biochar soil application, Biomass 2020, 15<sup>th</sup> Global Summit and Expo on Biomass and Bioenergy; Rome, Italy- September 21-22, 2020

(https://biomass.expertconferences.org/abstract/2020/possibilities-to-increase-carbon-sequestration-in-forest-ecosystems-through-biochar-soil-application)