

Prevalence of Soil Transmitted Helminthes Infections and its Determinants among Primary School Children in Gena Bossa Tiworeda, Ethiopia: Cross-Sectional Study

Teklemariam Ergat Yarinbab^{1*} and Abebe Demissie Darcha²

¹Department of Population and Family Health, Institute of Health, Jimma University, Ethiopia

²Department of Epidemiology, Institute of Health, Jimma University, Ethiopia

*Corresponding authors: Teklemariam Ergat Yarinbab, Department of Population and Family Health, Institute of Health, Jimma University, Ethiopia, Tel: +251996337146; E-mail: teklemariam36@gmail.com

Received date: December 07, 2018; Accepted date: May 21, 2019; Published date: May 31, 2019

Copyright: © 2019 Yarinbab TE, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Soil-Transmitted Helminths (STH) infection is a major public health problem among school-aged children in developing countries. In Ethiopia, school-aged children have been identified as a high-risk group of population to be infected with Soil Transmitted Helminthes. Effective prevention and control of STH infection require the identification of risk factors among high-risk groups. Therefore, the aim of this study was to assess the prevalence of Soil Transmitted Helminthes infections and its determinants among primary school children in Gena Bossa Woreda, Ethiopia.

Methods: Cross-sectional study was conducted in Gena Bossa Woreda, Ethiopia. The data were collected using pre-tested and structured questionnaires. Simple random sampling technique was used. Data analysis was done by SPSS for Windows Version 20.0. Bivariate and Multivariate logistic regressions analyses were conducted. p-value <0.05 was used to declare statistically significant variables.

Result: About 303 (97.7%) of the study subjects participated in the study. Above one-third (38.3%) of the study, participants were tested positive for at least one of the STH species. *lumbricoides* (42.1%) was the predominant parasite followed by Hookworms (37.4%) and *T. trichiura* (11.2%). Lack of private latrines (AOR=4.12, 95% CI: 1.64 and 3.37), not wearing shoes always (AOR=1.80, 95% CI: 1.01, 3.23), age of children (5 to 10 years) (AOR=2.43, 95% CI: 1.42 and 4.16) and not knowing causative agents of STH infection (AOR=2.60, 95% CI: 1.37 and 4.93) were found to be determinants of Soil Transmitted Helminthes Infections.

Conclusion: Lack of private latrines, not wearing shoes always, age of children (5 to 10 years) and not knowing causative agents of STH infections were found to be major determinants of STH Infections. Therefore; public health policymakers and stakeholders working in the area should focus their intervention against STH infections on integrated control programs including regular health education on children's shoes wearing habits, raising awareness of school children and their families on STH infections and promoting private latrines in every household.

Keywords: Determinants; Soil-transmitted helminths; Gena bossa woreda; Ethiopia

Abbreviations: AOR: Adjusted Odds Ratio; SNNPR: Southern Nations Nationalities and Peoples' Region; SPSS: Statistical Package for Social Sciences; STH: Soil Transmitted Helminthes

Background

Soil-Transmitted Helminths (STH) refers to a group of parasites that live in the soil in warm and humid climates and are transmitted to humans by faecally contaminated soil. Major STHs to humans are *Ascaris lumbricoides* (Roundworm), *Trichuris trichiura* (Whipworm), and Hookworm (*Necator americanus* and *Ancylostoma duodenale*) [1,2].

Globally, 2 billion people are infected with soil-transmitted helminths. STH infections are most prevalent in tropical and sub-tropical regions of the developing world where adequate water and

sanitation are lacking [1,3]. In Sub-Saharan Africa, STHs infects over 90 million school children [2,4].

Children are an important high-risk group for soil-transmitted helminths. They are largely affected with STH infection which negatively affects their physical growth, intellectual development, and resistance to other diseases that are contributing significantly to school absenteeism. Furthermore, a large proportion of school children are likely not to be aware of the need for good personal and environmental hygiene and are continuously being exposed to contaminated soil and water [2,5,6].

A national school health and nutrition survey done in 2005-2006 revealed that the overall national prevalence of any soil transmitted helminthes infection was 29.8% with a variable degree of prevalence among regions. Accordingly, SNNPR and Gambella Region contributed the highest prevalence of STH infections i.e. 51% each. Infection of *Ascaris lumbricoides* was the most prevalent (23.2%), followed by Hookworms (9.1%) and *Trichuris trichiura* (7.4%) [7].

There was no prior research conducted in the study area on STH infections. Due to the lack of data about STH infection in the study area, there has not been a deworming program in Gena Bossa Woreda. Therefore, it is important to assess the prevalence of STH infections and its determinants in Gena Bossa Woreda, Ethiopia.

Materials and Methods

Study area and period

The study was conducted in Gena Bossa Woreda from Aug 05/2018 to Aug 28/2018. Gena Bossa Woreda is found in Dawuro Zone, Ethiopia and it is located 578 KM Southwest of Addis Ababa. It has one functional health center and two health posts. There are four governmental primary schools with a total of 3262 school children and 49 teaching sections within a four compound. The total estimated number of population was 17,891 (Projected from the 2015 National Census) (Figure 1).




Figure 1: Location of Karawo town in Gena Bosa woreda, Dawuro zone. Study design: Cross-sectional study was conducted.

Inclusion and exclusion criteria

Inclusion criteria: Primary school children from age 5 to 14 years old were included in the study.

Exclusion criteria: Children who were clinically ill and treated with anti-helminths drugs within the last 2 weeks before the study were excluded from the study.

Sample size determination

The sample size was calculated using a single population proportion formula. The prevalence of STH Infections 27.9% among primary school children of Libo-Kemkem District, Northwest Ethiopia [8], 5% margin of error, 95% confidence level was considered. For the source population was <10,000, the population Correction formula was applied. Finally, a 10% non-response rate was considered. Hence, the total sample size was calculated to be 310.

Data quality control

Questionnaires were prepared in English and translated into the local language (Dawurot'uwa) and vice versa. A pre-test was

conducted on 5% of the sample. Data collectors were given short training on data collection methods. Close supervision was conducted throughout the data collection. Data entry and analysis was conducted carefully.

Data collection tools and procedures

Data were collected using structured questionnaires. Three B.Sc. degree holder public health professionals and three nurses participated in the data collection. Two supervisors participated in the data collection. Short orientation was given to the data collectors and supervisors.

Data processing and analysis

Data were analyzed by SPSS windows version 20. Frequencies and percentages of different variables were computed to summarize the data. A bivariate logistic regression model was fitted as a primary method of analysis. Finally, p-value <0.05 in multivariate analysis was used to declare statistically significant variables.

Ethical approval and consent to participate

The Institutional Review Board (IRB) of Jimma University approved this study. The participants were well informed about the purposes of the study, and oral consents were obtained accordingly. The participants' rights to refuse or withdraw from participating in the study and confidentiality issues were considered.

Definition of terms

- Neglected tropical diseases:** A group of diseases that historically has been overlooked
- Soil-transmitted helminths:** is a term referring to a group of parasitic diseases caused by nematode worms that are transmitted to humans by fecal contaminated soil
- School-age children:** Children aged 5 to 14 years who may or may not be enrolled in School
- School children:** Children aged 5 to 14 years who are enrolled in school
- School-based deworming:** Periodic drug treatment for children in schools based on prevalence to tackle STH infections [2]

Results

Socio-demographic characteristics

Out of 310 sampled primary school children, 303 students participated in the study with the overall response rate of 97.7%. Majority of them, 176 (58.1%), were females and 7 participants were excluded from analysis due to insufficient stool samples obtained.

The children's age ranges from a minimum of 5 years to 14 years with a mean age of 10 (\pm 2.6) years. Two hundred thirty (75.9%) of children's caregivers were females. Among the caregivers, 130 (42.9%) were housewives followed by government workers 51 (16.8%), merchants 41 (13.5%) and private employed 36 (11.9%) (Table 1).

Characteristics (n=303)	Frequency	Percent (%)
Sex of children		
Female	176	58.1
Male	127	41.9

Age of child		
5-10	167	55.1
11-14	136	44.9
Child school grade		
1-4	142	46.9
5-6	90	29.7
7-8	71	23.4
Sex of child caregivers		
Female	230	75.9
Male	73	24.1
The educational level of child caregivers		
Primary education completed	153	50.5
Secondary education completed	91	30
College/university	40	13.2
No education	19	6.3
Occupation of caregivers		
Housewife	130	42.9
Government employee	51	16.8
Merchant	41	13.5
Private/NGO employed	36	11.9
Home servant	19	6.3
Farmer	18	5.9
Others	8	2.7
Annual household income in ETB		
14400 to 28099	104	34.3
>28100	101	33.3
<14400	98	32.3
Family size		
4-6	194	64
s<3	73	24.1
>7	36	11.9

Table 1: Socio-demographic characteristics of school children, Gena Bossa Woreda, Ethiopia, August 2018.

Prevalence of STH infection

One hundred sixteen (38.3%) of the children became positive for at least one of the soil-transmitted helminths species. Besides, *A. lumbricoides* was the predominant parasite (38.8%) followed by *T. trichiura* (34.5%). Female children were more infected with STH infections (Figure 2).




Figure 2: Prevalence of STH among school children, Gena Bossa Woreda, Ethiopia, August 2018.

Factors associated with STH infection

Multivariate logistic regression analysis revealed that lack of private latrines, not wearing shoes always, age of children (5 to 10 years) and not knowing causative agents of STH infections were found to be major determinants of STH Infections.

School children aged 5 to 10 years old were 2.4 times (AOR=2.43, 95% CI=1.42, 4.16) more likely to have STH infections than those aged 11 to 14 years old. Besides, school children who wear shoes sometimes/not at all wear/were nearly two times (AOR=1.80, 95% CI=1.01, 3.23) more likely to be infected with STH than children who wear shoes always.

Furthermore, school children who do not have a private latrine were four times (AOR= 4.12, 95% CI=1.64, 3.37) more likely to have STH infection as compared to those who have private latrines (Table 2).

Variables	STH infection		COR (95% CI)	AOR (95% CI)
	Yes	No		
	No (%)	No (%)		
Child age category				
5-10	75 (44.9)	92 (55.1)	1.88 (1.17-3.04)*	2.43 (1.42-4.16)**
11-14	41 (30.1)	95 (69.9)	1	1
The educational level of child caregivers				
No education	9 (47.4)	10 (52.6)	2.10 (0.68-6.47)	0.94 (0.25-3.54)
Primary education completed	67 (43.8)	86 (56.2)	1.81 (0.86-3.84)	1.00 (0.42-2.40)
Secondary education completed	28 (30.8)	63 (69.2)	1.03 (0.46-2.33)	0.75 (0.30-1.85)
College/ university	12 (30.0)	28 (70.0)	1	1
Annual household income in ETB				
<14400	39 (39.8)	59 (60.2)	1.64 (0.90-2.96)	1.48 (0.74-2.97)

14400 to 28099	48 (46.2)	56 (53.8)	2.12 (1.19-3.79)	1.75 (0.90-3.38)
>28100	29 (28.7)	72 (71.3)	1	1
Family size				
<3	33 (45.2)	40 (54.8)	1	1
4-6	63 (32.5)	131 (67.5)	0.58 (0.33-1.01)	0.70 (0.37-1.31)
>7	20 (55.6)	16 (44.4)	1.51 (0.67-3.38)	1.69 (0.69-4.16)
Availability of private latrine				
Yes	99 (35.9)	177 (64.1)	1	1
No	17 (63.0)	10 (37.0)	3.09 (1.34-6.89)*	4.12 (1.64-3.37)**
Child shoes wearing the habit				
Always	55 (29.9)	129 (70.1)	1	1
Sometimes/not at all	61 (51.3)	58 (48.7)	2.46 (1.52-3.98)*	1.80 (1.01-3.23)**
A short and clean finger				
Yes	70 (34.3)	134 (65.7)	1	1
No	46 (46.5)	53 (53.5)	1.66 (1.01-2.71)	1.17 (0.65-2.11)
Hear about STH				
Yes	89 (76.3)	156 (63.7)	1	1
No	27 (46.6)	31 (53.4)	1.52 (0.85-2.72)	0.71 (0.35-1.45)
Know causative agents of STH infection				
Yes	66 (30.8)	148 (69.2)	1	1
No	50 (56.2)	39 (43.8)	2.87 (1.72-4.78)*	2.60 (1.37-4.93)**
**Significant at p-value <0.05				

Table 2: Factors associated with STH infection among school children, Gena Bossa Woreda, Ethiopia, Aug 2018.

Discussion

The current study revealed that the prevalence of STH infections among school children was 38.3%. This finding was larger as compared to the study findings from Libo-Kemkem District, Northwest Ethiopia [9] and Umolante District, SNNPR Ethiopia [8]. In contrast, the overall prevalence in the current study was lower than the findings of studies conducted in Jimma Zone, Ethiopia [10] and in Nigeria [11]. The differences among the study findings might be due to differences in living standards and socio-demographic characteristics of the populations.

School children whose family did not have private latrine were four times more likely to be infected by STH infections when compared to those whose families possess private latrines. This finding was in agreement with findings of other studies in Jimma Zone, Ethiopia [12]. Here, the absence of latrines in both setups might have exposed school children to parasites because of the existing improper defecation system [13].

Besides, children who did not wear shoes always were two times more likely to be infected with STH than their counterparts [14]. This finding is similar to the study finding from Motta Town, Ethiopia [10]. The possible reason may be the similarity of the existing socio-economic status of the population in the two settings.

Moreover, children whose caregivers did not know the causative agents for STH infections were about three times more likely to be infected with STH infection than those whose families were aware of the same. This finding was similar to the study findings from Nigeria and Ethiopia [4,10]. This implies that knowledge about the agents and transmission mechanisms of the STH infection may contribute to the prevention of STH infection [15].

Conclusion

The study revealed that lack of private latrines, not wearing shoes always, age of children (5 to 10 years) and not knowing causative agents of STH infections were found to be major determinants of STH Infections.

Recommendation

Ethiopian Federal Ministry of Health in collaboration with other stakeholders should focus their intervention against STH infections on integrated control programs including regular health education on children's shoes wearing habits, raising awareness of school children and their families on STH infections and promoting private latrines in every household. Dawro Zonal Health Department and Gena Bossa District Health Office should also give due attention to accessibility and proper utilization of private latrines.

Declarations

Ethical approval

Ethical clearance letter was received from Jimma University. Parents were asked to sign the assent forms if they agreed on their children to be involved in the study. Permission to collect a stool sample from pupils was sought from the head teachers of the schools involved in the study and informed verbal consent was obtained from the pupils themselves. Confidentiality issues were considered. School children who were tested positive for STH infection were treated.

- **Consent to publish:** Not applicable
- **Availability of data and materials:** All data included in the article
- **Competing interests:** Authors' declare that they have no competing interests
- **Funding:** No funding was received for this study

Authors' Contribution

AD developed the proposal, supervised the data collection and conducted the data analysis. TE participated in data analysis, conducted the report writing and drafted the manuscript. Both authors critically reviewed and approved the final manuscript.

Acknowledgment

First of all, we thank Jimma University for supporting this research activity by supplying all the necessary materials. We also would like to

thank the Gena Bossa Woreda Office of Education and Schools for their good collaboration. Finally, our great appreciation goes to the data collectors and supervisors.

References

1. PAHO (2011) Addressing soil-transmitted helminths in Latin America and the Caribbean.
2. WHO (2012) Eliminating soil-transmitted helminthiasis as a public health problem in children and progress report 2001-2010 and Strategic Plan 2011-2020, Geneva, Switzerland.
3. PAHO/WHO (2015) Soil-transmitted helminthiasis, key facts on neglected infectious diseases.
4. Babatunde SK, Adedayo MR, Ajiboye AE, Sunday O, Ameen N (2013) Soil-transmitted helminth infections among school children in rural communities of moro local government area, kwara state, Nigeria. *Afr J Microbiol Res* 7: 5148-5153.
5. Munisi DZ, Kihamia CM, Jones C, Msoffe PL (2014) Soil-transmitted helminthes infections, malnutrition and anaemia among primary school children in northern Tanzania. *IMTU Med J* 5: 15-21.
6. WHO (2011) Weekly epidemiological record of NTDs.
7. Jemaneh L (2001) Soil-transmitted helminthes infections and schistosomiasis mansoni in school children from Chilga district, Northwest Ethiopia. *Ethiop J Health Sci* 11: 79-87.
8. Addisu T (2015) A survey of soil-transmitted helminths infections and schistosomiasis mansoni among school children in Libo-Kemkem district, northwest Ethiopia: Cross-sectional study. *Am J Health Res* 3: 57.
9. Alemu M (2014) Prevalence of intestinal schistosomiasis and soil-transmitted helminthiasis among primary school children in Umolante district, South Ethiopia. *Clin Med Res* 3: 174.
10. Asemahagn MA (2014) Parasitic infection and associated factors among the primary school children in Motta town, Western Amhara, Ethiopia. *Am J Public Health Res* 2: 248-254.
11. Egwunyenga OA, Ataikiru DP (2005) Soil-transmitted helminthiasis among school-age children in ethiope East local government area, delta state, Nigeria. *Afr J Biotechnol* 4.
12. Emanu D (2015) Prevalence and intensity of soil-transmitted helminths among school-aged children in sigmo primary school, Jimma zone, south-western Ethiopia. *Clin Med Res* 4: 98-103.
13. Mekonnen Z, Getachew M, Bogers J, Vercruysse J, Levecke B (2019) Assessment of seasonality in soil-transmitted helminth infections across 14 schools in Jimma Town, Ethiopia. *Pan Afr Med J* 4: 32-36.
14. Chard AN, Baker KK, Tsai K, Levy K, Sistrunk JR, et al. (2019) Associations between soil-transmitted helminthiasis and viral, bacterial, and protozoal enteroinfections: A cross-sectional study in rural Laos. *Parasit Vectors* 12: 216.
15. Ercumen A, Benjamin-Chung A, Arnold BF, Lin A, Hubbard AE, et al. (2019) Effects of water, sanitation, handwashing and nutritional interventions on soil-transmitted helminth infections in young children: A cluster-randomized controlled trial in rural Bangladesh. *PLoS Negl Trop Dis* 13: e0007323.