OPEN ACCESS Freely available online

ISSN: 2155-9546
Journal of Aquaculture Research & Development

Short Communication

Probiotics in Fish Culture: Role in Growth, Immunity, and Disease Management

Carmela Venter*

Department of Marine Biology and Fisheries, University of Padua, Padua, Italy

DESCRIPTION

Probiotics are living microorganisms that can provide beneficial effects to aquatic animals when supplied in suitable amounts. In modern aquaculture, probiotics have gained considerable attention because intensive fish culture can create conditions that favor the accumulation of organic wastes and the growth of harmful microorganisms. High stocking densities, poor water quality, nutritional deficiencies, and environmental stress can reduce fish health and increase susceptibility to disease. The application of probiotics offers a biological approach to improving water quality, digestion, immunity, and overall fish performance.

Probiotic microorganisms used in aquaculture commonly include beneficial species of bacteria such as *Bacillus, Lactobacillus, Lacto coccus*, and certain other microorganisms. These organisms may be administered through feed or directly into the culture water. Their effectiveness depends on the microbial strain, dosage, method of application, fish species, developmental stage, environmental conditions, and duration of treatment. Therefore, the selection of an appropriate probiotic is an important consideration in aquaculture management [1].

One of the major functions of probiotics is the improvement of digestive processes. Beneficial microorganisms can produce enzymes such as proteases, lipases, and amylases that assist in the breakdown of dietary nutrients. Improved digestion may increase the availability of proteins, carbohydrates, and lipids for absorption. As a result, probiotic supplementation may contribute to better feed utilization, growth rate, and feed conversion efficiency in cultured fish [2].

Probiotics can also influence the microbial population within the gastrointestinal tract. The digestive system of fish contains a diverse community of microorganisms that can affect digestion, metabolism, and health. Beneficial bacteria may compete with harmful microorganisms for nutrients and attachment sites within the intestine. By influencing the composition of the intestinal microbiota, probiotics can help create conditions that are less favorable for certain opportunistic pathogens [3].

Another important function of probiotics is the production of antimicrobial substances. Some beneficial bacteria produce organic acids, bacteriocins, enzymes, and other compounds that can inhibit the growth of harmful microorganisms. These substances may interfere with pathogen attachment, reproduction, or survival. Consequently, probiotics can contribute to biological protection against certain bacterial diseases without relying exclusively on chemical treatments or antibiotics [4].

The immune system of fish can also be influenced by probiotic supplementation. Fish possess both innate and adaptive immune mechanisms, with innate immunity providing an important first line of defense against pathogens. Probiotics may stimulate immune-related cells and increase the activity of defense mechanisms such as phagocytosis, lysozyme activity, complement activity, and other immune responses. Enhanced immune function may improve the ability of fish to resist infections and recover from environmental stress [5].

Probiotics may also improve resistance to disease by strengthening the intestinal barrier. The intestinal surface acts as an important protective boundary between the internal tissues of fish and the external aquatic environment. Beneficial microorganisms can influence mucus production and epithelial integrity, thereby supporting the physical and biological defenses of the gastrointestinal tract. A healthy intestinal environment is particularly important in intensive culture systems where fish are continuously exposed to microorganisms [6].

Water quality management is another important application of probiotics. Certain bacterial strains can participate in the decomposition of organic matter and the transformation of nitrogenous compounds. Uneaten feed, feces, and other organic materials can increase ammonia and other waste products in aquaculture systems. Beneficial microbial communities can assist in the biological processing of these substances, potentially

Correspondence to: Carmela Venter, Department of Marine Biology and Fisheries, University of Padua, Padua, Italy, Email: carmela.venter@unipd.it

Received: 17-Aug-2026, Manuscript No. JARD-26-32974; **Editor assigned:** 20-Aug-2026, PreQC No. JARD-26-32974 (PQ); **Reviewed:** 03-Sep-2026, QC No. JARD-26-32974; **Revised:** 10-Sep-2026, Manuscript No. JARD-26-32974 (R); **Published:** 17-Sep-2026, DOI: 10.35248/2155-9546.26.17.1146

Citation: Venter C (2026). Probiotics in Fish Culture: Role in Growth, Immunity, and Disease Management. J Aquac Res Dev. 17.1146.

Copyright: © 2026 Venter C. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

improving water quality when combined with appropriate aeration, filtration, and routine management [7].

The reduction of ammonia is particularly important because elevated concentrations can cause physiological stress and damage to fish. Ammonia can interfere with gill function, metabolism, and osmoregulation. Some probiotic and bioaugmentation systems promote microbial processes that transform ammonia into less harmful nitrogen forms. However, probiotics should not be considered a replacement for proper water-quality management, since filtration, aeration, stocking density, feeding practices, and water exchange remain important [8].

Probiotics may also contribute to improved growth performance. Fish receiving suitable probiotic supplementation may show increased weight gain, improved specific growth rate, and better feed conversion. These effects may result from improved digestion, enhanced nutrient absorption, healthier intestinal microbiota, and reduced energy expenditure associated with disease and physiological stress. However, positive effects are not universal, and different probiotic strains may produce different results [9].

Stress is a significant concern in aquaculture because fish may experience stress during transportation, handling, grading, stocking, changes in temperature, fluctuations in dissolved oxygen, and variations in salinity. Probiotic supplementation may help fish cope with some forms of physiological stress by supporting intestinal health and immune function. Reduced stress can contribute to improved feeding activity, growth, disease resistance, and survival [10].

CONCLUSION

Erythrocyte osmoregulation represents an important cellular component of aquatic animal physiology. Through controlled ion movement, membrane transport, volume adjustment, and metabolic regulation, erythrocytes maintain conditions that permit continued oxygen transport under changing environmental circumstances. Further investigation across cultured species can improve knowledge of species-specific responses and help aquaculture researchers establish suitable environmental conditions for healthy blood function and efficient production.

REFERENCES

1. Dash P, Tandel RS, Bhat RA, Mallik S, Pandey NN, Singh AK, et al. The addition of probiotic bacteria to microbial floc: Water quality, growth, non-specific immune response and disease resistance of *Cyprinus carpio* in mid-Himalayan altitude. Aqua. 2018; 495:961-969.
2. Amir I, Zuberi A, Kamran M, Imran M. Evaluation of commercial application of dietary encapsulated probiotic (*Geotrichum candidum* QAUGC01): Effect on growth and immunological indices of rohu (*Labeo rohita*, Hamilton 1822) in semi-intensive culture system. Fish Shellfish Immunol. 2019;95:464-472.
3. Tachibana L, Telli GS, de Carla Dias D, Goncalves GS, Ishikawa CM, Cavalcante RB, et al. Effect of feeding strategy of probiotic *Enterococcus faecium* on growth performance, hematologic, biochemical parameters and non-specific immune response of Nile tilapia. Aquac. Rep. 2020;16:100277.
4. Van Doan H, Lumsangkul C, Hoseinifar SH, Tongsiri S, Chitmanat C, Musthafa MS, et al. Modulation of growth, innate immunity, and disease resistance of Nile tilapia (*Oreochromis niloticus*) culture under biofloc system by supplementing pineapple peel powder and *Lactobacillus plantarum*. Fish Shellfish Immunol. 2021;115:212-220.
5. Rohani MF, Islam SM, Hossain MK, Ferdous Z, Siddik MA, Nuruzzaman M, et al. Probiotics, prebiotics and synbiotics improved the functionality of aquafeed: Upgrading growth, reproduction, immunity and disease resistance in fish. Fish Shellfish Immunol. 2022;120:569-589.
6. Jiang Z, Tian J, Bai X, McClements DJ, Ma C, Liu X, et al. Improving probiotic survival using water-in-oil-in-water (W1/O/W2) emulsions: Role of fish oil in inner phase and sodium alginate in outer phase. Food Chem. 2023;417:135889.
7. Tamilselvan M, Raja S. Exploring the role and mechanism of potential probiotics in mitigating the shrimp pathogens. Saudi J Biol Sci. 2024;31(3):103938.
8. El-Son MA, Elbahnaswy S, Khormi MA, Aborasain AM, Abdelhaffez HH, Zahran E. Harnessing the fish gut microbiome and immune system to enhance disease resistance in aquaculture. Fish Shellfish Immunol. 2025;163:110394.
9. Shija VM, Tran NT, Shehata AI, Zheng X, Zhang M, Bakky MA, et al. Synergistic impacts of probiotics and prebiotics (synbiotics) in fish farming. Fish Shellfish Immunol. 2026:111358.
10. Vahdat S, Noori F, Van Stappen G, Agh N. Dietary inclusion of agricultural by-products, algae, and probiotics in a dry feed: Effects on growth performance, survival, immune response, biochemical indices, and reproductive traits of *Artemia parthenogenetica*. Aquac. Rep. 2026;46:103342.