

Original Paper

PURIFICATION AND CHARACTERIZATION OF *Aeromonas media* KLU 11.16 CHITOSANASE ISOLATED FROM SHRIMP WASTE

Ekowati Chasanah¹, Gintung Patantis¹, Dewi Seswita Zilda¹, Mahrus Ali² and Yenny Risjani³

¹ Research Center of Marine and Fisheries Product Processing and Biotechnology,
Jl. KS. Tubun, Jakarta, Indonesia; ekowati_ch@yahoo.com

² Department of Fisheries, Faculty of Agriculture, University of Lampung, Jl. Prof. Sumantri Brojonegoro,
Lampung

³ Faculty of Fisheries and Marine Sciences, Brawijaya University, Malang

Received : August, 10, 2011 ; Accepted : September, 26, 2011

ABSTRACT

*Our previous study found that KLU 11.16, isolated from shrimp waste secreted chitinolytic enzymes. The crude enzyme was interesting since their chitooligosaccharide was able to inhibit some pathogenic bacteria. In this study we report a purification and characterization of the chitosanase enzyme produced and the identification of the KLU 11.16. Purification of the enzyme was done two steps by ion exchange chromatography followed by gel filtration. Two out of 4 peaks from Gel Filtration step, i.e. fraction 16 and 33 were capable of hydrolyzing 100% deacetylated chitosan, indicating that both fractions contained chitosanase enzyme. The enzyme from fraction 16 had approximate molecular weight of 98.3 kDa. The enzyme worked optimally at temperature of 30°C, and pH 6. Addition of Ca²⁺, Fe²⁺, K⁺, Na⁺ ions in the form of Cl₂ salt and detergent Triton X-100 increased the enzyme activity, while Co²⁺, Mn²⁺ and Zn²⁺ ions in the same concentration decreased the enzyme activity. Addition of EDTA and SDS significantly decreased the enzyme activity. Molecular based identification revealed that KLU 11.16 was 99% similar to *Aeromonas media*.*

Keywords : purification; characterization; *Aeromonas media* ; chitosanase

Correspondence: Phone: +62-21-53650158; Fax: +62-21-53650157; E-mail: ekowati_ch@yahoo.com

INTRODUCTION

Chitosanase (EC 3.2.1.132) is one of chitinolytic enzymes group, a member of Glycoside Hydrolase (GH) which catalyze a hydrolysis process of β -1,4 linkages between N-acetyl-D-glucosamine and D-glucosamine residue in a partly acetylated chitosan and between D-glucosamine residue of 100% deacetylated chitosan. Chitosanases are categorized in three main classes based on their substrate specificities, i.e. the first class is the chitosanases that degrade chitosan upon recognizing a GlcNAc-GlcN bond, those that recognize both the GlcNAc-GlcN and the GlcNGlcN bond, and those that are specific to the GlcNGlcN bond only. However, the ability to hydrolyze 100% deacetylated chitosan was used to differentiate this enzyme from other enzyme degrading 1,4- β -glucosidic linkage (Park *et al.*, 1999; Chulhong *et al.*, 2011).

There is growing attention to the chitinolytic enzyme especially-chitosanase enzymes since the chitooligosaccharides produced by this enzyme are useful for the application of various industries. In the form of oligomer (chitooligosaccharide), the chitosan has better solubility in water while maintaining and even better performing biological activities such as antimicrobial and protective effects against infection with some pathogens (Jeon & Kim, 1998; Tsai *et al.*, 2000; Kumar *et al.*, 2005, Chasanah *et al.*, 2006b; Yan Wang *et al.*, 2007; Liang *et al.*, 2010), antitumor activity and immune-enhancing effects (Murao *et al.*, 1992; Suzuki, 1996; Nam *et al.*, 1999; Kendra and Kim, 1998, Jeon *et al.*, 2000; Wahyuni *et al.*, 2007). Enzymatic process of chitooligosaccharide production is more preferable compared to chemical and physical process due to safety reason. Among the enzyme

degrading chitosan such as cellulose, pectinase and lysozyme, chitosanase has ability to hydrolyze and produce specific oligosaccharide from chitosan (chitooligosaccharide) with high degree of deacetylation.

Worldwide annual recovery of chitin from the processing of marine crustaceans is approximated by 37,300 metric tons (MT), while potency of shrimp and crustacean waste in Indonesia is estimated $\pm 76,657 - 114,986$ MT of shrimp shell and 3,643 – 4,128 MT of crab shell equals to 12,045 – 17,867 MT a year chitin (Fawzya, 2009). The used of Crustacean waste for chitin production by proteolytic enzymatic methods has been reported (Risjani and Yuniarta, 2008; Liang *et al.*, 2010). Commercial interest in the utilization of chitin and its derivatives has led to the need for inexpensive and reliable sources of enzymes. Production of a suitable and stable enzymes is an important element in the utilization of shellfish wastes, not only solving environmental problems but also promoting the economic value of marine products.

Our group has isolated chitinolytic bacteria from various Indonesia marine environment/and traditional fishery products, *i.e.* from several sponges (Uria *et al.*, 2005, and Chasanah *et al.*, 2007), *terasi* (Zilda *et al.*, 2006 and Noviendri *et al.*, 2006) and from shrimp waste (Chasanah *et al.*, 2009a). One isolate from shrimp processing waste, KLU 11.16, was further studied due to its potential. Previous result showed that chitooligosaccharide produced by the crude enzyme was capable of inhibiting *Pseudomonas aurogenosa* and *V. parahaemolyticus* (Chasanah *et al.*, 2009b). This paper will report purification and characterization of the KLU 11.16 and the isolate identification.

MATERIALS AND METHODS

Production and extraction of enzyme.

Bacterial cells were cultivated in the Minimal Synthetic Medium (MSM) supplemented with 0.5% colloidal chitin as the sole carbon source (Chasanah *et al.*, 2006a). The medium consisted of 0.1% K_2HPO_4 , 0.01% $MgSO_4 \cdot 7H_2O$, 0.1% NaCl, 0.7% $(NH_4)_2SO_4$, 0.05% yeast extract. One full loop of the fresh bacterial cells grown on an agar plate was picked up and inoculated into the

20-ml starting medium. After 15 hours of incubation in the shaking water bath (37°C, 100 rpm), 10% (v/v) of the resulting culture was transferred into the 80-ml production medium, followed with an one-day incubation at the same temperature and shaking as above. Crude chitosanase enzyme was obtained by centrifuging the resulting culture at 9,000 $\times g$ (4°C, 25 min).

Purification of the enzyme.

Purification was initiated by concentrating the enzyme extract by ammonium sulphate precipitation technique. A series concentration of 20% - 80% (saturation) ammonium sulphate was added slowly to enzyme extract and kept overnight at 4°C. The mixture was then centrifuged at 12,000 $\times g$ (4°C, 6 min). The precipitate obtained was suspended with 15 ml 0.05 M phosphate buffer, pH 7.0 and the suspension was dialyzed with the same buffer to remove the salt. Purification was conducted by Ion exchange and Gel Filtration chromatography. A 1 ml of dialysed sample, diluted with 4 mL milliQ water, was loaded in chromatography column containing DEAE Sepharose Hi Trap FF matrix (Amersham) previously equilibrated with Tris-Cl 0.02M buffer (pH 8). Targeted enzyme was eluted with gradient buffer of 0 - 1 M 0.02M NaCl Tris-Cl pH 8 buffer. Peaks showing chitosanase activity was checked for purity by SDS-PAGE. A 5 mL of unpure fraction resulted from previous purification above, showing chitosanase activity was loaded in chromatography containing Sephadex G-100. Before loaded with the sample, the column was previously equilibrated with enzyme buffer (0.05M phosphate buffer pH 6.0). The targeted enzyme was eluted with the same enzyme buffer with elution speed of 0.5 ml/h

Assay of chitosanase activity.

Assay of chitosanase activity was done based on the method described by Yoon, *et al.* (2000) with modification. The substrate, 0.1% soluble chitosan (100 μ l), was added with phosphate buffer (0.05M,) pH 6.0 in an eppendorf tube. Then 100 μ l of enzyme solution was added into the tube followed with a 30-min incubation at 37°C. The enzyme reaction was stopped by

immersing the tubes for 5 minutes at boiling water to stop the reaction. The resulting mixture (200 μ l) was put in a reaction tube filled with 800 μ l water. After adding with 1 ml of Schales reagent, it was boiled for 15 min. and then centrifuged at 8.000 rpm for 10 min. The absorbance of the supernatant was measured at 420 nm using spectrophotometer. One unit of chitosanase activity was defined as the amount of enzyme that produce reducing sugar corresponding to 1 μ mol of D-glucosamine per minute.

Measurement of protein content.

The protein content of the crude enzyme was estimated according to the Lowry method with Bovine Serum Albumin (BSA) as standard (Bollag and Edelstain, 1991). Samples (100 μ L) was mixed with Lowry reagent (900 μ L), incubated for 15 minutes, and followed by addition of 3 mL Follin reagent. After vortexing and incubating for 45 minutes, the absorbancy to of the mixture was read using spectrophotometry within 540 nm wavelength. Blank was aquadest to replace the samples, while standard curve was set at concentration range of 0.05 – 0.2 mg/mL of BSA.

Determination of optimum pH and temperature.

The optimal pH of chitosanase activities was determined at 37°C by using overlapping pH ranges. To generate such pH ranges, the following buffer used were 0.2 M citrate buffer (pH 4.0-6.0), 0.2 M phosphate buffer (pH 6.0-8.0), 0.2 M borate buffer (pH 8.0-9.0), 0.2 M glycine buffer (pH 9.0-10), 0.2 M NaOH-borate buffer (pH 10). The optimal temperature of the enzyme activity was measured at different temperatures (30, 37, 50, 60, 70°C) at optimum pH. The measurement procedure and conditions followed the standard assay as described previously.

Effect of metals.

The effects of divalent cations as chlorides metals on the enzyme activity were measured at the final concentration of 1 mM. The tested divalent cations included Mg^{2+} , Mn^{2+} , Li^{2+} , Ni^{2+} , Co^{2+} , Zn^{2+} , Ba^{2+} and Ca^{2+} . As control, the enzyme solution without addition of any metals

was used. Monovalent cations (K^+ , Na^+ , NH_4^+) as well as a trivalent cation (Fe^{3+}) were also tested in this experiment. The effect of EDTA on the enzyme activity was determined in the same concentration.

SDS-PAGE and detection of the enzyme by zymogram.

SDS-PAGE and zymogram were performed using 10% (w/v) polyacrylamide gel. The substrates used for zymogram were 0.1% of 85% deacetylated degree soluble chitosan. Samples were run without boiling with addition of gliserol and 2.5% SDS in 125 mM Tris-HCl (pH 6.7). Electrophoresis was run at constant current of 50 and 25 mA, 100 volt at 40°C. After separation, the gel was renatured following procedure of Chasanah (2004) for chitosanase enzyme.

RESULTS AND DISCUSSION

Identification of KLU 11.16

KLU 11.16 was gram negative, rod (basil) bacteria. Based on 16S rRNA, the bacteria was closed to *Aeromonas media* as shown in phylogenetic tree (**Fig. 1**). The isolate has been isolated from exoskeleton of shrimp which has been sampled at one of the shrimp industries at Muara Baru, Jakarta. The pH and temperature of the waste was 8 and 20°C. It was likely that the shrimp has been treated with water containing chlorine. Therefore, it was suspected that the microbial flora contained in the shrimp waste was the one which can withstand with chlorine and low temperature. From the results (**Fig. 1**), the microbe has been closely identified as *Aeromonas media*. The microbe could be from the water where the shrimp has been cultivated or could be from contamination. Janda & Duffey (1988) and Altwegg & Geiss (1989) reported that *Aeromonas* spp. has been isolated from water, but some of them were implicated as a cause of diarrheal on man. *Aeromonas media* has been isolated from environment (water), infected fish and some of them have been found at diarrhea patient stools. Further study on this bacteria found that this bacteria was potentially enterotoxigenic (Singh, 2000).




Fig. 1. Phylogenetic tree of KLU 11.16 isolate

Aeromonas sp producing chitinolytic enzymes has been reported by Mitsutomi (1990), Huang (1996), Choi *et al.*, (2003), Kojima (2005), Kuk (2005 and 2006), and Sun *et al.*, (2009). *Aeromonas hydrophyla* was reported to produce 94 kDa chitinase while *Aeromonas schubertii* has been reported to produce 30 and 38 kDa chitinase. Charles-Rodrigues *et al.*, (2008) has isolated *Aeromonas* sp and produced a 20 kDa chitosanase from the bacteria, while Sun *et al.*, (2009) reported a 70 kDa chitosanase from *Aeromonas* sp screened from soil. This report is likely the first report on chitosanase from *Aeromonas media*. This isolate secreted not only chitinolytic enzymes but also

proteolytic enzyme (Chasanah *et al.*, 2009b) when cultivated in chitin medium.

Purification and characterization of chitosanase

Ammonium sulphate precipitation method was used to concentrate the extracellular enzyme produced by the *Aeromonas media* KLU 11.16 before entering purification step. **Fig. 2** shows that the enzyme could be concentrated by addition of 50% (saturation) ammonium sulphate. The extract was then dialysed using 10,000 Da molecular weight cut off (MWCO) membrane .




Fig. 2. Optimization of ammonium sulphate addition to concentrate the enzyme

Fig. 3 shows result of first step purification. There were 2 peaks of chitosanase at fraction no. 8 and 9. Result on SDS-PAGE showed that both fraction had 3 bands, *i.e.* 22,4

kDa, 28,6 kDa dan 98,3 kDa. Therefore, both fractions were mixed and runned at second system of purification, *i.e.* Gel filtration.




Fig. 3. Purification profile of the enzyme using DEAE *Sepharose*




Fig. 4. Gel Filtration profile using *Sephadex G-100*

Using Sephadex G-100, there were 4 peaks of chitosanase, *i.e.* fraction 16, fraction 21, fraction 27 and fraction 33. Activity and zymogram based assay using 100% deacetylated soluble chitosan as substrate found that among the 4 fractions only 2 fractions, *i.e.* fraction 16 and 33 were capable of degrading 100% deacetylated chitosan, assuring that both fractions contains chitosanase enzyme. The ability to degrade 100% deacetylated chitosan have been used to

differentiate chitosanase among other chitinolytic enzymes such as chitinase (Chasanah, 2004; Park *et al.*, 1999). Result of SDS-Page (**Fig. 6**) and Zymogram (not presented) of the fraction 16 was a single band of protein enzyme. The molecular weight was around 98.3 kDa. This fraction was used further for characterization study.




Fig. 5. Enzyme assay of 4 fractions resulted from Gel Filtration chromatography using 85% and 100% deacetylated chitosan

Pure enzyme performed best at 30⁰C (**Fig. 7**) and at pH 6 (**Fig. 8**). Activity of the pure enzyme was increased by 1 mM Ca²⁺, Fe²⁺,

K⁺, Li⁺ dan Na⁺ ion (in the form of Cl₂ salt), ranging from 148% to 340%. While addition of Co²⁺, Mn²⁺ dan ion Zn²⁺ in the same




Fig. 6. SDS- PAGE of the protein enzyme from purification step

Note : SDS-PAGE (left) : M = LMWMarker ; SFC = crude enzyme, 9 = fraction 9 of DEAE Sepharose FF, 16 = fraction 16 of Gel Filtration Sephadex G-100




Fig. 7. Optimum temperature of fraction 16 KLU 11.16 chitosanase

concentration have decreased the enzyme activity (Fig. 9). Addition of additives, *i.e.* EDTA and SDS in the concentration of 1mM decreased significantly the chitosanase activity, while addition of β -mercaptoethanol decreased enzyme activity by 68%, but detergent Triton X-100 increased the enzyme activity (Fig. 10). Addition of additives, *i.e.* EDTA and SDS in the concentration of 1mM decreased significantly (100%) the chitosanase, while addition of β -mercaptoethanol decreased enzyme activity by 68%. Detergent Triton X-100 increased the enzyme activity. Compared to other chitosanases studied so far, this chitosanase is considered unusual due to its high molecular weight (98.3 kDa). Another *Aeromonas* chitosanase having high molecular weight has been reported by Sun *et al.*, (2009). *Aeromonas* sp. HG08 isolated

from soil secreted the chitosanase (named AsChi) with molecular weight of 70 kDa. The enzyme performed well at optimum pH of 6.0 and temperature of 55°C. On the contrary of *Aeromonas media* KLU 11.16, the activity of *Aeromonas* sp HG08 chitosanase was markedly enhanced by Mn^{2+} and inhibited by Fe^{3+} , Cu^{2+} , Ag^+ and Hg^{2+} . Recent report on chitosanase produced by marine bacterium *Bacillus subtilis* CH-2 has been isolated from the intestine of *Sebastiscus marmoratus* (scorpion fish) (Chulhong *et al.*, 2011). The molecular mass of the purified chitosanase was detected as 29 kDa, and the enzyme was performed well at the pH and temperature of 5.5 and 60°C, respectively. The purified chitosanase was continuously thermostable at 40°C.




Fig.8. Optimal pH of fraction 16 KLU 11.16 chitosanase




Fig. 9. Effect of ion metal to the fraction 16 KLU 11.16 chitosanase activity




Fig. 10. Effect of additives to fraction 16 KLU 11.16 chitosanase

CONCLUSION

Chitosanase from *Aeromonas media* KLU 11.16 isolated from shrimp waste has been successfully purified by two steps purification. One of the two chitosanases (fraction 16) had approximate molecular weight of 98.3 kDa. The enzyme worked optimally at temperature of 30°C, and pH 6. Addition of Ca²⁺, Fe²⁺, K⁺, Na⁺ ions in the form of Cl₂ salt and detergent Triton X-100 increased the enzyme activity. The presence of Co²⁺, Mn²⁺ and Zn²⁺ ions and EDTA and SDS in the same concentration decreased the enzyme activity.

The utilization of shellfish (shrimp) waste as a sources of microbes and raw material of chitin, will not only solves environmental problems but also decreases the production cost of microbial chitosanases. The production of inexpensive chitosanase is an important element in the process of chitooligosaccharide production. From this finding, we have an alternative local bacterial chitosanase that can be use for enzymatically chitooligosaccharide production. Since properties of the enzymatically produced chitooligosaccharide, albeit its application, is very depending on the enzyme used, further study on the action of the pure enzyme to hydrolyze chitosanase is very

urgently needed. Since the chitosanase isolated was quite unique, having high molecular weight, and firstly reported from *Aeromonas media*, further protein characterization such as its N-terminal amino sequence might also be worthed to add scientific data on bacterial chitosanase from *Aeromonas media* KLU 11.16 which has been isolated from Indonesia.

REFERENCES

- Altwegg M. and H.K. Geiss 1989. CRC Crit. Rev. Microbiol 16 : 253-286
- Charles-Rodriguez AV., J.E. Mauricio-Benavides, Y. Garza-Garcia, C.N. Aguilar J. Rodriguez and J.C. Contreras-Esquivel 2008. Chitosanase production by a new bacterial sources. *Res J. Biol. Sci.* 3(8) : 957-963
- Chasanah, E. *et. al.* 2007. Research on Depolymerization of Chitosan by Enzymatic Methods to Produce Chitooligosaccharides (in Indonesian). Technical Reports. Research Centre for Marine and Fisheries Product Processing and Biotechnology. Department of Marine and Fisheries, Jakarta.
- Chasanah, E., M. Ilmi, dan Mangunwardoyo. 2009a. Penapisan bakteri kitinolitik dari limbah pengolahan udang. *Jurnal Pasca Panen dan Bioteknologi Kelautan dan Perikanan* vol.4 (1) (in Indonesian).
- Chasanah E., Ali M. and Ilmi M. 2009b. Characteristics of extracellular enzymes from bacteria isolated from shrimp waste cultivated on chitin medium. Presented at International Seminar 2009 : From Ocean for Food, Energy and Sustainable Resources and Environmenton. Unair, Surabaya, Indonesia :18 November 2009.
- Chasanah, E., P. Hariyadi, A.B. Witarto, J.K. Hwang, M.T. Suhartono. 2006a. Biochemical Characteristic of Chitosanase from the Indonesian *B. licheniformis* MB-2. *J. Mol. Biotechnol* 33 (2) : 93 – 102.
- Chasanah E., Meidina, M.T.Suhartono 2006b. Antibacterial potency of chitosan oligomer produced by *Bacillus licheniformis* MB-2 chitosanase. *Indonesian Fisheries Research Journal* Vol. 14 (2) : 91-95
- Choi, Y. J., E. J. Kim, Z. Piao, Y. C. Yun, Y.C. Shin. 2003. Purification and characterization of chitosanase from *Bacillus* sp. strain KCTC 0377BP and its application for the production of chitosan oligosaccharides. *Appl. Environ. Microbiol.* 70 (8) : 4522-453.
- Chulhong O., Zoysa MD., Do-Hyung Kang¹, Lee Y., Whang I., Nikapitiya C., Soo-Jin Heo¹, Kon-Tak Yoon¹, Affan A., and Lee J. 2011. Isolation, Purification, and Enzymatic Characterization of Extracellular Chitosanase from Marine Bacterium *Bacillus subtilis* CH2. *J. Microbiol. Biotechnol.* 21(10), 1021–1025
- Fawzya, YN., Pratitis, A. and Chasanah, E. 2009. Characaterization of chitosanase from KPU 2123 bacterial isolate and its application to produce chitosan oligomer. (in Indonesian).. *J. Penelitian Pengolahan Produk dan Bioteknologi Kelautan dan Perikanan.* vol.4 (1)
- Fawzya Y.N. and S.Wibowo 2009. Exploration of Indonesia Marine Chitinolytic Enzymes and their application. *Indonesian Marine and Fisheries Product Processing and Biotechnology.* Research Center for Marine and Fisheries Product Processing and Biotechnology. Jakarta
- Janda JM., and P.S. Duffey 1988. Mesophilic aeromonads in human disease : current taxonomy, laboratory identification and infectious disease spectrum. *Rev Infect Dis* 10 : 980-997.
- Jeon, Y.J. and S.K. Kim, 2000. Potential Immuno-Stimulating Effect of Antitumoral Fraction of Chitosan Oligosaccharides. The Korean Society for Chitin and Chitosan Department of Agricultural. Dongguk University, Gyeongju. Korea. 6 :4

- Kendra, Y.J. and S.K. Kim, 1998. Bioactivities of chitosan Oligosaccharides and their Derivatives in *Abstract of Asia Pasific Chitin and Chitosan Symposium*, pp. 64, Kilung, Taiwan.
- Liang, T.W., K. Yi-Hsuan, W. Pei-Chen, W. Chuan-Lu, N.A. Dzung and W. San-Lang. 2010. Purification and Characterization of a Chitosanase and a Protease by Conversion of Shrimp Shell Wastes Fermented by *Serratia Marcescens Subsp. Sakuensis* TKU019. *Journal of the Chinese Chemical Society*, 57, 857-863
- Kumar A.B.V., M.C.Varadaraj L.R. Gowda, R.N.Tharanathan. 2005. Characterization of chitooligosaccharides prepared by chitosanolytic with the aid of papain and pronase, and their bactericidal action against *Bacillus cereus* and *Escherichia coli*. *Biochemical Journal Immediate Publication*. BJ20050093.
- Murao, S., T. Kawada, H. Itoh, H. Oyama, and T. Shin. 1992. Purification and Characterization of a Novel Type of Chitinase from *Vibrio alginolyticus* TK-22. *Biosci. Biotech. Biochem.* 56(2):368-369.
- Nam, M.Y., S. Yun-Hee, K. Se-Kwon, K. Cheorl-Ho, and N. Kyung-Soo, 1999. Inhibitory Effect of Chitosan Oligosaccharides on the Growth of Tumor Cells. The Korean Society for Chitin and Chitosan Department of Agricultural. Dongguk University, Gyungju. Korea. Volume 4 Issue 4.
- Noviendri, D., Y.N. Fawzya, dan E. Chasanah, 2006. Characterization and kinetics of chitinase produced by T5a1 bacteria isolated from terasi (in Indonesian). *J. Penelitian Pengolahan Produk dan Bioteknologi Kelautan dan Perikanan*.
- Risjani Y. and Yuniarta. 2008. *Mucor pusillus* production and Aspartic Protease Enzyme Activity in the bioconversion of Crustacean waste *Penaeus sp* and *Portunus pelagicus*. (in Indonesian) *Agritek*, 16 (1): 30-33
- Singh D.V. 2000. A putative heat-labile enterotoxins expressed by strains of *Aeromonas media*. *J. Med. Microbiol.* Vol 49 : 685-689
- Sun Y., J. Zhang and S.Wang 2009. Purification and characterization of the chitosanase from *Aeromonas sp.* HG08. *African Journal of Biotechnology* Vol. 8 (12), pp. 2830-2834, 17 June, 2009.