OPEN ACCESS Freely available online

ISSN: 2155-9546
Journal of Aquaculture Research & Development

Perspective

Quantitative Assessment of Intestinal Villi Morphometry in Cultured Aquatic Species

Robson Marins*

Department of Marine Sciences, University of Algarve, Faro, Portugal

DESCRIPTION

Intestinal villi morphometry is an important area of investigation in aquaculture because the physical organization of the intestinal surface influences how efficiently cultured animals process and absorb nutrients. The intestine is not simply a passage through which feed travels. Its internal surface contains folds, villi, epithelial cells, mucus-producing cells, blood vessels, and associated tissues that collectively contribute to digestion, absorption, protection, and transport. Quantitative examination of these structures can provide useful information about intestinal condition and may help researchers evaluate how feed composition, environmental conditions, developmental stage, and husbandry practices affect digestive performance.

Villi are projections of the intestinal mucosa that increase the available surface area for contact between digested nutrients and absorptive epithelial cells. Their dimensions differ among species and along different sections of the digestive tract. Fish, shrimp, and other cultured aquatic organisms have distinct intestinal arrangements, meaning that morphometric measurements must be interpreted according to species-specific anatomy. Parameters such as villus length, villus width, epithelial thickness, villus density, mucosal thickness, and the ratio between villus length and width can provide numerical descriptions of intestinal architecture.

In fish, intestinal villi are generally covered by epithelial cells responsible for nutrient uptake and interaction with the intestinal contents. These cells are supported by connective tissue containing blood vessels and other structural components. The dimensions of the villi can change according to nutritional status, age, water conditions, disease exposure, and feed formulation. Young fish may show progressive changes in villus development as their digestive system matures. Consequently, age and body size should be recorded carefully when comparing morphometric observations between experimental groups.

Histological examination is commonly used to evaluate intestinal morphology. Tissue samples are collected from defined intestinal regions, preserved, processed, sectioned, stained, and examined using light microscopy. Digital imaging systems can then be used to measure selected structures. Consistent orientation of tissue sections is particularly important because an oblique section may make villi appear shorter or wider than they actually are. Researchers should therefore use comparable anatomical regions and standardized sectioning procedures when conducting quantitative assessments.

Villus length is one of the frequently measured characteristics. Longer villi may provide a greater epithelial surface for nutrient interaction, although length alone does not determine intestinal function. Villus width is another useful parameter because unusually broad or narrow structures may reflect differences in tissue organization. Villus density can provide additional information by describing the number of villi within a defined mucosal area. When these measurements are considered together, they offer a more complete representation of intestinal architecture than any single measurement.

Epithelial thickness can also be evaluated because the intestinal epithelium forms the immediate interface between the intestinal contents and underlying tissues. Changes in epithelial thickness may occur in response to dietary ingredients, microbial conditions, physical irritation, or physiological adaptation. Goblet cell abundance is another useful observation. These cells produce mucus that contributes to lubrication and protection of the intestinal lining. Recording goblet cell distribution alongside villus dimensions can therefore provide additional information about mucosal condition.

Dietary composition is strongly connected with intestinal morphology. Protein sources, lipid levels, carbohydrate composition, fiber content, mineral concentrations, and functional feed ingredients can influence the structure of the digestive tract. Alternative protein ingredients, for example, may contain compounds that alter intestinal responses when included at high levels or without appropriate processing. Conversely, suitable feed ingredients may support favorable epithelial organization and nutrient utilization. Morphometric

Correspondence to: Robson Marins, Department of Marine Sciences, University of Algarve, Faro, Portugal, E-mail: robson.marins@ualg.pt

Received: 03-Aug-2026, Manuscript No. JARD-26-32967; **Editor assigned:** 06-Aug-2026, PreQC No. JARD-26-32967 (PQ); **Reviewed:** 20-Aug-2026, QC No. JARD-26-32967; **Revised:** 27-Aug-2026, Manuscript No. JARD-26-32967 (R); **Published:** 03-Sep-2026, DOI: 10.35248/2155-9546.26.17.1140

Citation: Marins R (2026). Quantitative Assessment of Intestinal Villi Morphometry in Cultured Aquatic Species. J Aquac Res Dev. 17.1140.

Copyright: © 2026 Marins R. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

analysis allows researchers to compare these effects using measurable anatomical variables.

Feed additives are also being investigated in relation to intestinal structure. Probiotics, prebiotics, organic acids, enzymes, plant-derived compounds, and other dietary components may influence the microbial environment or digestive processes within the intestine. Their effects should not be judged solely by changes in growth rate. Combining growth measurements with villus morphology, digestive enzyme activity, nutrient retention, and intestinal microbial observations can provide a broader interpretation of animal responses.

Environmental conditions can influence intestinal tissues as well. Temperature, salinity, dissolved oxygen, water quality, and stocking density may affect feeding activity and metabolic demand. Exposure to unsuitable environmental conditions can modify energy allocation and may be accompanied by alterations in digestive tissues. For this reason, intestinal morphometry should be assessed together with relevant water-quality measurements whenever environmental treatments are part of an aquaculture experiment.

CONCLUSION

Overall, intestinal villi morphometry provides a practical quantitative method for evaluating digestive tissue organization in cultured aquatic animals. Measurements of villus length, width, density, epithelial thickness, mucosal thickness, and related cellular features can contribute valuable information to nutrition and health studies. When combined with growth performance, feed utilization, biochemical indicators, microbial analysis, and controlled environmental observations, morphometric data can strengthen the interpretation of aquaculture experiments and support more precise evaluation of factors affecting intestinal development and nutrient absorption.