OPEN ACCESS Freely available online

ISSN: 2155-9546
Journal of Aquaculture Research & Development

Commentary

Quorum-Sensing Mechanisms and Microbial Interactions in Aquaculture

Wenbo Gui*

Department of Aquatic Animal Science, Shanghai Ocean University, Shanghai, China

DESCRIPTION

Aquaculture production depends on a stable interaction between cultured animals, microorganisms, water quality, feed, and environmental conditions. Microbial communities are present in ponds, tanks, biofilters, sediments, cages, and the digestive systems of aquatic animals. These microorganisms do not exist as isolated populations. Many bacteria communicate with one another through chemical signals that coordinate changes in collective behavior. This process is known as quorum sensing. In aquaculture, quorum sensing has gained attention because bacterial communication can influence biofilm formation, enzyme secretion, nutrient utilization, motility, virulence-associated activity, and interactions between microorganisms and cultured animals.

Quorum sensing generally occurs when bacteria release small signaling molecules into their surrounding environment. As the bacterial population increases, the concentration of these molecules also rises. Once a suitable concentration is reached, bacterial cells detect the signals and alter the expression of selected genes. The resulting response can involve a large group of cells rather than an individual bacterium. This communication system allows microorganisms to respond collectively to changes in population density and environmental conditions.

Different bacterial groups use different signal molecules. N-acyl homoserine lactones are widely associated with communication among many Gram-negative bacteria, while autoinducer-2 has been associated with communication involving several bacterial groups. Gram-positive bacteria commonly use short signaling peptides. Aquaculture environments can contain numerous bacterial species, so several communication systems may operate simultaneously. Their interactions can influence the structure and activity of microbial communities within culture systems.

Biofilm formation is one of the important processes associated with bacterial communication. Biofilms consist of microbial cells attached to a surface and surrounded by an extracellular matrix. In aquaculture facilities, biofilms can develop on tank walls, pipes, aeration equipment, filters, cages, feeding systems, and other submerged surfaces. Quorum sensing can regulate bacterial behaviors involved in attachment, extracellular polymer production, and biofilm maturation. Mature biofilms can provide microorganisms with protection from environmental fluctuations and may contain bacteria with different physiological characteristics.

The relationship between quorum sensing and fish or shellfish health is particularly relevant. Several aquatic pathogens use chemical communication systems to coordinate behaviors associated with infection. *Vibrio* species are especially important in marine and brackish-water aquaculture. Certain *Vibrio* populations can use quorum-sensing signals to regulate extracellular enzymes, surface attachment, motility, and other cellular activities. These processes may affect the interaction between bacteria and shrimp, fish, or other cultured organisms. Understanding these communication mechanisms can therefore contribute to improved disease management.

Quorum sensing is not limited to harmful microorganisms. Beneficial bacteria can also use communication systems to regulate their activities. Probiotic bacteria used in aquaculture may interact with other microorganisms through signaling compounds and may influence the microbial composition of the surrounding water or intestinal tract. Some probiotic strains have been investigated for their ability to interfere with communication processes used by undesirable bacteria. Such interactions may reduce the expression of selected bacterial behaviors without necessarily eliminating the target microorganisms.

The interruption of bacterial communication is commonly described as quorum quenching. Quorum-quenching activity can occur through several mechanisms. Some microorganisms produce enzymes that degrade signaling molecules, while others modify the signals so that they are no longer recognized by bacterial receptors. Certain compounds may also interfere with signal production or signal detection. In aquaculture, these mechanisms are being investigated as possible approaches for reducing undesirable bacterial activity.

Correspondence to: Wenbo Gui, Department of Aquatic Animal Science, Shanghai Ocean University, Shanghai, China, E-mail: wenbo.gui@shou.edu.cn

Received: 29-Jun-2026, Manuscript No. JARD-26-32754; **Editor assigned:** 01-Jul-2026, PreQC No. JARD-26-32754 (PQ); **Reviewed:** 15-Jul-2026, QC No. JARD-26-32754; **Revised:** 22-Jul-2026, Manuscript No. JARD-26-32754 (R); **Published:** 29-Jul-2026, DOI: 10.35248/2155-9546.26.17.1131

Citation: Gui W (2026). Quorum-Sensing Mechanisms and Microbial Interactions in Aquaculture. J Aquac Res Dev. 17:1131.

Copyright: © 2026 Gui W. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

CONCLUSION

Aquaculture quorum sensing represents an important area of microbial research because bacterial behavior is influenced not only by the presence of microorganisms but also by communication among them. Signal production, signal detection, biofilm development, microbial competition, and pathogen activity can affect culture conditions and animal health. Continued research into signaling molecules, quorum-quenching organisms, microbial communities, and environmental factors can contribute to more precise microbial management. The combination of molecular biology, microbiology, chemical analysis, and aquaculture engineering can provide a broader understanding of microbial communication and its practical relevance to fish and shellfish production.