

Regulation of Oncogene Expression via Targeting i-Motif in Nucleic Acids

Prashansa Agrawal*

Department of Chemistry, Case Western Reserve University, Cleveland, Ohio, 44106, USA

*Corresponding author: Prashansa Agrawal, Department of Chemistry, Case Western Reserve University, Cleveland, Ohio, 44106, USA, Tel: +1 216-368-2404; E-mail: prashansa.agrawal@case.edu

Received date: December 28, 2016; Accepted date: January 02, 2017; Published date: January 05, 2017

Copyright: © 2017 Agrawal P. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Editorial

The basic molecules required for sustaining life on earth include carbohydrates, lipids, proteins, and nucleic acids (DNA and RNA). DNA is considered to be the building block of life and can adopt various types of secondary structures. Gehring et al. revealed for the first time that DNA sequences having long cytosine stretch can form unconventional secondary structures called i-Motifs [1]. These i-Motifs are very dynamic intercalated, quadruple-helical structures and known to exist under acidic conditions. These cytosine stretches are not erratically located throughout the genome but enriched in the promoter regions of several oncogenes within the genome and in telomeres, the terminal regions of chromosomes. The formation and stabilization of i-motif may play a crucial role in gene expression [2-4], such as, the stabilization of the human telomeric i-motif by carboxylated single-walled carbon nanotubes inhibit telomerase that leads to telomere uncapping, DNA damage response and apoptosis [5].

These i-Motif structures are formed by hemi protonated cytosine-cytosine base pairs, which require protonation of one of the cytosines in the base pair (that is done by lowering the physiological pH *in vitro*). However, recent study using molecular crowding conditions, mirrored by carboxyl modified single-walled carbon nanotubes, have been shown to selectively induce human telomeric i-Motifs formation even at pH 8.0 [6]. The i-Motif structures can be formed from four strands, two strands or single strand of DNA. This single strand i-motif is referred to as intramolecular i-Motif which is considered to be potentially biologically relevant and is divided into two classes: class I having shorter loops regions (e.g., telomeric i-Motif) [7,8] and class II that has longer loops regions (bcl2 promoter i-Motif) [3]. The class II i-Motifs are more stable than class I i-Motifs due to extra stabilizing interactions induced within the longer loop regions.

The significance of i-motif structures has not been explored much *in vivo*. Its potential role in gene regulation has been proposed only recently [2-4,9-11]. Some examples include, B-cell lymphoma gene 2 (a pro-survival oncoprotein) which is over expressed in some cancers like breast cancer, pancreatic cancer and under expressed in some neurodegenerative diseases such as Alzheimer's and Parkinson's disease, thus leaving the cells more susceptible to apoptosis. The i-Motif formation on the cytosine rich bcl2 promoter region provides a mechanism for modulation of gene expression on being targeted by the compounds, IMC 48 (cholestane derivative) which up regulates transcription whereas IMC 76 (pregnanol derivative) down regulates gene expression [9]. Thus targeting the bcl2 i-motif by these compounds can help in regulating the gene expression. Another example is c-myc promoter i-Motif. This c-myc gene codes for transcription factor that has various function such as regulation of cell cycle, progression and cell growth [12]. The gene expression of c-myc is controlled by several proteins which bind to single strand cytosine

rich repeats (e.g., heterogeneous ribonucleoprotein K). With the formation i-Motif on the cytosine rich strand followed by its stabilization through small molecule compound may lead to suppression of gene expression [13].

Besides these above-mentioned i-Motifs, there are other promoter i-Motifs that have been under study, like vascular endothelial growth factor promoter region can form an i-Motif *in vitro* at acidic pH [14]. The cytosine rich promoter region of c-kit, kras, PDGF, c-myc, HIF-1 α and hTERT are also known to form i-Motif structures [15-17]. The cytosine rich repeats in RNA can also form i-Motif [18] but they are highly unstable due to the steric interactions in the sugar and repulsive interactions between 2' hydroxyl groups of the ribose sugars in the minor groove. Besides this, other less stable i-Motifs are: DNA-RNA hybrid i-Motif [19] and peptide nucleic acids i-Motif [20].

Targeting i-motif structure with small molecules may lead to prospective cure for cancer and other genetic disease and hence can be a potential target for drug design and modulation of gene expression. At present only few compounds have been tested that bind to i-Motif structures but they also show non-specific binding to other targets as well. The first binding study on telomeric i-Motif was done in 2000 [21]. They showed binding of TMPyP4 to telomeric i-motif. Besides this, there are other telomeric i-motif targeting compounds, such as, BisA [22], phenanthrolines [23], perylene-neomycin conjugate [24], and crystal violet [25], amino acid complexes of Terbium [26], ruthenium complexes [27] and mitoxantrone [28]. The bcl2 i-Motif targeting compounds are IMC 48 and IMC 76 [9]. In addition, i-Motif structure has vast application in the field of nanotechnology, for example, as sensors to map pH changes in living cells [29,30], switches for logic operations [31-33], designing Nano machines [34,35] and assembly of gold nanoparticles.

At present, i-Motif target based drug design is an emerging field. Since targeting i-Motif with small molecules is currently a challenging problem as there are not any compounds/drugs in the market that can specifically bind to a particular i-Motif. Researchers still have to go long way in order to find a potential drug candidate that can target, induce and stabilize i-Motif formation in the promoter region of oncogene thereby regulating its gene expression, which can be a breakthrough development in the field of cancer therapy.

References

1. Gehring K, Leroy JL, Gueron M (1993) A tetrameric DNA structure with protonated cytosine. cytosine base pairs. Nature 363: 561.
2. Sun D, Hurley LH (2009) The Importance of Negative Superhelicity in Inducing the Formation of G-Quadruplex and i-Motif Structures in the c-Myc Promoter: Implications for Drug Targeting and Control of Gene Expression. J Med Chem 52: 2863.

3. Brooks TA, Kendrick S, Hurley L (2010) Making sense of G-quadruplex and i-motif functions in oncogene promoters. *FEBS J* 277: 3459.
4. Choi J, Kim S, Tachikawa T, Fujitsuka M, Majima TJ (2011) pH-Induced Intramolecular Folding Dynamics of i-Motif DNA. *Am Chem Soc* 133: 16146-16153.
5. Chen Y, Qu K, Zhao C, Wu L, Ren J, et al. (2012) Insights into the biomedical effects of carboxylated single-wall carbon nanotubes on telomerase and telomeres. *Nat Comm* 3: 1074.
6. Li X, Peng Y, Ren J, Qu X (2006) Carboxyl-modified single-walled carbon nanotubes selectively induce human telomeric i-Motif formation. *Proc Natl Acad Sci USA* 103: 19658-19663.
7. Phan AT, Gueron M, Leroy JL (2009) The solution structure and internal motions of a fragment of the cytidine-rich strand of the human telomere. *J Mol Biol* 299: 123.
8. Leroy JL, Gueron M, Mergny JL, Helene C (1994) Intramolecular folding of a fragment of the cytosine-rich strand of telomeric DNA into an i-motif. *Nucleic Acids Res* 22: 1600-1606.
9. Kendrick S, Kang H, Alam MP, Madathil MM, Agrawal P, et al. (2014) The Dynamic Character of the BCL2 Promoter i-Motif Provides a Mechanism for Modulation of Gene Expression by Compounds that Bind Selectively to the Alternative DNA Hairpin Structure. *J Am Chem Soc* 136: 4161-4171.
10. Brazier JA, Shah A, Brown GD (2012) i-Motif formation in gene promoters: unusually stable formation in sequences complementary to known G-quadruplexes. *Chem Comm* 48: 10739-10741.
11. Kendrick S, Akiyama Y, Hecht SM, Hurley LH (2009) The i-Motif in the bcl-2 P1 Promoter Forms an Unexpectedly Stable Structure with a Unique 8:5:7 Loop Folding Pattern. *J Am Chem Soc* 131: 17667-17676.
12. Gonzalez V, Hurley LH (2010) The c-MYC NHE III1: Function and Regulation. *Annu Rev Pharmacol Toxicol* 50: 111-129.
13. Brooks TA, Hurley LH (2010) Targeting MYC Expression through G-Quadruplexes. *Genes Cancer* 1: 641-649.
14. Guo K, Gokhale V, Hurley LH, Sun D (2008) Intra molecularly folded G-quadruplex and i-motif structures in the proximal promoter of the vascular endothelial growth factor gene. *Nucleic Acids Res* 36: 4598-4608.
15. Manzini G, Yathindra N, Xodo LE (1994) Evidence for intramolecularly folded i-DNA structures in biologically relevant CCC-repeat sequences. *Nucleic Acids Res* 22: 4634-4640.
16. Day HA, Pavlou P, Waller ZAE (2014) i-Motif DNA: Structure, stability and targeting with ligands. *Bioorg Med Chem* 22: 4407-4418.
17. Lacroix L, Mergny JL, Leroy JL, Helene C (1996) Inability of RNA To Form the i-Motif: Implications for Triplex Formation. *Biochem* 35: 8715-8722.
18. Collin D, Gehring K (1998) Stability of Chimeric DNA/RNA Cytosine Tetrads: Implications for i-Motif Formation by RNA. *J Am Chem Soc* 120: 4069-4072.
19. Fedoroff OY, Rangan A, Chemeris VV, Hurley LH (2000) Cationic Porphyrins Promote the Formation of i-Motif DNA and Bind Peripherally by a Nonintercalative Mechanism. *Biochemistry* 39: 15083-15090.
20. Alberti P, Ren J, Teulade-Fichou MP, Guittat L, Riou JE, et al. (2001) Interaction of an acridine dimer with DNA quadruplex structures. *J Biomol Struct Dyn* 19: 505.
21. Wang L, Wu Y, Chen T, Wei C (2013) The interactions of phenanthroline compounds with DNAs: preferential binding to telomeric quadruplex over duplex. *Int J Biol Macromol* 52: 1-8.
22. Xue L, Ranjan N, Arya DP (2011) Synthesis and spectroscopic studies of the aminoglycoside (neomycin)-perylene conjugate binding to human telomeric DNA. *Biochemistry* 50: 2838-2849.
23. Ma DL, Kwan MHT, Chan DSH, Lee P, Yang H, et al. (2011) Crystal violet as a fluorescent switch-on probe for i-motif: label-free DNA-based logic gate. *Analyst* 136: 2692-2696.
24. Shi S, Zhao J, Geng X, Yao T, Huang H, et al. (2010) Molecular "light switch" for G-quadruplexes and i-motif of human telomeric DNA: [Ru(phen)₂(dppz)]²⁺. *Dalt Trans* 39: 2490-2493.
25. Wright EP, Day HA, Ibrahim AM, Kumar J, Boswell LJE, et al. (2016) Mitoxantrone and Analogues Bind and Stabilize i-Motif Forming DNA Sequences. *Sci Rep* 6: 39456.
26. Modi S, Swetha MG, Goswami D, Gupta GD, Mayor S, et al. (2009) A DNA Nanomachine that maps spatial and temporal pH changes inside living cells. *Nat Nanotechnol* 4: 325.
27. Surana S, Bhat JM, Koushika SP, Krishnan Y (2011) An autonomous DNA nanomachine maps spatiotemporal pH changes in a multicellular living organism. *Nat Commun* 2: 340.
28. Elbaz J, Wang ZG, Orbach R, Willner I (2009) pH-Stimulated Concurrent Mechanical Activation of Two DNA "Tweezers". A "SET-RESET" Logic Gate System. *Nano Lett* 9: 4510-4514.
29. Yang Y, Liu G, Liu H, Li D, Fan C, et al. (2010) An Electrochemically Actuated Reversible DNA Switch. *Nano Lett* 10: 1393-1397.
30. Li T, Ackermann D, Hall AM, Famulok M (2012) Input-Dependent Induction of Oligonucleotide Structural Motifs for Performing Molecular Logic. *J Am Chem Soc* 134: 3508.
31. Liu D, Balasubramanian S (2003) A Proton-Fuelled DNA Nanomachine. *Angew Chem Int Ed* 42: 5734-5736.
32. Wang ZG, Elbaz J, Willner I (2011) DNA Machines: Bipedal Walker and Stepper. *Nano Lett* 11: 304-309.
33. Wang W, Liu H, Liu D, Xu Y, Yang Y, et al. (2007) Use of the Interparticle i-Motif for the Controlled Assembly of Gold Nanoparticles. *Langmuir* 23: 11956-11959.
34. Sharma J, Chhabra R, Yan H, Liu Y (2007) pH-driven conformational switch of "i-motif" DNA for the reversible assembly of gold nanoparticles. *Chem Commun* 2007: 477-479.
35. Balasubramanian V, Huwylar J (2015) Nanomedicine in cancer therapy: Challenges, opportunities, and clinical applications. *J Control Release* 200: 138-157.