



Secondary Metabolites from *Pseudophaeolus soloniensis* : Isolation, Characterization and Antifungal Potential against Plant Pathogens

Maenia Castillo *

Department of Natural Products and Microbial Biotechnology, Federal University of Paraná, Curitiba, Brazil

DESCRIPTION

Plant diseases caused by phytopathogenic fungi represent a persistent challenge for agriculture because they can reduce crop productivity, compromise food quality and increase production costs. The continued dependence on conventional fungicides has also raised concerns involving environmental persistence, resistance development and effects on beneficial organisms. These concerns have encouraged greater interest in natural products obtained from fungi and other biological sources. Wood-inhabiting fungi are particularly promising because they produce diverse secondary metabolites that can participate in ecological interactions and may possess antimicrobial properties. *Pseudophaeolus soloniensis*, a wood-associated fungal species, is therefore of interest as a potential source of biologically active compounds. Secondary metabolites are chemically diverse molecules that are not necessarily required for basic cellular growth but can provide ecological advantages. Fungi produce compounds belonging to several structural groups, including phenolic substances, terpenoid derivatives, aromatic compounds, polyketides and other specialized metabolites.

The production can vary according to nutritional conditions, developmental stage, substrate and environmental factors. Because of this chemical diversity, fungal secondary metabolism represents an important reservoir of molecules that may interfere with the growth or development of microorganisms. Obtaining metabolites from *Piptoporellus soloniensis* generally begins with careful preparation of fungal material followed by extraction using appropriate solvents. Solvent polarity influences which classes of compounds are recovered, so sequential or complementary extraction approaches can provide a broader chemical profile. The resulting extracts can be concentrated and separated into fractions according to their chemical characteristics. Chromatographic techniques are commonly useful for separating complex mixtures, while spectroscopic approaches can provide information about molecular structures.

Combining chromatographic and spectroscopic data allows individual constituents to be distinguished and their chemical identities to be evaluated.

The antimicrobial potential of fungal metabolites is particularly relevant when directed toward phytopathogenic fungi. Plant pathogens can interfere with seed germination, root development, vascular function and photosynthetic tissues, depending on the species involved. Natural compounds that inhibit fungal growth may act through several mechanisms, including disruption of cell membranes, interference with energy metabolism, inhibition of essential enzymes or disturbance of cellular signalling. The effectiveness of a metabolite can vary considerably among fungal species because pathogens differ in their physiological characteristics and sensitivity to chemical compounds. Assessment of antifungal activity requires more than observing whether an extract produces a visible inhibition zone. Quantitative approaches can determine the concentration needed to restrict fungal growth and can help distinguish weak activity from substantial inhibition. Comparing crude extracts with purified fractions is also important because biological effects may result from one dominant compound, several active metabolites or interactions among multiple constituents. Chemical fractionation can consequently reveal whether antimicrobial activity becomes stronger as specific compounds become enriched.

The chemical complexity of fungal extracts creates both opportunities and challenges. A crude extract may contain numerous molecules with complementary biological effects, but its composition can make reproducibility difficult. Environmental conditions, culture substrate, extraction procedures and storage can influence metabolite concentrations. Consistent cultivation and standardized extraction are therefore important when evaluating the potential of *Piptoporellus soloniensis* as a source of antifungal agents. Natural products from wood-decaying fungi may also offer structural diversity that

Correspondence to: Maenia Castillo, Department of Natural Products and Microbial Biotechnology, Federal University of Paraná, Curitiba, Brazil, Email: maenia.castillo@ufpr.br

Received: 30-Aug-2025, Manuscript No. JDMGP-25-32404; **Editor assigned:** 02-Sep-2025, PreQC No. JDMGP-25-32404; **Reviewed:** 16-Sep-2025, QC No. JDMGP-25-32404; **Revised:** 22-Sep-2025, Manuscript No. JDMGP-25-32404; **Published:** 30-Sep-2025, DOI: 10.35248/2153-0602.25.16.392

Citation: Castillo M (2025). Secondary Metabolites from *Pseudophaeolus soloniensis*: Isolation, Characterization and Antifungal Potential against Plant Pathogens. J Data Mining Genomics Proteomics. 16:392.

Copyright: © 2025 Castillo M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

differs from compounds currently used in agricultural formulations. Molecules with unfamiliar chemical scaffolds can be valuable because they may interact with fungal targets through mechanisms distinct from those of established fungicides. Such diversity is particularly relevant to the continuing problem of fungicide resistance. Nevertheless, antimicrobial activity in laboratory conditions does not automatically indicate suitability for agricultural application. Further evaluation would be required to determine stability, plant compatibility, environmental behavior, toxicity and effectiveness under practical crop conditions.

The relationship between chemical composition and antifungal activity can provide important clues about the biological role of secondary metabolites in *Piptoporellus soloniensis*. Compounds

produced by a wood-associated fungus may contribute to competition with other microorganisms inhabiting the same ecological niche. Their activity against plant-pathogenic fungi could therefore reflect ecological functions that have developed through long-term microbial interactions. Characterizing these compounds expands understanding of fungal chemical diversity while also identifying molecules with possible practical value. The exploration of *Piptoporellus soloniensis* secondary metabolites demonstrates the potential of fungal biodiversity as a source of antimicrobial chemistry. Isolation and structural characterization can reveal compounds that merit further evaluation, while antifungal testing helps connect chemical identity with biological function.