OPEN ACCESS Freely available online

ISSN: 2155-9546
Journal of Aquaculture Research & Development

opinion

Semi-Intensive Shrimp Culture: Production Practices, Water Management, and Sustainable Farm Performance

Flávio Leite*

Department of Aquatic Systems, Federal University of Santa Catarina, Florianopolis, Brazil

DESCRIPTION

Semi-intensive shrimp culture is a widely used production approach that combines controlled pond management with natural biological processes. It is positioned between extensive farming, where shrimp depend mainly on naturally occurring food organisms, and intensive production, where stocking densities and management inputs are considerably higher. Semi-intensive systems generally use moderate stocking densities, prepared feeds, water exchange, aeration when needed, pond preparation, and regular observation of shrimp and environmental conditions. The objective is to produce a reliable shrimp harvest while maintaining suitable water quality and controlling production costs.

The success of semi-intensive shrimp farming depends on the interaction between shrimp biology, pond characteristics, feed management, water quality, weather, and farm practices. Species such as Litopenaeus vannamei and Penaeus monodon are commonly associated with pond-based shrimp production. Before stocking, farmers normally assess the suitability of the site, water source, soil characteristics, drainage arrangements, pond dimensions, and accessibility. A suitable location should provide dependable water while reducing the possibility of contamination from agricultural, domestic, or industrial sources.

Pond preparation is an important activity before each production cycle. Farmers may remove excessive organic material, repair pond banks, clear unwanted vegetation, and inspect inlet and outlet structures. The pond bottom can be dried between production cycles when environmental conditions permit. Drying helps reduce accumulated organic residues and can improve bottom conditions before filling. Liming materials may be applied according to soil characteristics and farm management requirements. After preparation, water is introduced through properly screened inlet structures to reduce the entry of unwanted fish and other organisms.

Water quality management begins before shrimp are stocked and continues throughout the culture period. Temperature, dissolved oxygen, pH, salinity, alkalinity, ammonia, nitrite, and water transparency are among the variables that can influence shrimp performance. Temperature affects metabolic activity and feed consumption, while dissolved oxygen is required for respiration and supports microbial processes within the pond. Low oxygen concentrations can reduce feeding activity and increase physiological stress. Regular measurements allow farmers to identify changes before they become severe.

Pond water contains a complex community of microorganisms, algae, zooplankton, and organic particles. These organisms contribute to the biological productivity of the culture environment. Phytoplankton can contribute oxygen during daylight and provide food for small aquatic organisms. However, excessive algal growth can create substantial changes in oxygen concentration, particularly during nighttime and early morning. Farmers therefore need to observe water color, transparency, oxygen concentration, and other indicators instead of relying on visual appearance alone.

Stocking density is another major factor affecting production. In semi-intensive systems, density is selected according to pond size, water availability, aeration capacity, feed resources, farmer experience, and expected environmental conditions. Excessive stocking can increase competition for oxygen and feed while increasing organic loading. Lower densities may reduce production per unit area but can provide greater environmental stability. Selecting healthy post-larvae from a reliable hatchery is therefore an important step in establishing a productive crop.

Acclimation should be performed carefully when post-larvae are transferred from hatchery conditions to pond water. Differences in temperature, salinity, and other water properties can affect shrimp during transfer. Gradual adjustment allows the animals to adapt to the new environment. Farmers can observe shrimp activity after stocking and monitor survival through sampling and routine pond observations. Early detection of unusual swimming behavior, weak feeding response, or changes in distribution can provide useful information about pond conditions.

Correspondence to: Flávio Leite, Department of Aquatic Systems, Federal University of Santa Catarina, Florianopolis, Brazil, E-mail: flavio.leite@ufsc.br

Received: 29-Jun-2026, Manuscript No. JARD-26-32807; **Editor assigned:** 01-Jul-2026, PreQC No. JARD-26-32807 (PQ); **Reviewed:** 15-Jul-2026, QC No. JARD-26-32807; **Revised:** 22-Jul-2026, Manuscript No. JARD-26-32807 (R); **Published:** 29-Jul-2026, DOI: 10.35248/2155-9546.26.17.1130

Citation: Leite F (2026). Semi-Intensive Shrimp Culture: Production Practices, Water Management, and Sustainable Farm Performance. J Aquac Res Dev. 17:1130.

Copyright: © 2026 Leite F. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Feed represents one of the major operating expenses in shrimp culture. Semi-intensive farms commonly use formulated feeds designed to supply protein, lipids, carbohydrates, vitamins, minerals, and other nutrients. Feeding schedules should consider shrimp size, biomass, water temperature, feeding response, and environmental conditions. Feed trays or sampling methods can help determine whether shrimp are consuming the supplied ration. Excess feed should be avoided because uneaten material decomposes and contributes to organic loading and deterioration of water quality.

As shrimp grow, farmers can estimate biomass through periodic sampling. Sampling provides information about average body weight, size distribution, survival estimates, and growth rate. These measurements can be used to adjust feed quantities and estimate the expected harvest period. Growth can vary between ponds because of differences in water quality, stocking density, feed distribution, weather, and bottom conditions. Maintaining production records enables farmers to compare results between cycles and identify practices associated with better performance.

Aeration can be incorporated into semi-intensive systems when oxygen demand increases. Mechanical aerators increase water movement and improve gas exchange at the water surface. Their use may be especially valuable during periods of high biomass, cloudy weather, excessive algal activity, or elevated nighttime oxygen demand. Aerator operation should be based on measured conditions and farm requirements rather than unnecessary continuous operation. Efficient use of equipment can help control electricity consumption while maintaining an acceptable culture environment.

Disease prevention is an important component of shrimp farming. Disease problems may arise from infectious agents, environmental stress, poor nutrition, unsuitable stocking practices, or interactions among several factors. Farm biosecurity can include controlled access, equipment sanitation, screening of incoming water, removal of dead animals, proper disposal of waste, and careful selection of post-larvae. Farmers should also avoid transferring equipment between ponds without suitable cleaning because equipment can contribute to the movement of pathogens and contaminated material.

Biosecurity is particularly important in areas where multiple shrimp farms operate close together. Water movement between farms can increase the possibility of disease transmission when discharge and intake systems are poorly separated. Maintaining appropriate water exchange practices, using screened water inlets, and monitoring shrimp health can reduce avoidable risks. Farmers should work with aquatic animal health professionals when unusual mortality or disease symptoms are observed.

Sediment management also influences the performance of semi-intensive ponds. Shrimp waste, uneaten feed, dead plankton, and other organic materials gradually accumulate on the pond bottom. Areas receiving repeated feed deposition may develop higher organic loads than other parts of the pond. Proper feeding, water circulation, periodic pond drying, and suitable sludge management can help maintain better bottom conditions. Wastewater should be handled responsibly to reduce effects on surrounding waterways.

Environmental sustainability can be improved through efficient use of feed, water, energy, and other farm inputs. Reducing unnecessary water exchange can lower the volume of discharge while maintaining suitable pond conditions through careful monitoring. Improved feed conversion can reduce production costs and decrease nutrient release. Solar-powered aeration and efficient pumps may also reduce dependence on conventional electricity in suitable locations.

Harvest planning begins well before shrimp reach market size. Farmers monitor growth, survival, market requirements, water conditions, and seasonal factors when deciding the appropriate harvest period. Harvesting can involve partial removal of shrimp or complete draining of the pond. Proper handling after capture is important because shrimp quality can decline rapidly under warm conditions. Rapid cooling, clean handling surfaces, appropriate containers, and hygienic transportation help maintain product quality.

Record keeping supports better decision-making across successive culture cycles. Records can include stocking date, post-larval quantity, feed use, water quality measurements, mortalities, treatments, sampling results, electricity consumption, labor requirements, harvest weight, and financial outcomes. Comparing these records can reveal changes in feed conversion, survival, growth, and profitability. Digital farm records can make this process easier and allow farmers to examine production trends over several cycles.

CONCLUSION

Semi-intensive shrimp culture can provide a practical balance between natural pond productivity and controlled production inputs. Its performance depends less on a single technique and more on consistent management of stocking, nutrition, water quality, animal health, pond conditions, and harvesting. Farmers who maintain regular observations and respond to measured changes can improve production stability while reducing avoidable resource use. Continued development in sensor monitoring, automated feeding, water-quality analysis, disease screening, and environmentally responsible pond management can further improve the efficiency of semi-intensive shrimp production. With appropriate site selection, sound farm hygiene, careful feeding, responsible water management, and accurate production records, semi-intensive culture can remain an important method for producing shrimp for domestic and international markets.