



Soil Sensor Networks for Real-Time Monitoring and Efficient Agricultural Resource Management

Imane Duric *

Department of Agricultural Technology, University of Queensland, Brisbane, Australia

DESCRIPTION

Soil sensor networks are becoming increasingly useful in modern agriculture because they provide continuous information about conditions within the soil. Agricultural fields can contain substantial variation in moisture, temperature, electrical conductivity, nutrient availability, and other physical characteristics. Conventional soil sampling provides valuable measurements, but samples are normally collected at specific locations and times. Sensor networks provide a different approach by allowing repeated observations from multiple points across a field. These observations can help farmers understand how soil conditions change over time and support decisions related to irrigation, crop establishment, nutrient management, and field operations.

A soil sensor network generally consists of several sensing devices positioned at selected locations within an agricultural area. Each sensor measures one or more soil characteristics and transfers the collected information to a central platform or local data unit. Depending on the design, communication may occur through wireless networks, radio systems, cellular connections, or other digital communication technologies. The collected information can then be displayed through computers, tablets, or mobile applications. This arrangement allows users to monitor field conditions without physically visiting every measurement location.

Soil moisture is one of the most commonly monitored variables in sensor-based agricultural systems. Water availability strongly influences seed germination, root development, nutrient movement, and crop growth. Soil moisture sensors can provide measurements at different depths, allowing farmers to observe changes within the root zone. When moisture values fall below an appropriate range, irrigation can be scheduled according to crop requirements. When adequate moisture remains in the soil, irrigation can be delayed. This approach can reduce unnecessary water application and support more efficient use of irrigation resources.

Different types of soil moisture sensors are available for agricultural applications. Some systems estimate water content by measuring electrical properties of the soil, while others use methods based on dielectric characteristics. Sensor selection depends on soil texture, installation depth, expected accuracy, environmental conditions, and available budget. Sensors must also be installed correctly because poor placement can produce measurements that do not represent the surrounding soil. Regular inspection and calibration are useful for maintaining reliable measurements throughout the growing season.

Soil temperature is another important variable that can be monitored through sensor networks. Temperature affects seed germination, microbial activity, root growth, and nutrient transformations. In some crops, soil temperature is an important factor when determining suitable planting conditions. Sensors placed at different depths can show how quickly the soil warms during the day and cools during the night. Continuous temperature records can therefore provide information that is difficult to obtain through occasional manual measurements.

Electrical conductivity is also widely measured in sensor-based soil monitoring. It is influenced by factors such as soil texture, moisture, dissolved salts, and nutrient concentration. Spatial differences in electrical conductivity may indicate changes in soil properties across a field. When conductivity information is combined with soil sampling and geographic mapping, farmers can identify areas that may require further investigation. This information can contribute to management zone development and support more efficient field sampling.

Some advanced sensor systems can measure chemical characteristics such as nitrate concentration or soil acidity. These sensors can provide information about nutrient conditions, although their accuracy may vary according to soil type, moisture status, temperature, and sensor design. Laboratory analysis remains important for confirming soil nutrient levels. Sensor measurements can complement laboratory testing by providing more frequent observations and identifying locations where additional samples may be useful.

Correspondence to: Imane Duric, Department of Agricultural Technology, University of Queensland, Brisbane, Australia, E-mail: imane.duric@agtech.com

Received: 01-Dec-2025, Manuscript No. AGT-25-32508; **Editor assigned:** 03-Dec-2025, Pre QC No. AGT-25-32508 (PQ); **Reviewed:** 17-Dec-2025, QC No. AGT-25-32508; **Revised:** 23-Dec-2025, Manuscript No. AGT-25-32508 (R); **Published:** 31-Dec-2025, DOI:10.35248/2168-9891.25.14.434

Citation: Duric I (2025) Soil Sensor Networks for Real-Time Monitoring and Efficient Agricultural Resource Management. Agrotechnology. 14:434.

Copyright: © 2025 Duric I. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

CONCLUSION

Soil sensor networks provide a practical method for collecting frequent information about agricultural soil conditions. By monitoring moisture, temperature, electrical conductivity, and selected chemical properties, these systems can support irrigation planning, nutrient management, crop monitoring, and resource allocation. Their effectiveness depends on suitable

sensor selection, correct installation, reliable communication, accurate calibration, and appropriate interpretation of the collected information. As sensor costs decrease and digital agricultural systems become more accessible, soil sensor networks can become an increasingly useful component of data-based farm management, supporting efficient production while conserving water, nutrients, energy, and soil resources.