ISSN: 2157-7471
Journal of Plant Pathology & Microbiology

OPEN ACCESS Freely available online

Commentary

Suppression and Spread Dynamics of Bacterial Wilt in Agricultural Systems

Vesna Dragomirovic*

Department of Plant Pathology, University of Novi Sad, Novi Sad, Serbia

DESCRIPTION

Bacterial wilt is a destructive plant disease that affects a wide range of cultivated crops across tropical, subtropical, and warm temperate regions. It is caused mainly by soil- and water-borne bacterial species such as *Ralston solanacearum*, which invades plant vascular tissues and disrupts normal water transport. Once inside the host, the pathogen multiplies rapidly within xylem vessels, leading to blockage of water movement and progressive wilting symptoms even when soil moisture is adequate. The disease is known for its persistence in soil and water environments, making long-term agricultural management highly challenging.

The infection process often begins through natural openings in roots or through wounds caused by farming tools, nematodes, or insect feeding. After entry, bacterial cells colonize the intercellular spaces and later move into vascular tissues. The blockage of xylem vessels is not only due to bacterial proliferation but also due to plant defence responses such as production of gums and teloses, which further restrict water flow. This dual effect results in leaf yellowing, drooping, and eventual collapse of the entire plant. In severe cases, the plant dies rapidly after the first visible symptoms appear.

Environmental conditions strongly influence disease development. Warm temperatures and high soil moisture levels promote bacterial multiplication and movement through the soil water film. Poor drainage, continuous cropping of susceptible hosts, and infested irrigation water significantly increase disease incidence. Infected plant debris left in fields can serve as a long-term reservoir, allowing the pathogen to survive between growing seasons. Additionally, asymptomatic carrier plants may contribute to unnoticed spread across agricultural areas.

Host range is another concern in bacterial wilt management. The pathogen can infect solanaceous crops such as tomato, potato, eggplant, and pepper, as well as banana, peanut, and several ornamental plants. Genetic variation within pathogen populations leads to differences in virulence and host specificity. This variability complicates breeding efforts aimed at developing resistant cultivars, as resistance effective in one region may fail in another due to pathogen diversity.

Management strategies rely on an integrated approach combining cultural, biological, and genetic methods. Crop rotation with non-host species helps reduce soil inoculum levels over time. However, because the bacterium can survive in soil and water for extended periods, rotation alone is often insufficient. Soil polarization and improved drainage systems can reduce bacterial survival and limit disease spread. The use of pathogen-free planting material is essential to prevent introduction into clean fields.

Biological control has gained attention as an environmentally friendly strategy. Certain beneficial microorganisms, including *Bacillus* and *Pseudomonas* species, can suppress bacterial wilt through competition, antibiosis, and induction of plant defence responses. These microbes colonize the rhizosphere and reduce pathogen establishment by occupying ecological niches and producing inhibitory compounds. Organic amendments such as compost and farmyard manure also enhance microbial diversity in soil, indirectly reducing disease pressure.

Plant resistance breeding remains an important long-term solution. Resistant varieties exhibit reduced vascular colonization or enhanced ability to limit bacterial multiplication. Advances in molecular biology have enabled identification of resistance-related genes and quantitative trait loci associated with partial resistance. However, complete immunity is rare, and resistance is often influenced by environmental conditions and pathogen strain variation.

Diagnostic techniques have improved early detection of bacterial wilt in fields and nurseries. Traditional methods based on symptom observation are now complemented by molecular tools such as polymerase chain reaction assays and loop-mediated isothermal amplification. These techniques allow rapid identification of the pathogen even in latent infections, supporting timely intervention measures and reducing further spread.

Correspondence to: Vesna Dragomirovic, Department of Plant Pathology, University of Novi Sad, Novi Sad, Serbia, E-mail: vesna.dragomirovic.pp@uns.edu.rs

Received: 27-Feb-2026, Manuscript No. JPPM-26-31639; **Editor assigned:** 02-Mar-2026, PreQC No. JPPM-26-31639 (PQ); **Reviewed:** 16-Mar-2026, QC No. JPPM-26-31639; **Revised:** 23-Mar-2026, Manuscript No. JPPM-26-31639 (R); **Published:** 30-Mar-2026, DOI: 10.35248/2157-7471.26.17.783

Citation: Dragomirovic V (2026). Suppression and Spread Dynamics of Bacterial Wilt in Agricultural Systems. J Plant Pathol Microbiol. 17:783.

Copyright: © 2026 Dragomirovic V. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Climate change and global trade patterns are expected to influence the distribution of bacterial wilt in the coming decades. Rising temperatures may expand the geographic range of the pathogen, while increased movement of agricultural products can introduce it into new regions. Strengthening quarantine measures and monitoring systems will therefore be important to limit long-distance dissemination. Understanding the ecological behavior of the pathogen within soil and plant systems remains an active area of research. Interactions between microbial communities, soil properties, and host physiology determine disease outcomes in complex ways. Continued investigation into these interactions can support the development of more effective and sustainable management practices.