OPEN ACCESS Freely available online

ISSN: 2155-9546
Journal of Aquaculture Research & Development

Opinion Article

Temporal Patterns of Nutrient Release and Assimilation in Formulated Aquafeeds

Hellen Pirarat*

Department of Aquaculture, Kasetsart University, Bangkok, Thailand

DESCRIPTION

Evaluating the nutritional efficiency of formulated feeds in aquaculture has traditionally relied on static measurements, primarily through the calculation of apparent digestibility coefficients. While these values provide a useful baseline for comparing ingredients, they offer a single endpoint measurement that conceals the dynamic nature of digestion. Digestion is a continuous physiological process that unfolds over time rather than a simple binary outcome. The rate at which dietary components are hydrated, structurally disassembled, enzymatically cleaved, and absorbed across the intestinal epithelium is influenced by a complex interplay of physical, chemical, and biological factors. These factors include the physical architecture of the feed, raw ingredient characteristics, processing conditions, digestive morphology, enzyme kinetics, environmental temperature, host developmental stage, and overall physiological status. Evaluating digestion as a time-dependent kinetic process provides critical insights into how effectively a diet meets the metabolic demands of farmed aquatic species for growth, energy maintenance, and health.

The process of digestion commences immediately upon feed ingestion. As feed enters the gastrointestinal tract, it undergoes rapid hydration while being subjected to mechanical forces, endogenous digestive secretions, enzymatic activity, and interactions with host microbial populations. Complex dietary macromolecules must be systematically broken down into bioavailable components: proteins into short peptides and free amino acids, triacylglycerol's into free fatty acids and monoacylglycerols, and complex carbohydrates into simple monosaccharides. The velocity of these biochemical transformations is strongly dictated by the physical integrity of the feed matrix and the inherent physiological capabilities of the target species. Nutrients housed within soluble or highly porous matrices are rapidly liberated and absorbed, whereas nutrients locked within dense, highly cross-linked protein structures or tough plant cell wall matrices require prolonged residence time and extensive enzymatic activity before absorption can occur. Efficient digestion therefore depends on the synchronization of feed breakdown, enzyme secretion, nutrient absorption, and gastrointestinal transit to maximize nutrient utilization and minimize waste production.

A complete kinetic evaluation of feed breakdown requires tracking multiple parameters throughout the passage of chime through the gastrointestinal tract. Key kinetic indicators include the initial rate of nutrient disappearance, the latency period prior to significant hydrolysis, the time required to achieve peak absorption rates, the total bioavailable fraction, and the mass of indigestible residue remaining in the hindgut. Two distinct aquafeed formulations can possess identical total protein digestibility percentages while exhibiting drastically different temporal release profiles. A rapid, unsustained influx of amino acids into the bloodstream may overwhelm cellular transport systems and exceed the rate of protein synthesis, leading to the catabolic oxidation of excess amino acids for energy rather than their retention in muscle tissue. Conversely, an excessively slow release rate can result in intact protein reaching the distal intestine, promoting unwanted microbial fermentation and increasing nitrogenous waste production.

The physical properties of an aquafeed play an essential role in dictating the rate of hydration and subsequent enzymatic breakdown within the digestive lumen. Characteristics such as particle size, pellet hardness, bulk density, porosity, moisture absorption capacity, and water stability determine how quickly a feed matrix disintegrates inside the gut. Processing technologies, including thermal extrusion, conditioning, pelleting, drying, and fine grinding, alter the microstructural arrangement of ingredients. Reducing raw material particle size increases the surface area available for digestive enzymes, thereby accelerating hydrolysis. However, extremely fine grinding or over-processing can compromise the physical integrity of pellets in water, leading to nutrient leaching before ingestion or negatively altering feed intake behaviors. Feed manufacturers must therefore balance physical stability during handling and feeding with appropriate physical breakdown kinetics inside the gut.

Each major macronutrient class exhibits unique breakdown dynamics influenced by ingredient origin and structural configuration. Protein digestion depends heavily on molecular conformation, tertiary structure, and the presence of heat-

Correspondence to: Hellen Pirarat, Department of Aquaculture, Kasetsart University, Bangkok, Thailand, E-mail: hellen.p@ku.ac.th

Received: 17-Aug-2026, Manuscript No. JARD-26-32983; **Editor assigned:** 20-Aug-2026, PreQC No. JARD-26-32983 (PQ); **Reviewed:** 03-Sep-2026, QC No. JARD-26-32983; **Revised:** 10-Sep-2026, Manuscript No. JARD-26-32983 (R); **Published:** 17-Sep-2026, DOI: 10.35248/2155-9546.26.17.1155

Citation: Pirarat H (2026). Temporal Patterns of Nutrient Release and Assimilation in Formulated Aquafeeds. J Aquac Res Dev. 17.1155.

Copyright: © 2026 Pirarat H. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

induced cross-links or anti-nutritional factors. Ingredients such as fishmeal, terrestrial animal by-product meals, plant protein concentrates, insect meals, and single-cell proteins differ substantially in their susceptibility to pepsin, trypsin, and chymotrypsin.

CONCLUSION

Evaluating aquafeed digestibility as a dynamic kinetic process provides a far more comprehensive picture of nutrient availability than single-point digestibility values alone. Understanding the temporal breakdown and absorption of nutrients enables feed formulators to design diets with greater precision, align ingredient selection with species-specific digestive physiology, optimize feeding management strategies, reduce nutrient losses into the environment, and rigorously assess novel sustainable ingredients for the future of aquaculture. Carbohydrate digestion varies widely among aquatic species based on their evolutionary adaptations. Lipids follow a distinct temporal path because hydrophobic fat molecules must first undergo emulsification. Endogenous bile salts and phospholipids emulsify fats into tiny droplets, dramatically expanding the surface area for pancreatic lipase activity.