

# Effects of Spearmint (*-Carvon*) Oil and Methyl Salicylate Oil Emulsion on Anesthesia of Common Carp (*Cyprinus carpio* L., 1758)

Zahra Roohi\* and Mohammad Reza Imanpoor

Fishery Science, Department of Fishery and Environment, Gorgan University of Agriculture Science and Natural Resources, Gorgan, Iran

## Abstract

The aim of this study was to identify the times of anesthetic induction and recovery in common carp (*Cyprinus carpio*) with average body weights of  $11.59 \pm 1.23$  g, at water temperatures of  $23^\circ\text{C}$  that were exposed to the spearmint (*-Carvon*) oil and methyl salicylate oil emulsion (CMSE). Common carp were placed in 1-L aquaria containing different concentrations of CMSE (263, 395, 526, 658 and 789  $\mu\text{L/L}$ ). The different treatments resulted in average induction and recovery times ranging from  $80.33 \pm 5.13$  s to  $305.67 \pm 21.13$  s and  $87.67 \pm 19.39$  s to  $194.33 \pm 27.09$  s, respectively. Induction and recovery time decreased significantly with increasing of the concentration of CMSE ( $P < 0.05$ ). Opercular rate firstly increased and then slowly decreased with increasing the concentration of anesthetic. Glucose levels were significantly affected by CMSE concentration ( $P < 0.05$ ). Lower levels of plasma glucose after anesthesia and recovery belong to treatment of 526  $\mu\text{L/L}$ . A second experiment was conducted in which common carp were immersed in either 526  $\mu\text{L/L}$  CMSE or 150 mg/L clove powder in order to compare them with each other. Induction was quicker in the CMSE group; however, recovery was quicker in the clove powder group. Also, lower levels of plasma glucose and opercular rate belong to CMSE. No mortality was observed in the study. Applications of CMSE in 526  $\mu\text{L/L}$  level, seems to be suitable on anesthesia of common carp.

**Keywords:** Anesthesia; Induction time; Glucose; Spearmint; Common carp

## Introduction

Fish used in farming or research are exposed to many situations that are stressful or painful to the animals, including handling [1], confinement, netting, weighing, vaccination, blood sampling [2,3], transport [4], and surgical procedures [5,6]. The stress response is an adaptive physiological function that responds to a perceived threat to homeostasis, which in the short term preserves the health and viability of the stressed individual [7]. However, when the stress is excessive in either intensity or duration, the stress response can result in undesirable consequences, such as illness and mortality [8,9]. The use of anesthetics and sedatives may be beneficial in fish transportation by reducing physical activity and fish stress during handling [10-12] and to reduce mortality when exposed to severe and repeated stressors [13]. Though anesthetics can be a valuable tool to ensure animal welfare during these events, these agents can also have unwanted side effects and should therefore be used with caution [14].

Choosing an anesthetic must be attributed to several characteristics including its efficacy, availability; cost-effectiveness; ease of use; nature of the study; and safety for the user including fish, humans and the environment [15-17].

To minimize the effects of stress on fish, investigators worldwide have begun to examine the use of natural products with anesthetic properties. Examinations of chinook salmon (*Oncorhynchus tshawytscha*), rainbow trout (*Oncorhynchus mykiss*), common carp (*Cyprinus carpio*) and persian sturgeon (*Acipenser persicus*) anesthetized with clove oil [6,15,18,19] and silver catfish (*Rhamdia quelen*) anesthetized with essential oil (EO) of *Lippia alba* [20,21], *Ocimum gratissimum* [22], *Aloysia triphylla* [23], and *Hesperozygis ringens* [24] have led to the development of anesthesia protocols in which products extracted from natural sources were employed with efficacy to treat fish.

One of these potential fish anesthetic compounds is an emulsified mixture of two common human food substances: spearmint oil (*-carvone*) and methyl salicylate. Spearmint (*Mentha spicata*) has been consumed by humans for millennia; it has been written into mythological and religious tales, and today mint is the third most popular flavor in the world behind vanilla (*Vanilla planifolia*) and citrus [25]. It is used in cuisine, candy, chewing gum, cosmetics, toothpaste, tobacco products, and pharmaceutical preparations [26]. Carvone is the component of the spearmint oil that gives it a minty fragrance and flavor. Caraway (*Carum carvi*) seeds, dill weed (*Anethum graveolens*), mandarin orange (*Citrus reticulata*), peels, peppermint (*Mentha piperita*), and many fruits contain mixtures of both enantiomers of carvone and related substances that give them their unique fragrances. *I*-Carvone can be biosynthesized from tangerine peels or extracted from spearmint plants [27]. *I*-Carvone's anesthetic effect on the peripheral and central nervous systems include central nervous system depression, antinociceptive effects, sedation, and anticonvulsant-like activity [28-30].

Methyl salicylate can be synthesized by esterifying salicylic acid with methanol, but for centuries it was distilled from twigs of sweet birch (*Betula lenta*) and eastern teaberry (*Gaultheria procumbens*). Methyl salicylate is purported to be a nonsteroidal anti-inflammatory drug with pharmacological effects comparable to salicylic acid [31,32].

\*Corresponding author: Zahra Roohi, Fishery Science, Department of Fishery and Environment, Gorgan University of Agriculture Science and Natural Resources, Gorgan, Iran, Tel: +989106071079; E-mail: roohi26\_jut@yahoo.com

Received January 04, 2014; Accepted January 29, 2014; Published February 07, 2014

Citation: Roohi Z, Imanpoor MR (2014) Effects of Spearmint (*-Carvon*) Oil and Methyl Salicylate Oil Emulsion on Anesthesia of Common Carp (*Cyprinus carpio* L., 1758). J Aquac Res Development 5: 221 doi:10.4172/2155-9546.1000221

Copyright: © 2014 Roohi Z, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Therefore, the aim of this study was to examine the efficacy and determine the optimum concentration of Spearmint (*I*-Carvon) Oil and Methyl Salicylate Oil Emulsion (CMSE) as common carp anesthetic and compared with commonly fish anesthetic clove powder. The physiological response of plasma glucose was also subsequently investigated.

## Materials and Methods

### Animal

These experiments were conducted on common carp with a mean length of  $9.8 \pm 0.6$  cm and weight of  $11.59 \pm 1.23$  g. The experiment was carried out in 2 m<sup>3</sup> tank. All of the fish were starved for 24 h prior to experiment [33]. Experiment was performed out at 23°C water temperature and in triplicate.

#### Immersion anesthetic formulation and preparation

Five different concentrations of CMSE (263, 395, 526, 658 and 789 µL/L) were chosen according to the previous studies. The CMSE is composed of 28.4% *I*-carvone (pmentha- 6,8-dien-2-one), 4% methyl salicylate (methyl-2- hydroxybenzoate), 25% glycerin, and 5% polysorbate 80 in an aqueous solution. The ingredients are blended aggressively until a brilliant white emulsion forms.

### Second experiment

In order to compare CMSE with clove powder, empirical doses of 526 µL/L CMSE and 150 mg/L clove powder used to anesthetize common carp.

### Opercular rate (OpR)

Opercular rate as a function of concentration and time under anesthesia, was recorded during induction of anesthesia and recovery time.

### Sampling and analyses

Blood was collected from the caudal vein of each fish using nonheparinized 3-mL syringes. Plasma was obtained after blood centrifugation (7 min, 5,000 rpm) at room temperature. The collected serum was frozen at -20 °C until plasma glucose analyses, using commercial kits (Pars Azmun Co. Ltd., Tehran, Iran), could be performed.

After adding the anesthetic agent, three fish was stocked into the each aquarium. In this study, induction time, recovery time, opercular rate, plasma glucose was noted and fish were maintained there for 72h in order to observe possible mortality.

### Statistical analysis

Data were statistically analyzed by the SPSS 16 software. One-way analysis of variance (ANOVA) was employed to compare the means of factors. Where significant F-ratios were calculated by ANOVA, the Duncan test was applied to identify which means was different.

## Results

The results obtained from the experiment are shown in Figures 1-3. All fish exposed to 263, 395, 526, 658 and 789 µL/L dose of CMSE at the experiment were anesthetized and recovered from anesthesia. No mortality occurred during the experiments.

In CMSE application, induction time for common carp changed




Figure 1: Induction and recovery time (mean ± SD) of common carp anesthetized with CMSE.




Figure 2: Opercular rate (OpR) at induction and recovery time of fish anesthetized with CMSE.




Figure 3: Glucose level profile of common carp after anesthesia and recovery.

between  $80.33 \pm 5.13$ - $305.67 \pm 21.13$  second (Figure 1). There was significant difference in the induction time at the different concentrations of the CMSE tested ( $P < 0.05$ ). Behavioral anesthesia recovery occurred in  $< 200$ s for all anesthetized fish (Figure 1). There was significant difference in the recovery time at the different concentrations of the CMSE tested ( $P < 0.05$ ).

According to the results, fish anesthetized with 263 µL/L showed a

significant decrease in the induction opercular rate compared to other concentrations of the CMSE ( $P < 0.05$ ). The Highest of the opercular rate at induction and recovery time belong to concentration of 526  $\mu\text{L}$  (Figure 2). There was significant difference in the opercular rate at recovery time ( $P < 0.05$ ).

In the case of plasma glucose after anesthesia and recovery, lower levels belong to treatment of 526  $\mu\text{L}$  (Figure 3). The glucose levels were significantly affected by CMSE concentration ( $P < 0.05$ ). There was significant difference in the glucose levels after anesthesia and recovery under the different concentrations of the CMSE tested.

In the second experiment, fish anesthetized in CMSE reached stage 4 anesthesia in 137 s. Common carp anesthetized in clove powder reached 4 anesthesia in 157 s (Figure 4). However, common carp anesthetized in CMSE, had slower recovery time from stage 4 anesthesia than clove powder (Figure 4).

Common carp anesthetized in CMSE, had lowest opercular rate at induction and recovery time (Figure 5). Also, lower levels of plasma glucose after anesthesia and recovery belong to CMSE (Figure 6).

## Discussion

In fish farming, anesthetics are necessary to lower the level of stress, to prevent any injuries of fish during handling such as particularly in artificial reproduction, marking, biometric evaluation etc [34]. Many factors affect the efficacy of anesthetics in fish. These factors can be divided into biological and environmental categories, and include




Figure 6: Plasma glucose at induction and recovery times for common carp anesthetized to stage 4 anesthesia with CMSE or clove powder.

species, body size, the density of the fish in a bath, and water quality. An ideal anesthetic for fish should induce anesthesia in less than 3 to 5 min, with total loss of equilibrium and muscle tone, should allow an uneventful and rapid (i.e. less than 10 min) recovery, should leave low tissue residues and be safe to users as well as be inexpensive and easy to use [35,36].

In this study, we found CMSE to be an effective immersion fish anesthetic. An increasing concentration of the CMSE proportionally decreased the time required for anesthesia induction (Figure 1). This result is in accordance with previous findings in *Hippocampus reidi* exposed to the essential oil of *Lippia alba* [37] and *Cyprinus carpio* anesthetized with  $\text{NaHCO}_3$  [38]. In our study, recovery time is usually faster at highest concentrations of anesthetic, and recovery time decrease with an increase in the concentration of the CMSE. This our result is consistent with previous findings in russian sturgeon [39] and *Rutilus frissi kutum* [40] anesthetized with clove oil. This can indicate that induction time and recovery time of the fish can vary depending on concentration of anesthetic [36, 41]. Fish anesthetized with 789  $\mu\text{L}$  concentration did not show any significant difference in the recovery time compared to 526 and 658  $\mu\text{L}$  concentrations. So the 526  $\mu\text{L}$  concentration is suggested as the optimal concentration for deep anesthesia induction.

Figure 2 shows the changes in the opercular rates of the common carp during anesthesia and recovery. Fish absorb anesthetic substances mainly through their gills [42]. Opercular rate firstly increased and then slowly decreased with increasing the concentration of anesthetic. This result is in accordance with previous findings in *Cyprinus carpio* anesthetized with  $\text{NaHCO}_3$  [38]. In our study, when fish were exposed to 526  $\mu\text{L}$  of CMSE, highest opercular rates were observed during anesthesia or recovery. Different fish species have different opercular rates, which can reflect upon the rate of anesthetics absorption and upon anesthesia induction.

Smith [43] summarized reports on changes in plasma biochemical parameters due to fish stress. Based on his results, stress may decrease or increase blood glucose levels in different fish 1997; [44-46] In this study, significant differences in the plasma glucose levels after anesthesia and after recovery between the groups were observed ( $P < 0.05$ ). This our result is in accordance with findings of Mousavi [47]. The blood glucose levels of fish anaesthetized with 526  $\mu\text{L}$  decreased significantly compared to other concentrations. Significant changes in glucose levels suggest that anesthesia with 526  $\mu\text{L}$  dose of CMSE would be lower stress response compared to higher and lower doses.




Figure 4: Anesthesia induction and recovery times for common carp anesthetized to stage 4 anesthesia with CMSE or clove powder.




Figure 5: Opercular rate at induction and recovery times for common carp anesthetized to stage 4 anesthesia with CMSE or clove powder.

According to the results obtained from the second experiment, CMSE had a slightly quicker induction time to stage 4 anesthesia than clove powder. The difference in induction time for stage 4 anesthesia between CMSE and MS-222 is not due to differences in fish size. Total body mass and length were normally distributed. However, CMSE had a slower recovery time from stage 4 anesthesia than clove powder (Figure 4). The opercular rate and glucose concentration of common carp was significantly reduced by CMSE and increased by clove powder. These findings demonstrate that CMSE has comparable induction and recovery times (stage 4 anesthesia) to clove powder.

In conclusion, the result of this study indicated that CMSE (526  $\mu$ I/L) can induce safe and valid anesthesia in common carp that was higher (257  $\mu$ I/L) than result of specie atlantic salmon (*Salmo salar sebago*) in study of Russell Danner et al. [48] So, CMSE applications can be a good alternative to other anesthetics.

## References

- Sandodden R, Finstad B, Iversen M (2001) Transport Stress in Atlantic Salmon (*Salmo Salar* L.): Anaesthesia and Recovery. *Aquaculture Research* 32: 87-90.
- Ackerman JL, Bellwood DR (2002) Comparative Efficiency of Clove Oil and Rotenone for Sampling Tropical Reef Fish Assemblages. *Journal of Fish Biology* 60: 893-901.
- Baker DW, Wood AM, Litvak MK, Kieffer JD (2005) Haematology of Juvenile *Acipenser Oxyrinchus* and *Acipenser Brevirostrum* at Rest and Following Forced Activity. *Journal of Fish Biology* 66: 208-221.
- McGovern-Hopkins K, Tamaru CS, Takeshita G, Yamamoto M (2003) Procedural Guide For The Artificial Insemination of The Lyretail Swordtail (*Xiphophorus Helleri*). University of Hawai'i Sea Grant College Program School of Ocean and Earth Science and Technology; Available Online at (Accessed April 2009).
- Kiessling A, Johansson D, Zahl IH, Samuelsen OB (2009) Pharmacokinetics, Plasma Cortisol And Effectiveness of Benzocaine, MS-222 and Isoeugenol Measured in Individual Dorsal Aorta-Cannulated Atlantic Salmon (*Salmo Salar*) Following Bath Administration. *Aquaculture* 286:301-308.
- Wagner GN, Singer TD, Mckinley RS (2003) The Ability of Clove Oil and MS-222 To Minimize Handling Stress In Rainbow Trout (*Oncorhynchus Mykiss* Walbaum). *Aquaculture Research* 34:1139-1146.
- Ashley PJ (2007) Fish Welfare: Current Issues in Aquaculture. *Applied Animal Behaviour Science* 104:199-235.
- Iwama G, Afonso L, Todgham A, Ackerman P, Nakano K (2004) Are Hsps Suitable For Indicating Stressed States In Fish? *The Journal Of Experimental Biology* 204:15-19.
- Barcellos LJG, Kreutz LC, Quevedo RM, Fioreza I, Rodrigues LB, Soso AB, Ritter F, Conrad J, Cericato L, Fagundes M, Terra SR (2003) Haematological And Biochemical Characteristics Of Male Jundia (*Rhamdia Quelen* Quoy & Gaimard Pimelodidae): Changes After Acute Stress. *Aquaculture Research* 34: 1465-1469.
- Rotllant J, Balm PHM, Pérez-Sánchez J, Bonga SEW, Tort L (2001) Pituitary and Interrenal Function in Gilthead Sea Bream (*Sparus Aurata* L., Teleostei) After Handling and Confinement Stress. *General Comp Endocrinol* 121:333-342.
- Skjervold PO, Fjoera SV, Østby PB, Einem O (2001) Livechilling and Crowding Stress before Slaughter of Atlantic Salmon (*Salmo Salar*). *Aquaculture* 192:265-280.
- Mckim JM, Kolanczyk RC, Lien GJ, Hoffman AD (1999) Dynamics of Renal Excretion of Phenol and Major Metabolites in The Rainbow Trout (*Oncorhynchus Mykiss*). *Aquatic Toxicology*. 45: 265-277.
- Strange RJ, Schreck CB (1978) Anesthetic And Handling Stress on Survival And Cortisol Concentration in Yearling Chinook Salmon (*Oncorhynchus Tshawytscha*). *Journal of The Fisheries Research Board of Canada* 35:345-349.
- Zahl IH, Samuelsen O, Kiessling A (2012) Anaesthesia of Farmed Fish: Implications for Welfare. *Fish Physiology and Biochemistry* 38:201-218.
- Cho GK, Heath DD (2000) Comparison of Tricaine Methanesulphonate (MS 222) and Oil Clove Anaesthesia Effects on The Physiology of Juvenile Chinook Salmon (*Oncorhynchus Tshawytscha*) (Walbaum). *Aquaculture Research* 31:537-546.
- Iversen M, Finstad B, Mckinley RS, Eliassen RA (2003) The Efficacy Of Metomidate, Clove Oil, AQUI-Sk And Benzoakr As Anesthetics In Atlantic Salmon (*Salmo Salar* L.) Smolts And Their Potential Stress-Reducing Capacity. *Aquaculture* 221: 549-566.
- Mylonas CC, Cardinaletti G, Sigelaki I, Polzonetti-Magni A (2005) Comparative Efficacy of Clove Oil and 2-Phenoxyethanol as Anesthetics in the Aquaculture of European Sea Bass (*Dicentrarchus Labrax*) and Gilthead Sea Bream (*Sparus Aurata*) at Different Temperatures. *Aquaculture* 246: 467-481.
- Velisek J, Svobodova Z, Plackova V, Groch L, Nepechalova L (2005) Effects of Clove Oil Anaesthesia on Common Carp (*Cyprinus Carpio* L.). *Veterinary Medicine Czech* 50:269-275.
- Imanpoor MR, Bagheri T, Hedayati SAA (2010) The Anesthetic Effects of Clove Essence in Persian Sturgeon, *Acipenser Persicus*. *World Journal of Fish and Marine Sciences* 2: 29-36.
- Cunha MA, Barros FMC, Garcia LO, Veeck APL, Heinzmann BM, Loro VL, Emanuelli T, Baldissarotto B (2010) Essential Oil of *Lippia Alba*: A New Anesthetic For Silver Catfish, *Rhamdia Quelen*. *Aquaculture* 306:403-406.
- Heldwein CG, Silva LL, Reckziegel P, Barros FMC, Bürger ME, Baldissarotto B, Malmann CA, Schmidt D, Caron BO, Heinzmann BM (2012) Participation of Gabaergic System In The Anesthetic Effect of The Essential Oil Of *Lippia Alba* (Mill.) N.E. Brown. *Brazilian Journal of Medical and Biological Research* 45:436-443.
- Silva LL, Parodi TV, Reckziegel P, Garcia VO, Bürger ME, Baldissarotto B, Malmann CA, Pereira AMS, Heinzmann BM (2012) Essential Oil of *Ocimum Gratissimum* L.: Anesthetic Effect, Mechanism of Action and Tolerance in Silver Catfish. *Rhamdia Quelen Aquaculture* 350-353:91-97.
- Gressler LT, Riffel APK, Parodi TV, Saccol EMH, Koakoski G, Costa ST, Pavanato MA, Heinzmann BM, Caron B, Schmidt D, Liesuy SF, Barcellos LJG, Baldissarotto B (2012) Silver Catfish *Rhamdia Quelen* Immersion Anaesthesia With Essential Oil of *Aloysia Triphylla* (L'Hérit) Britton Or Tricaine Methanesulfonate: Effect On Stress Response And Antioxidant Status. *Aquaculture Research*. Doi:10. 1111/Are.12043.
- Silva LL, Silva DT, Garlet QI, Cunha MA, Malmann CA, Baldissarotto B, Longhi SJ, Pereira AMS, Heinzmann BM (2013) Anesthetic Activity of Brazilian Native Plants in Silver Catfish (*Rhamdia Quelen*). *Neotropical Ichthyology* 11: 443-451.
- Hayes JR, Stavanja MS, Lawrence BM (2007) Biological And Toxicological Properties Of Mint Oils And Their Major Isolates: Safety Assessment. Pages 422-491 in Lawrence BM, Editor. *Mint: The Genus Mentha*. Boca Raton, Florida: CRC Press, Taylor & Francis Group.
- Chen C, Isabelle LM, Pickworth WB, Pankow JF (2010) Levels Of Mint And Wintergreen Flavorants: Smokeless Tobacco Products Vs. Confectionery Products. *Food And Chemical Toxicology* 48:755-763.
- De Carvalho CCCR, Da Fonseca MMR (2006) Carvone: Why and How Should One Bother To Produce This Terpene. *Food Chemistry* 95:413-422.
- De Sousa DP, De Farias Nobrega FF, De Almeida RN (2007) Influence of The Chirality of (R)-(2)-A and (S)-(+)-Carvone in The Central Nervous System: A Comparative Study. *Chirality* 19: 264-268.
- Goncalves JC, Oliveira Fde S, Benedito RB, De Sousa DP, De Almeida RN, De Araujo DA (2008) Antinociceptive Activity of (2)-Carvone: Evidence of Association With Decreased Peripheral Nerve Excitability. *Biological And Pharmaceutical Bulletin* 31:1017-1020.
- Oliveira FDS, De Sousa DP, De Almeida RN (2008) Antinociceptive Effect of Hydroxydihydrocarvone. *Biological And Pharmaceutical Bulletin* 31:588-591.
- Stevens SG, Warren B (1964) Determination of Methyl Salicylate In Pharmaceutical Preparations. *Journal of Pharmacy and Pharmacology* 16:SUPPL:32-34T.
- Warren DE, Matsumoto S, Roessig JM, Cech JJ (2004) Cortisol Response of Green Sturgeon to Acid-Infusion Stress. *Comparative Biochemistry and Physiology. A. Physiology* 137: 611-618.
- Weyl O, Kaiser H, Hecht T (1996) On The Efficacy And Mode of Action Of 2-Phenoxyethanol as an Anaesthetic for Goldfish, *Carassius Auratus* (L.), At Different Temperatures and Concentrations. *Aquaculture Research* 27: 757-764.

34. Hamackova J, Kouril J, Kozak P, Stupka Z (2006) Clove Oil As An Anaesthetic For Different Freshwater Fish Species. Bulgarian Journal Of Agriculture Science 12: 185-194.
35. Gilderhus PA, Marking LL (1987) Comparative Efficacy of 16 Anaesthetic Chemicals in Rainbow Trout. North American Journal of Fisheries Management 7: 288-292.
36. Hseu JR, Yeh SL, Chu YT, Ting YY (1998) Comparison of Efficacy of Five Anesthetic in Goldlined Sea Bream, *Sparus Sarba*. Acta Zoologica Taiwanica 9: 35-41.
37. Da Cunha MA, Da Silva BF, Cariello Delunado FA, Cristina Benovit S, De Cavalho Gomes L, Marita Heinzmann B, Baldisserotto B (2011) Anesthetic Induction and Recovery of *Hippocampus Reidi* Exposed to The Essential Oil *Lippia Alba*. Neotropical Ichthyology 9: 683-688.
38. Altun T, Bilgin R, Danabas D (2009) Effects of Sodium Bicarbonate on Anaesthesia of Common Carp (*Cyprinus Carpio* L., 1758) Juveniles. Turkish Journal of Fisheries and Aquatic Sciences 9: 29-31.
39. Akbulut B, Cakmak E, Aksungure N, Cavdar Y (2011) Effect Of Exposure Duration on Time To Recovery From Anaesthesia of Clove Oil In Juvenile of Russian Sturgeon. Turkish Journal of Fisheries and Aquatic Sciences 11: 463-467.
40. Javahery S, Haji Moradlu A, Ghornani R (2012) Efficacy of Clove Oil as an Anaesthetic Fo Two Size Of *Rutilus Frisii Kutum*. Global Veterinaria 9: 319-322.
41. Booke HE, Hollender B, Lutterbie G (1978) Sodium Bicarbonate, An Inexpensive Fish Anesthetic For Field Use. Progressive Fish Culturist 40: 11-13.
42. Ferreira JT, Smith GL, Schoonbee HJ (1984) The Effect of The Anaesthetic Benzocaine-Hydrochloride on Freshwater Quality. Water SA 5: 123-127.
43. Smith LS (1982) Introduction of Fish Physiology. TFH Publications, 352
44. Biron M, Benfey TJ (1994) Cortisol, Glucose And Hematocrit Changes During Acute Stress, Cohort Sampling, And The Diel Cycle In Diploid And Triploid Brook Trout (*Salvelinus Fontinalis* Mitchell). Fish Physiology And Biochemistry 13: 153-160.
45. Wendelaar Bonga SE (1997) The Stress Response in Fish. Physiological Reviews 77: 591-625.
46. Barcellos L, Kreutz L, Quevedo R (2006) Previous Chronic Stress Does Not Alter The Cortisol Response To An Additional Acute Stressor In Jundiá (*Rhamdia Quelen* Quoy and Gaimard) Fingerlings. Aquaculture 253:317-321.
47. Mousavi SM, Majidi Nasab E, Yavari V, Ranjbarzadeh Ghatrami E, Jalili MR (2012) Effects of Two Anaesthetic Regimes, MS-222 And Eugenol, on Plasma Biochemical Profile In *Barbus Sharpeyi*. Comparative Clinical Pathology 21: 859-863.
48. Russell Danner G, Muto KW, Zieba AM, Stillman CM, Seggio JA, Taiq Ahmad S (2011) Spearmint (*l*-Carvone) Oil and Wintergreen (Methyl Salicylate) Oil Emulsion is an Effective Immersion Anesthetic of Fishes. Journal of Fish and Wildlife Management 2: 146-155.