

The Flavonoids in *Citrus madurensis* Lour and their Anti-Hepatitis B Virus Activity

Pei-Hsin Shie¹, Ray-Ling Huang² and Horng-Liang Lay^{1*}

¹Department of Plant Industry, National Pingtung University of Science and Technology, Pingtung County, Taiwan

²Department of Biological Science and Technology, Meiho University, Pingtung County, Taiwan

Abstract

In this study, we evaluated anti-hepatitis B virus (HBV) activity of the extracts (dichloromethane, ethyl acetate, *n*-butanol, acetone, and methanol respectively) of fruit (peels and pulp) of *Citrus madurensis* Lour. (calamondin) using HBV transected cell line MS-G2. The ethyl acetate extracts from the peels strongly reduced the HBsAg expression of HBV in dose 50 µg/mL. According to the previous result, we chose this extract to elute by column chromatography and then separately eluted into 7 fractions. In dose in 50 µg/mL, fractions 2, 3, and 4 had lower HBsAg expression (compare with control). Furthermore, fraction 3 had a strongest inhibitory ability. The flavonoids (hesperidin, diosmin, neohesperidin, nobiletin, tangeretin, and 5-hydroxy-3',4',6,7,8-pentamethoxyflavone) were analysis in each fractions by high-performance liquid chromatography (HPLC). The results shown fraction 2, 3, 4 were detected containing nobiletin, tangeretin, and 5-hydroxy-3',4',6,7,8-pentamethoxyflavone. The anti-HBV activity of the flavonoids show that nobiletin, tangeretin, and also had a lower HBsAg expression. The IC₅₀ value were nobiletin (33.9 µM), tangeretin (20.7 µM), and 5F (5.12 µM), respectively. Taken together, we suggest that 5F could be used as a standard marker for the anti-HBV effect of calamondin.

Keywords: *Citrus madurensis* Lour.; Citrus flavonoids; Hepatitis B virus; High performance liquid chromatography

Introduction

Citrus madurensis Lour. (calamondin), a perennial tree in the family Rutaceae, is a important citrus tree in Taiwan where its juice and fruit are popular among consumers. However, juice production often results in a considerable amount of waste, such as peels, seeds and pulp; fortunately, the waste are rich in flavonoids, carotenoids, polyphenols, and limonoids compounds which all have excellent bioactivities [1-10] and the development of these compounds could create additional value.

The human hepatitis B virus (HBV) is a global disease that is especially prevalent in Asia and Africa. Infections could induce acute hepatitis that could cause the infected into chronic hepatitis carriers. Approximately one-quarter to one-third of those infected will develop early liver cirrhosis and are at elevated risk of hepatocellular carcinoma [11,12]. Current treatments include Interferon- α , Lamivudine, Adefovir, Entecavir, and Telbivudine, but these drugs can only reduce the activity or temporally inhibit the reproduction of HBV and cannot eradicate HBV from the human body. Therefore, in recent years, developing antiviral drugs from natural products has become an important research topic.

Research has revealed that flavonoids in citrus demonstrated antiviral activities [13-17], but little research has been conducted on the anti-HBV properties of citrus. So far the only citrus anti-HBV research consisted of isolated imperatorin purified from pomelo (*C. grandis* (L.) Osbeck) peels, in which findings displayed that imperatorin possessed excellent inhibition effect on the surface antigen and e-antigen of HBV [18]. The purpose of this study is to investigate the anti-HBV activity of calamondin and citrus flavonoids. We used HBV transected cell line MS-G2 to assess that anti-HBV ability.

Materials and Methods

Plant material

Calamondin fruit were bought at random on 9. Oct. 2009 at Jiou-

Ju Township, Pingtung County, Taiwan, and verified by Professor Chung-Ruey Yen. Voucher specimens (Specimen No. CM2009001) were maintained in the laboratory of the corresponding author in the Department of Plant Industry, NPUST.

Crude extracts preparation from calamondin

The peel and pulp were separate from of fruit, that were freeze-dried and powdered, then the dried material was extracted by ultrasonic extraction for 30 min. The extraction was carried out using organic solvents (dichloromethane, ethyl acetate, *n*-butanol, acetone, and methanol respectively). The filtrate was concentrated by evaporation under reduced pressure (ca. 42°C) and further freeze-dried. The dried extract powder was used in anti-HBV assays.

The better anti-HBV activities were evaluated from the different organic solvent extracts described above, and the one with the lowest HBsAg expression was chosen for column chromatography to clarify the bioactive fractions. 200 grams of dried sample powder were extracted with ethyl acetate (2L \times 3) then filtered. The filtrate was concentrated and freeze-dried to yield the ethyl acetate extract (13.8 g).

The ethyl acetate extract was chromatographed by a silica gel column (3.5 i.d. \times 30 cm) and eluted step wise with *n*-hexane \rightarrow 80% *n*-hexane-20% ethyl acetate \rightarrow 60% *n*-hexane-40% ethyl acetate \rightarrow 40% *n*-hexane-60% ethyl acetate \rightarrow 20% *n*-hexane-80% ethyl acetate \rightarrow ethyl acetate \rightarrow methanol.

***Corresponding author:** Horng-Liang Lay, Department of Plant Industry, National Pingtung University of Science and Technology, 1, Shue Fu Rd. Neipu, Pingtung, 91201, Taiwan, Tel: 886-8-7740365; Fax: 886-8-7740415; E-mail: layhl@mail.npust.edu.tw

Received January 09, 2013; Accepted May 15, 2013; Published May 20, 2013

Citation: Shie PH, Huang RL, Lay HL (2013) The Flavonoids in *Citrus madurensis* Lour and their Anti-Hepatitis B Virus Activity. Pharm Anal Acta 4: 239. doi:10.4172/2153-2435.1000239

Copyright: © 2013 Shie PH, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citrus flavanoids of calamondin

Pure compounds of hesperidin, diosmin, neohesperidin, and synephrine were acquired from Sigama Chemical Co.; tangeretin (1), nobiletin (2), and 5-hydroxy-3',4',6,7,8-pentamethoxyflavone (3) were purified from fruit peels of calamondin.

Dry fruit peels were extracted with methanol. After evaporation of the solvent, the residue was partitioned between water and ethyl acetate. The ethyl acetate extract was chromatographed on a silica gel column and eluted with gradients of *n*-hexane: ethyl acetate to yield compound 1~3. The physical and ¹H NMR spectroscopic data of 1~3 were in agreement with the reported data for tangeretin, nobiletin, and 5-hydroxy-3',4',6,7,8-pentamethoxyflavone [19-21]. Using osthole as a positive controls that origin, isolation and structure identification to refer publication [22].

Cell culture and measurement of HBsAg

The human HBV transected cell line MS-G2 was cultured in Dulbecco's modified Eagle medium (DMEM) containing 10% (v/v) FBS (fetal bovine serum), 2.5 µg/mL amphotericin B, 1% penicillin streptomycin solution, and 0.1 µM MEM (modified eagle medium) non-essential amino acid solution at 37°C under 5% CO₂. Cell were seeded in 24-well culture plates at a density of 2×10⁵ cells/mL per well. Cells were properly attached after overnight and changed the test samples, which were all dissolved in DMSO. The concentration of DMSO in the culture medium was kept below 2.5 µL/mL to ensure it did not affect cell growth. The culture media were collected on Day 3 for the measurement of hepatitis B surface antigen (HBsAg). The HBsAg expression in the culture medium was measured using ELISA kits (General Biologicals, Co., Hsinchu, Taiwan) following the Manufacturer's recommendations. Each experiment was performed in triplicate. The expression was calculated as follows:

$$\text{Expression (\%)} = [\text{OD}_{540}(\text{sample}) / \text{OD}_{540}(\text{control})] \times 100\%$$

Cell viability assay

Cytotoxicity was assessed by the MTT assay as previously described [23]. Cells were cultured for 48h with tested compounds. The cells with media alone were used as a negative control. After 48 hours, MTT (1 mg/mL) reagent was added to each well. After 4 hours at 37°C, the medium was then replaced by 400 µL of DMSO and mixed thoroughly to dissolve the formazan crystals. Absorbance was measured at 540 nm. Cell viability was determined as a percentage of the negative control.

$$\text{Cell viability (\%)} = [\text{OD}_{540}(\text{sample}) / \text{OD}_{540}(\text{control})] \times 100\%$$

Cell viability above 80% can be regarded as indicating nontoxicity, and the pharmacy and dose can then be used for investigating HBsAg expression in advance.

HPLC analysis

High Pressure Liquid Chromatography (HPLC) was conducted with a Hitachi system equipped with a degasser DG-2410, pump L-7100, UV/Vis detector L-7420, and auto-sampler L-7200. Peak areas were calculated with D-7000 HSM software.

The samples were eluted using an Inertsil ODS-2 column (4.6 mm i.d.×250 mm) at 40°C. Elution was carried out on a gradient. Mobile phases were 10% (solvent A) and 70% (solvent B) acetonitrile (85% H₃PO₄ was added to adjust the pH value to 2.8). The gradient program was started with 100 : 0 (A : B), and A was gradually decreased to 90% after 10 min, 84% after 55 min, 51% after 65 min where it settled for 31

min, and then to 30% at 105 min and to 0% at 110 min. Re-equilibration of the column was achieved with a linear gradient to 100% A (initial condition) in 5 min, followed by 3 min of isocratic elution before the next injection. The flow rate was 1.0 mL/min, injected volume was 20 µL, and the components were detected at 220 nm.

Statistical analysis

All the experiments were carried out in triplicate and each experiment was repeated three times. The data were analyzed by SAS (Statistical Analysis System). One-way ANOVA with Duncan's multiple range test was used to determine the significance at *p*<0.05.

Results

Effect of solvent extracts on HBsAg expression

To investigate the effect of calamondin on HBsAg expression, five solvents to extract the peel and pulp of calamondin in respectively. The ethyl acetate and acetone extracts from the peels reduced the HBsAg expression of HBV by 41.6% and 71.4%, respectively, in doses 50 µg/mL (Figure 1). The ethyl acetate extract exhibited a high inhibitory effect. In pulp, the *n*-butanol extracts light reduced HBsAg expression to 78.7%, four other solvents extract had no inhibitory effect on the HBsAg of HBV (expression were higher than 80%). The solvent control had no inhibition effect on the HBsAg of HBV.

Effect of different fractions from ethyl acetate extract on HBsAg expression

According to the previous result, the extracts of ethyl acetate from the peels had the highest inhibitory effect on HBsAg expression. We chose this extract to elute by column chromatography and then separately eluted into 7 fractions. The dose at 50 µg/mL, fractions 2, 3, and 4 had lower HBsAg expression (Figure 2). Furthermore, fraction 3 had a strongest inhibitory effect (HBsAg expression 40.4%) then fraction 2 and 4 (HBsAg expression 70.6% and 52.5%, respectively).




Figure 1: Effects of different solvent extracts of each part of calamondin on the HBsAg expression of HBV in (A) peel, and (B) pulp. Values are expressed as mean ± S.D. (percentage of untreated cells) of three experiments.

Effect of flavonoids of calamondin HBsAg expression

The effect on the HBsAg expression of HBV of citrus flavonoids was investigated with results shown in Figure 3. The results shown nobiletin, tangeretin, and 5F had lower HBsAg expression than other compounds. 100 and 50 μM doses of nobiletin lowered the HBsAg expression to 19.3% and 35.1%, respectively, and had a high inhibitory effect. Meanwhile, a 25 μM dose of nobiletin could still lower the HBsAg expression to 56.0% and had a medium inhibitory effect. 100, 50, and 25 μM doses of tangeretin greatly lowered the HBsAg expression to 33.1%, 38.4%, and 41.5%, respectively, and had a very high inhibitory effect. A 6.25 μM dose of 5F lowered the HBsAg expression to 42.1% and had a very strong inhibitory effect, while a 3.13 μM dose of 5F still lowered the HBsAg expression to 64.33% and had a medium inhibitory effect. The IC_{50} of nobiletin, tangeretin, and 5F were calculated. The IC_{50} of 5F was 5.12 μM and 5F had lowest inhibiting concentration followed by tangeretin (20.7 μM) and nobiletin (33.9 μM). The positive control

(osthole) have medium inhibition of HBsAg expression with 48.7% expression at 80 μM .

Quantitative analysis and fingerprint variation of different fractions

Ethyl acetate extracts of peels of calamondin were divided into 7 fractions by column chromatography elution. The fingerprint variation of each fraction was investigated using HPLC and the marker components were quantitatively analyzed, with results shown in Table 1 and Figure 4. In fraction 1, none of the 6 marker components were detected. In fraction 2, only tangeretin and 5F were detected at 47.5 and 26.0 mg/g D.W., respectively. In fraction 3, only nobiletin and tangeretin were detected at 291.2 and 45.0 mg/g D.W., respectively. In fractions 4, 5 and 6, only nobiletin was detected at 420.9, 2.2, and 2.2 mg/g D.W., respectively. In fraction 7, synephrine (without chromatogram), hesperidin, diosmin, and neohesperidin were detected at 12.8, 16.4, 3.9, and 16.7 mg/g D.W., respectively.

Discussion

Organic solvents with different polarities could induce different expressions of activity in their extracts [24-26]. Calamondin methanol extracts had no significant inhibitory effect on the HBV surface antigen or e-antigen in the pre-test of this study. Therefore, dichloromethane, *n*-butanol, ethyl acetate, acetone, and methanol were chosen for the extraction of calamondin in accordance with their polarities. Results exhibited a 50 $\mu\text{g}/\text{mL}$ dose of ethyl acetate extracts of calamondin peel significantly lowered the HBsAg expression of HBV. Meanwhile, MTT assay of the ethyl acetate peel extract displayed no toxicity to normal cells. Hence, organic solvents extracted in different polarities apparently can enhance expression activity and massively elute effective fractions.

According to effective extraction methods previously assessed with different solvents and the tracing of the effective fractions, higher activity expressions were expected. The ethyl acetate peel extracts of calamondin were analyzed using column chromatography and were divided into several fractions. Each fraction was processed using the HBV activity assays. Results showed fractions 2, 3, and 4 significantly lowered the HBsAg expressions of HBV without cytotoxicity with doses of 100, 50, and 50 $\mu\text{g}/\text{mL}$, respectively, but these dosages induced cytotoxicity on normal liver cells. It may be that the components which exhibited anti-HBV effects as well as induced cytotoxicity on normal cells were condensed in the column chromatography process; thus, enhancing their anti-virus activities and rendering them from toxic to normal cells.

Imperatorin, a constituent of the fruit peel of shaddock (*C. maxima* (Burm. f.) Merr. form. Buntan (Hay.) Hort.) and its biotransformation from fermentation using *Aspergillus flavus* displayed excellent anti HBV surface antigens [17]. The three polymethoxyflavones (PMFs) investigated here (nobiletin, tangeretin, and 5F) also had excellent inhibitory effects on the HBV surface antigen, and were higher more effective than imperatorin [17]. Although a 100 μM dose of 5F had lower inhibitory effect on the HBV surface antigen when compared to nobiletin or tangeretin, there was still a high inhibitory effect when the dose was reduced to 6.25 μM .

PMFs are a kind of flavonoid, a specific component of citrus, and they display anti-inflammatory, anti-mutation, and anti-tumor activities [27-30]. This study showed that nobiletin, tangeretin, and 5F all have excellent HBV surface antigen inhibitory effects. Real-time PCR tests will be needed to confirm that the effective compounds that were filtered out in this study could reduce HBV DNA. Further




Figure 2: Effects of different fractions (Fr.) from ethyl acetate extracts in peel of calamondin on the HBsAg expression of HBV. Values are expressed as mean \pm S.D. (percentage of untreated cells) of three experiments.





Figure 3: HPLC fingerprint of ethyl acetate extracts from peel of calamondin for anti-HBV activity. 1. hesperidin; 2. diosmin; 3. neohesperidin; 4. nobiletin; 5. tangeretin; 6. 5-hydroxy-3',4',6,7,8-pentamethoxyflavone.



investigation could examine the influence of intracellular molecules on HBV inhibition mechanism.

Conclusion

In this study, ethyl acetate peel extracts of calamondin have excellent inhibitory effects on HBsAg, and extracts of effective fractions were traced using column chromatography. Although these fractions could effectively reduce the HBsAg expression, they induced cytotoxicity in normal liver cells. Therefore, toxicity in host cells must be considered when using crude extracts to process anti HBV activities. Moreover, after processing anti hepatitis B virus activities using all the marker components in this study, three polymethoxyflavones, namely nobiletin, tangeretin, and 5-hydroxy-6,7,8,3',4'-pentamethoxyflavone, were found to significantly inhibit the hepatitis B virus surface antigen, with 5-hydroxy-6,7,8,3',4'-pentamethoxyflavone showing the strongest hepatitis B virus inhibitory effect. The results of this study could provide a reference for the development of hepatitis B treatments and diversify the use of citrus.

References

- Vikram A, Jayaprakasha GK, Patil BS (2007) Simultaneous determination of citrus limonoid aglycones and glucosides by high performance liquid chromatography. *Anal Chim Acta* 590: 180-186.
- Wang YC, Chuang YC, Ku YH (2007) Quantitation of bioactive compounds in citrus fruits cultivated in Taiwan. *Food Chem* 102: 1163-1171.
- Wang YC, Chuang YC, Hsu HW (2008) The flavonoid, carotenoid and pectin content in peels of citrus cultivated in Taiwan. *Food Chem* 106: 277-284.
- Hertog MG, Feskens EJ, Hollman PC, Katan MB, Kromhout D (1993) Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen Elderly Study. *Lancet* 342: 1007-1011.
- Miyagi Y, Om AS, Chee KM, Bennink MR (2000) Inhibition of azoxymethane-induced colon cancer by orange juice. *Nutr Cancer* 36: 224-229.
- Poulose SM, Harris ED, Patil BS (2005) Citrus limonoids induce apoptosis in human neuroblastoma cells and have radical scavenging activity. *J Nutr* 135: 870-877.
- Poulose SM, Harris ED, Patil BS (2006) Antiproliferative effects of citrus limonoids against human neuroblastoma and colonic adenocarcinoma cells. *Nutr Cancer* 56: 103-112.
- So FV, Guthrie N, Chambers AF, Moussa M, Carroll KK (1996) Inhibition of human breast cancer cell proliferation and delay of mammary tumorigenesis by flavonoids and citrus juices. *Nutr Cancer* 26: 167-181.
- Vanamala J, Leonardi T, Patil BS, Taddeo SS, Murphy ME, et al. (2006) Suppression of colon carcinogenesis by bioactive compounds in grapefruit. *Carcinogenesis* 27: 1257-1265.
- Vanamala J, Reddivari L, Yoo KS, Pike LM, Patil BS (2006) Variation in the content of bioactive flavonoids in different brands of orange and grapefruit juices. *J Food Compos Anal* 19: 157-166.
- Beasley RP, Hwang LY, Lin CC, Chien CS (1981) Hepatocellular carcinoma and hepatitis B virus. A prospective study of 22 707 men in Taiwan. *Lancet* 2: 1129-1133.
- Davey S (1996) State of the world's vaccines and immunization. Geneva: World Health Organization 76-82.
- Kaul TN, Middleton E Jr, Ogra PL (1985) Antiviral effect of flavonoids on human viruses. *J Med Virol* 15: 71-79.
- Rajnarayana KM, Reddy MS, Chaluvadi MR, Krishna DR (2001) Bioflavonoids classification, pharmacological, biochemical effects and therapeutic potential. *Indian J Pharmacol* 33: 2-16.
- De Meyer N, Haemers A, Mishra L, Pandey HK, Pieters LA, et al. (1991) 4'-Hydroxy-3-methoxyflavones with potent anticoronavirus activity. *J Med Chem* 34: 736-746.
- Burnham WS, Sidwell RW, Tolman RL, Stout MG (1972) Synthesis and antiviral activity of 4'-hydroxy-5,6,7,8-tetramethoxyflavone. *J Med Chem* 15: 1075-1076.
- Fortin H, Vigor C, Lohézic-Le Dévéhat F, Robin V, Le Bossé B, et al. (2002) In vitro antiviral activity of thirty-six plants from La Réunion Island. *Fitoterapia* 73: 346-350.
- Teng WY, Huang YL, Huang RL, Chung RS, Chen CC (2004) Biotransformation of imperatorin by *Aspergillus flavus*. *J Nat Prod* 67: 1014-1017.
- Kawaii S, Tomono Y, Katase E, Ogawa K, Yano M (1999) HL-60 differentiating activity and flavonoid content of the readily extractable fraction prepared from citrus juices. *J Agric Food Chem* 47: 128-135.
- Han S, Kim HM, Lee JM, Mok SY, Lee S (2010) Isolation and identification of polymethoxyflavones from the hybrid Citrus, hallabong. *J Agric Food Chem* 58: 9488-9491.
- Li S, Pan MH, Lai CS, Lo CY, Dushenkov S, et al. (2007) Isolation and syntheses of polymethoxyflavones and hydroxylated polymethoxyflavones as inhibitors of HL-60 cell lines. *Bioorg Med Chem* 15: 3381-3389.
- Huang RL, Chen CC, Huang YL, Hsieh DJ, Hu CP, et al. (1996) Osthole increases glycosylation of hepatitis B surface antigen and suppresses the secretion of hepatitis B virus in vitro. *Hepatology* 24: 508-515.
- Huang RL, Chen CC, Chang YS, Chen CF, Chang CG, et al. (2000) Anti-human hepatitis B virus activities of Chinese medicinal prescriptions for the treating for liver disease-anti-HBV effects of six traditional Chinese prescriptions. *J Chin Med* 11: 1-8.
- Anagnostopoulou MA, Kefalas P, Papageorgiou VP, Assimopoulou AN, Boskou D (2006) Radical scavenging activity of various extracts and fractions of sweet orange peel (*Citrus sinensis*). *Food Chem* 94: 19-25.
- Mohana DC, Raveesha KA (2007) Anti-fungal evaluation of some plant extracts against some plant pathogenic field and storage fungi. *J Agric Technol* 4: 119-137.
- Hirata T, Fujii M, Akita K, Yanaka N, Ogawa K, et al. (2009) Identification and physiological evaluation of the components from citrus fruits as potential drugs for anti-obesity and anticancer. *Bioorg Med Chem* 17: 25-28.
- Pan MH, Chen WJ, Lin-Shiau SY, Ho CT, Lin JK (2002) Tangeretin induces cell-cycle G1 arrest through inhibiting cyclin-dependent kinases 2 and 4 activities as well as elevating Cdk inhibitors p21 and p27 in human colorectal carcinoma cells. *Carcinogenesis* 23: 1677-1684.
- Miyazawa M, Okuno Y, Fukuyama M, Nakamura S, Kosaka H (1999) Antimutagenic activity of polymethoxyflavonoids from *Citrus aurantium*. *J Agric Food Chem* 47: 5239-5244.

29. Bas E, Recio MC, Giner RM, Máñez S, Cerdá-Nicolás M, et al. (2006) Anti-inflammatory activity of 5-O-demethylnobiletin, a polymethoxyflavone isolated from *Sideritis tragoriganum*. *Planta Med* 72: 136-142.
30. Imada K, Lin N, Liu C, Lu A, Chen W, et al. (2008) Nobiletin, a citrus polymethoxy flavonoid, suppresses gene expression and production of aggrecanases-1 and -2 in collagen-induced arthritic mice. *Biochem Biophys Res Commun* 373: 181-185.