

The Impact of Pharmacist Led Vancomycin Order Set Implementation in a Computerized-Prescriber-Order-Entry (CPOE) System at a Tertiary Care Centre: A Quasi Experimental Study

Nader Damfu, Mohammed Aseeri*, Allahna Davis, Hani Hasan and Sherine Ismail

King Abdul Allah Medical Research Center/ King Saud bin Abdul Aziz University for Health Sciences, National Guard Health Affairs, Jeddah, Saudi Arabia

Abstract

Introduction: Our hospital has developed a vancomycin protocol in order to improve vancomycin dosing and prevent improper sampling time of vancomycin levels and treatment failure. The study objective is to evaluate the impact of implementing a vancomycin order set integrated into a Computerized-Prescriber-Order-Entry (CPOE) system by pharmacist in adult surgical wards to optimize vancomycin dosing and monitoring.

Primary objective: Evaluation of the correct sampling time of vancomycin trough levels by implementing a vancomycin order set guiding nurses and physicians on the appropriate sampling time.

Secondary objectives: Evaluation of the impact of vancomycin order set integrated into CPOE system on vancomycin initial doses, appropriate use based on indication and time to reach target therapeutic level.

Method: This study was a prospective quasi-experimental study. Patients who are >18 years, admitted to surgical wards, receiving new vancomycin orders, creatinine clearance ≥ 15 ml/min and weighing 40-120 kg were eligible. We excluded patients receiving vancomycin loading doses, prophylaxis, treatment course <5 days and random or peak vancomycin levels. Educational sessions were provided to the surgical residents on the use the order set by pharmacy resident, subsequently vancomycin orders were then reassessed after the order set implementation.

Results: Total of 272 vancomycin trough levels were collected (136 levels in the pre-phase and 136 levels in the post-implementation phase) from 33 patients who met the study eligibility criteria. A 10% reduction in inappropriate vancomycin trough levels in the post-implementation phase was observed compared to the pre-phase ($p=0.478$) Appropriate vancomycin initial doses for the study patients were 2 out of 11 doses (18%) in the pre-phase and 22 doses in the post-phase (50%) ($p=0.078$).

Conclusion: The implementation of an institutional vancomycin order set did not result in a significant change in appropriate vancomycin initial dosing and trough level sampling time in the adult surgical wards in our hospital.

Keywords: Educational sessions; Vancomycin order set; Pharmacist; Computerized-Order-Prescriber-Entry (CPOE) system

Introduction

Vancomycin is the most widely used antibiotic for the treatment of serious gram-positive infections involving methicillin-resistant *Staphylococcus aureus* (MRSA) [1]. According to Infectious Disease Society of America (IDSA), the initial vancomycin dose is 15–20 mg/kg/dose based on the actual body weight, this dose is given intravenously every 8–12 h for adult patients with normal renal function [1]. Doses are adjusted according to the patient's estimated creatinine clearance and the maximum single dose of vancomycin is 2 g [1,2]. Inappropriate vancomycin dosing may lead to delays in therapeutic serum trough concentrations, the emergence of vancomycin resistant pathogens, and may ultimately culminate in treatment failure [3].

Trough serum concentrations have been recommended by IDSA for monitoring vancomycin nephrotoxicity [1]. Vancomycin serum or plasma trough concentrations are recommended as a surrogate marker of pharmacodynamics target attainment to predict vancomycin efficacy [4].

In patients with normal renal function, trough levels should be obtained once the patient has reached steady state prior to the fourth dose [1]. Minimum trough concentrations of vancomycin should be maintained above 10 mcg/ml in order to avoid the development of resistance. For most infections, trough serum concentrations of 10–15 mcg/ml are sufficient. However, for complicated infections including:

bacteraemia (caused by *Staphylococcus aureus*), endocarditis, osteomyelitis, meningitis, and hospital-acquired pneumonia, higher serum concentrations of 15–20 mcg/ml are required [1].

Improper sampling time can cause suboptimal dosing, lead to increased vancomycin resistance, prolonged hospital stay, and treatment failure. Morrison et al. retrospectively analyzed 2597 vancomycin trough levels during 13 months at a large medical centre. The results showed that 41.3% ($n=1075$) of vancomycin trough levels were drawn too early (defined as levels taken <10 h after administration of the previous dose). Consequently, in level which drawn too early, clinicians were more likely to decrease, discontinue, or hold a patient's vancomycin dose by (25.6%), or repeat the vancomycin level (29.2%). This may lead to an overestimation of patients' true trough levels,

***Corresponding author:** Mohammed Aseeri, King Abdul Allah Medical Research Center/ King Saud bin Abdul Aziz, University for Health Sciences, National Guard Health Affairs, Jeddah, King Abdul Aziz Medical City, Saudi Arabia, Tel: 966 2 624 0000; E-mail: AseeriMA@ngha.med.sa

Received: October 29, 2016; **Accepted** November 09, 2016; **Published** November 15, 2016

Citation: Damfu N, Aseeri M, Davis A, Hasan H, Ismail S (2016) The Impact of Pharmacist Led Vancomycin Order Set Implementation in a Computerized-Prescriber-Order-Entry (CPOE) System at a Tertiary Care Centre: A Quasi Experimental Study. J Pharmacovigil 4: 219. doi:10.4172/2329-6887.1000219

Copyright: © 2016 Aseeri M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

possible under dosing of vancomycin, or a high rate of repeated tests for vancomycin trough levels [5].

McCluggage et al., evaluated the implementation of a vancomycin nomogram in a Computerized-Prescriber-Order-Entry (CPOE) system on 522 vancomycin orders. A significant difference in the percentage of initial vancomycin orders that met nomogram recommendations in the post-implementation group (n=243) were observed compared to the pre-implementation group (n=279) (36% vs. 24%, p=0.0028). Therefore, integrating the vancomycin nomogram into the CPOE system increased the likelihood of patients receiving an initial vancomycin regimen that coincided with the nomogram's recommendations [6].

Traugott et al., evaluated the effects of Therapeutic Drug Monitoring (TDM) criteria in a CPOE system on the appropriateness of orders for vancomycin levels on 545 vancomycin serum levels in 200 patients (310 serum levels in 100 patients in the pre-implementation group and 235 serum levels in 100 patients in the post-implementation group) [7]. The percentage of appropriate orders for vancomycin levels significantly increased after criteria implementation (58% vs. 68% in the pre-implementation and post-implementation groups, respectively. p=0.02). The majority of inappropriate levels were due to improper timing of sample collections (55%) [7].

A vancomycin protocol was approved by the hospital antimicrobial committee; however, the dosing protocol has not been fully implemented or integrated into a CPOE order which might result in improper dosing, sampling time, and treatment failure. Therefore, we aim to evaluate the impact of implementing a vancomycin order set integrated into CPOE system in adult surgical patients to optimize vancomycin dosing and monitoring for better therapeutic outcomes.

Objectives

Primary objective

Evaluate the correct sampling time of vancomycin trough levels (identified from the hospital information system and patient charts).

Secondary objectives

1. Evaluate the impact of the vancomycin order set in a CPOE system on appropriateness of initial vancomycin dosing.
2. Evaluate the appropriateness of vancomycin use based on indication by reviewing cultures and patient charts.
3. Evaluate the time to reach vancomycin target trough level (by looking at the administration time of vancomycin from medication administration records, and sampling time of vancomycin levels and level results from the hospital information system).

Methods

Study design and setting

It is a quasi-experimental study conducted from July 2014 to May 2015 at adult surgical wards (90 beds) at a Tertiary Care Teaching Center, Jeddah Saudi Arabia.

Inclusion criteria

1. All adult patients, who are 18 years of age or older, admitted to surgical wards and receiving new vancomycin orders.
2. Patients with estimated Creatinine Clearance (CrCl) >15 ml/min.

3. Patients with total body weight of 40-120 kg.

Exclusion criteria

1. Patients with acute kidney injury or unstable kidney function.
2. Patients who received surgical prophylactic vancomycin doses.
3. Any patient that received a vancomycin loading dose (25-30 mg/kg/dose).
4. Patients on vancomycin <5 days.
5. Patients who had a random or peak level drawn.

This study was conducted in three phases (Figure 1).

First phase (pre-implementation phase over 5 months)

A daily report was generated by the Health Information System (HIS) to identify patients receiving vancomycin in adult surgical wards. Patients whose vancomycin orders met the eligibility criteria were included and data was collected using the vancomycin data collection sheet by pharmacy practice resident.

Second phase (electronic order set implementation and educational sessions for surgical residents over 1 month)

The hospital vancomycin dosing protocol was redesigned into an order set following Institute for Safe Medication Practices (ISMP) Standards for order sets [8], and subsequently integrated into a CPOE system through the HIS as vancomycin order set (Figure 2). Pharmacy practice resident conducted educational sessions for surgical residents on how to use the vancomycin order set.

Third phase (post-implementation phase over 5 months)

Similarly to first phase, a report of patients who received vancomycin was generated on daily basis and vancomycin orders met eligibility criteria were included. Subsequently, the pharmacy practice resident used the data collection sheet for documentation of study data and observed all vancomycin orders submitted via the order set in the HIS. Appropriate sampling time means that vancomycin trough level was taken within 1 h before the 4th dose. Appropriate vancomycin use based on indication refers to vancomycin orders which started empirically (duration less than 3 days) or given to treat gram positive cultures which are susceptible to vancomycin.

Statistical Analysis

A sample of 256 trough levels was estimated (128 before the implementation of the vancomycin order set and 128 after the implementation) to provide a 90% power to detect 20% difference for the correct sampling time before and after the implementation of the vancomycin order set with a type I error of alpha 0.05 [7].

1. Descriptive statistics are used to report baseline characteristics as Mean \pm SD for continuous variables and proportions for binary and categorical data.
2. Chi-square test is used to test for the proportion of appropriate vancomycin orders based on indication, sampling time and initial dosing.
3. Survival analysis is used to assess the median time to reach vancomycin target trough level and log rank test to compare median time to reach vancomycin level between pre-implementation phase and post implementation phase.
4. Data analysis was conducted using STATA version 14.




Figure 1: Study phases.

Excluding criteria	time within 1hr before the 4th dose:	***Dosing***	Patient weight 86-100	***Monitoring***
1. Patient less than 18 years old	-For most infections (e.g. skin and soft tissue infection): 10-15 mcg/ml.	CrCl > 60 ml/min	***Dosing***	<input type="checkbox"/> Vancomycin Level, Trough.
2. Cachectic patients with weight < 40 kg	-For complicated infections (e.g. bacteria (due to S.aureus), infection endocarditis, meningitis, hospital-acquired pneumonia, and osteomyelitis): 15-20 mcg/ml.	<input type="checkbox"/> Vancomycin 1000mg, IV, Q12h	CrCl > 60 ml/min	* DOSE TITRATION*
3. Obese patients with weight > 120 kg		CrCl = 30-60 ml/min	<input type="checkbox"/> Vancomycin 1500mg, IV, Q12h	<input checked="" type="checkbox"/> If targeted serum conc. achieved
4. Renal patients with CrCl < 15ml/min or patient on Dialysis		CrCl = 15-29 ml/min	CrCl = 30-60 ml/min	>Stable Kidney function: Draw Trough Co...
5. Patients with unstable kidney function		<input type="checkbox"/> Vancomycin 1000mg, IV, Q24h	CrCl = 15-29 ml/min	>UnStable Kidney function: Draw Trough ...
• If these situations, please contact the clinical pharmacists	Patient weight is 40 - 55 kg	***Monitoring***	<input type="checkbox"/> Vancomycin 1000mg, IV, Q48h	<input checked="" type="checkbox"/> If serum conc. are Higher than target conc.
	Dosing	<input type="checkbox"/> Vancomycin Level, Trough.	CrCl = 15-29 ml/min	>Requires prolong dosing interval and/or ...
Dosing and Monitoring	CrCl > 60 ml/min	Patient weight is 71-85 kg	<input type="checkbox"/> Vancomycin 1500mg, IV, Q48h	<input checked="" type="checkbox"/> If serum conc. are lower than target conc.
*****Dosing*****	<input type="checkbox"/> Vancomycin 750 mg IV q 12hrs	***Dosing***	***Monitoring***	>Requires either Increase in the dose or ...
•Initial Vancomycin dose (based on actual body weight) is 15-20 mg/kg per dose.	CrCl = 30-60 ml/min	CrCl > 60 ml/min	<input type="checkbox"/> Vancomycin Level, Trough.	
•Interval recommendation (based on CrCl obtained from face sheet option from the QuadraMed system.	<input type="checkbox"/> Vancomycin 750mg, IV, Q24h	CrCl = 30-60 ml/min	<input type="checkbox"/> Vancomycin 1500mg, IV, Q12h	
*****Monitoring*****	CrCl = 15-29 ml/min	CrCl = 30-60 ml/min	CrCl = 30-60 ml/min	
•Serum trough concentration sampling	<input type="checkbox"/> Vancomycin 750mg, IV, Q48h	CrCl = 15-29 ml/min	<input type="checkbox"/> Vancomycin 1750mg, IV, Q24h	
	Monitoring	<input type="checkbox"/> Vancomycin 1250mg, IV, Q12h	CrCl = 15-29 ml/min	
	<input type="checkbox"/> Vancomycin Level, Trough.	CrCl = 15-29 ml/min	<input type="checkbox"/> Vancomycin 1750mg, IV, Q48h	
	Patient weight 56-70 kg	***Monitoring***	<input type="checkbox"/> Vancomycin 1250mg, IV, Q48h	
	<input type="checkbox"/> Vancomycin Level, Trough.	<input type="checkbox"/> Vancomycin Level, Trough.	<input type="checkbox"/> Vancomycin 1750mg, IV, Q48h	

Figure 2: Vancomycin orders integrated into computerized prescriber order entry.

Ethics/IRB Approval

This study received an approval from the hospital ethical committee on July, 2014.

A research amendment was approved by IRB to change the effect sample size from 13% difference to 20% with a power of 90%, and total sample size from 462 levels to 256 levels (128 before the implementation and 128 after). This was done due to low number of vancomycin levels on surgical wards and the goal was to improve clinical significance by increasing the effect size to 20%.

Results

Data were collected prospectively over 11 months, starting from July 2014 to May 2015. Thirty three patients (11 patients in the pre-implementation phase and 22 patients in the post-implementation phase) were enrolled for a total of 272 vancomycin orders (Figure 3).

In the implementation phase, two educational sessions on use of vancomycin order set integrated into CPOE were conducted to 28 surgical residents out of 58 invited residents (48%) over one month period. Baseline characteristics of the study patients are shown in Table 1.

Characteristic	Pre-phase (n=11)	Post-phase (n=22)
Age (years) (mean ± SD)	50 ± 20	47 ± 20
Gender		
Males	9	19
Females	2	3
BMI (mean ± SD)	31 ± 10	27 ± 8
Creatinine Clearance (mL/min) (mean ± SD)	108 ± 47	122 ± 49
Indication		
1) Empirical	4	5
2) Bacteremia	2	1
3) Osteomyelitis	2	5
4) Skin and soft tissue infections	1	3
5) Meningitis	0	2
6) Post brain abscess extraction surgery	1	2
7) Infective Endocarditis	1	0
8) UTI	0	1
9) Peritonitis	0	2
10) Aspiration Pneumonia	0	1
Organism		
1) No Identified Organism	4	9
2) MRSA	2	3
3) Staphylococcus epidermis	5	4
4) <i>Enterococcus faecalis</i>	0	4
5) Bacillus	0	2
Duration of vancomycin treatment (days) (mean ± SD)	28 ± 22	23 ± 18

Table 1: Baseline characteristics.

Primary objective results

Correct sampling time of vancomycin trough levels: A total of 272 vancomycin trough levels were collected during the study period (136 levels the pre-implementation phase and 136 levels in the post-implementation phase) (Figure 4). Pre-implementation phase had 52.9% (72/136) appropriate trough levels, while the post-implementation phase had 62.5% (85/136) appropriate orders with a p-value of (0.478).

Secondary objectives results

Appropriate vancomycin initial dosing: We have 2 out of 11 patients (18%) in the pre-phase had appropriate initial vancomycin orders compared to 11 out of 22 patients (50%) with a p-value of 0.078.

Appropriate vancomycin use based on indication: Among the 33 patients who received vancomycin (11 in the pre-phase and 22 in the post-phase), 1 patient in the pre-phase and 1 patient in the post phase were inappropriately indicated for vancomycin (P=0.6).

Time to reach target therapeutic vancomycin level: Five patients (1 out of 11 in the pre-phase and 4 out of 22 in the post-phase) never reached their vancomycin therapeutic levels. The median time to reach vancomycin therapeutic level in the study was 65 h with an IQR of 6).

There is no statistically significance difference between pre and post in time to reach steady state concentration using log rank test (P value=0.9725) (Figure 5 for demonstration of Kaplan-Meier curves for survival estimates between pre and post-phase).

Discussion

The primary objective of the study showed a 9.6% difference in reduction of inappropriate vancomycin sampling time ordered between the post- implementation phase (62.5%) and the pre-phase

(52.9%), however, the results were not statistically significant (p=0.478). Our findings are comparable to Traugott et al. study [7], which had a larger sample size with 545 vancomycin serum levels in 200 patients (310 serum levels in 100 patients in the pre-implementation group and 235 serum levels in 100 patients in the post-implementation group). This study evaluated the effects of Therapeutic Drug Monitoring (TDM) criteria in a CPOE system on the appropriateness of orders for vancomycin serum levels, and the percentage of appropriate orders for vancomycin levels significantly increased after criteria implementation (58% vs. 68% in the pre-implementation and post-implementation groups, respectively; p=0.02) [7]. This study included all adult patients who met the study eligibility criteria in all hospital wards [7], compared to our study which only focused on adult surgical wards.

Our secondary objective was to evaluate the appropriateness of vancomycin initial dosing, showed an improvement in the post-phase from 2/11 patients (18%) to 11/22 patients (50%) but the findings were not significant with p=0.078. This could be attributed to small number of participants in our study which included only 33 patients and showed an improvement in the initial vancomycin dosing by 32%. These findings are consistent with McCluggage et al. [6], who evaluated the implementation of a vancomycin nomogram a CPOE system




Figure 3: Trial profile (for patients).




Figure 4: Trial profile (for vancomycin level orders).




Figure 5: Kaplan Meier estimate to demonstrate the differences between pre-phase and post-phase.

on 522 vancomycin orders, and showed a statistically significant difference in the percentage of initial vancomycin orders that met the study nomogram recommendations in the post-implementation group (n=243) compared to the pre-implementation group (n=279) (36% vs. 24%, p=0.0028) [6].

Our study evaluated the appropriate vancomycin use based on indication, and no statistically significant difference between the pre and post implementation groups were noted (P=0.6). Another study objective was to evaluate the time to reach target therapeutic vancomycin level, which did not show any statistically significant difference between in time to reach steady state concentration, the pre-phase and post-phase, p-value (0.9725).

Our study had several limitations; first, we did not assess the adherence of surgical resident to the use of vancomycin order set. Second, there were insufficient number of residents attended the educational sessions, total of 58 residents were invited and only 28 residents (48%) attended the sessions.

Additionally, we conducted educational sessions within one month, which might had an impact on negative findings, as it may be considered a short time for the residents to adapt on using the order set. On contrary to Hall et al. [9], who retrospectively studied vancomycin order set implementation through CPOE in emergency department, a period of 2-years between the pre-implementation and post-implementation groups was chosen to allow for implementation, physician education, and to acquire familiarity with the order set. The study resulted in a 21.9% increase in appropriate dosing in post-CPOE group compared to the pre-CPOE group (p<0.05) [9].

Third, we had a small number of vancomycin patients enrolled during the study period as we accounted for vancomycin orders in our sample size estimation and not number of patients. A larger sample might detect smaller differences between pre and post intervention. Despite we were aiming for a clinically meaningful difference of 20% between the study groups, this might have contributed to the negative findings of our study with detection of only 9.6% Fourth, the quasi-experimental study design using one point before and after for different population, has many limitations that might impact on internal validity such as lack of blinding, maturation, instrumentation, regression to the mean, selection bias and also difficulty to know if the intervention was only responsible for any observed changes [10].

Fifth, the intervention which was only focused on education about using the order set, which considered one of the least effective methods to reduce medication errors. Methods such as forced function or system alerts or reminders will be more effective [11].

And finally, the study has limited generalizability to surgical residents who may need more in depth knowledge in vancomycin dosing and monitoring.

Our study strengths included the quasi-experimental study design, actual implementation of an order set in a CPOE system, and although the findings are not statistically significant, they may provide clinically insights on role of pharmacist in initial assessment of vancomycin orders, future studies should assess the utility of forced function and clinical decision support system to enhance adherence and utilization of order sets in clinical practice to improve desired therapeutic outcomes.

Conclusion

The implementation of an institutional vancomycin order set and incorporated it in the hospital CPOE system did not result in a statistically significant change to appropriate vancomycin initial dosing and trough level sampling time in our hospital adult surgical wards however, a quasi-experimental, study with larger number of patients, use of forced functions to maximize adherence are highly recommended.

References

1. Rybak M, Lomaestro B, Rotschafer JC, Moellering R, Craig W, et al. (2009) Therapeutic monitoring of vancomycin in adult patients: a consensus review of the American Society of Health-System Pharmacists, the Infectious Diseases Society of America, and the Society of Infectious Diseases Pharmacists. *Am J Health Syst Pharm* 66: 82-98.
2. Liu C, Bayer A, Cosgrove SE, Daum RS, Fridkin SK, et al. (2011) Clinical Practice Guidelines by the Infectious Diseases Society of America for the Treatment of Methicillin-Resistant *Staphylococcus Aureus* Infections in Adults and Children. *Clin Infect Dis* 52: 1-38.
3. Rodman DP, McKnight JT, Rogers T, Robbins M (1994) The appropriateness of initial vancomycin dosing. *J Fam Pract* 38: 473-477.
4. Rybak MJ (2006) The pharmacokinetic and pharmacodynamics properties of vancomycin. *Clin Infect Dis* 42: S35-S39.
5. Morrison AP, Melanson SE, Carty MG, Bates DW, Szumita PM, et al. (2012) What proportions of vancomycin trough levels are drawn too early? Frequency and impact on clinical actions. *Am J Clin Pathol* 137: 472-478.
6. McCluggage L, Lee K, Potter T, Dugger R, Pakyz A (2010) Implementation and evaluation of vancomycin nomogram guidelines in a computerized prescriber-order-entry (CPOE) system. *Am J Health-Syst Pharm* 67: 70-75.
7. Traugott K, Maxwell P, Green K, Frei C, Lewis JS (2011) Effects of therapeutic drug monitoring criteria in a computerized prescriber-order-entry (CPOE) system on the appropriateness of orders for vancomycin level orders. *Am J Health-Syst Pharm* 68: 347-352.
8. <http://www.ismp.org/tools/guidelines/standardordersets.pdf>
9. Hall B, Montero, J, Cobian J, Regan T (2015) The effects of an electronic order set on vancomycin dosing in the ED. *Am J Emerg Med* 33: 92-94.
10. Krass I (2016) Quasi experimental designs in pharmacist intervention research. *Int J Clin Pharm* 38: 647-654.
11. https://www.ismp.org/newsletters/ambulatory/archives/200602_4.asp