



## Urinary Melatonin Levels and Endothelial Dysfunction in Non Obese Hypertension Subjects

Nakkeeran Mathivanan\*, Santha K, Inmozhi Sivakamasundari R

Department of Biochemistry, Rajah Muthiah Medical College, Annamalai University, Tamil Nadu, India

### ABSTRACT

**Introduction:** Melatonin, the principal product of the pineal gland, is metabolized in the liver mainly to 6-sulfatoxymelatonin. Melatonin is reason for diseases, like cancer, hypertension, depression and diabetes; its secretion is influenced by environmental light. As this substance is mostly excreted by urine, urinary 6-sulfatoxymelatonin is considered a good index of melatonin production. In vascular diseases, endothelial dysfunction is a systemic pathological state of the endothelium and can be broadly defined as an imbalance between vasodilatation and vasoconstriction substances produced by the endothelium. Endothelial dysfunction can result from and/or contribute to several disease processes, as occurs in hypertension, hypercholesterolemia, diabetes and it can also result from environmental factors, such as from smoking tobacco products and exposure to air pollution.

**Objective:** To investigate the association of urinary 6 sulfatoxymelatonin levels and endothelial dysfunction in hypertension subjects.

**Methods:** 63 hypertensive subjects without cause of any secondary hypertension. 35 subjects were control in this study, total of 98 subjects were studied. Subjects with secondary hypertension, past history of stroke, coronary artery disease, myocardial infarction were excluded from this study.

**Results:** The systolic and diastolic blood pressure considered to be elevated in hypertensive patients and statistically significant when compared to the control group. Our study shows a significant difference between urine aMT6s and NO in hypertension subjects compare to control.

**Conclusion:** The results suggest that melatonin and nitric oxide levels are low in subjects with hypertension, thus endothelial dysfunction precedes and predicts the development of hypertension.

**Keywords:** Urinary 6 sulfatoxymelatonin; Hypertension; Endothelial dysfunction

### INTRODUCTION

Another name of high blood pressure is hypertension. High blood pressure is the force exerted by the blood against the walls of the blood vessels. If, the blood pressure read shows >160 it is said to be hypertensive crisis, on this condition the person should wait for 2 or 3 minutes and then repeat the test. High blood pressure is one of the most common Non communicable disorders [1]. It is present in 60%-70% of the population over 60 years of age and may result in cardiovascular complication such as stroke, coronary heart disease, and heart failure. The most common type of hypertension can affects 95% of primary

hypertensive patients. It can be diagnosed when the systolic BP ranging between 120 mmHg and 138 mmHg and/or diastolic BP between 90 mmHg and 98 mmHg at three random checks and need medical monitoring and lifestyle changes. Melatonin is involved in a variety of diseases, including cancer, insomnia, depression, dementia, hypertension, and diabetes; its secretion is influenced by environmental light [2]. A pineal gland hormone secreted predominantly at night, is involved in not only regulation of circadian rhythm but also prevention of a variety of diseases, including cancer, insomnia, depression, dementia, hypertension and diabetes. Melatonin, the principal product of the pineal gland, is metabolized in the liver mainly to 6-

**Correspondence to:** Nakkeeran Mathivanan, Department of Biochemistry, Rajah Muthiah Medical College, Annamalai University, Tamil Nadu, India, Tel: 09443049480; E mail: dr.inmozhi@gmail.com

**Received:** 30-Dec-2020, Manuscript No. CMBO-23-7778; **Editor assigned:** 04-Jan-2021, PreQC No. CMBO-23-7778 (PQ); **Reviewed:** 18-Jan-2021, QC No. CMBO-23-7778; **Revised:** 31-Jan-2023, Manuscript No. CMBO-23-7778 (R); **Published:** 28-Feb-2023, DOI: 10.35841/2471-2663.23.9.149

**Citation:** Mathivanan N, Santha K, Sivakamasundari IR (2023) Urinary Melatonin Levels and Endothelial Dysfunction in Non Obese Hypertension Subjects. Clin Med Bio Chem. 9:149.

**Copyright:** © 2023 Mathivanan N, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

sulfatoxymelatonin. As this substance is mostly excreted by urine, urinary 6-sulfatoxymelatonin is considered a good index of melatonin production [3]. Melatonin is primarily involved in the regulation of biological rhythms. In addition, melatonin exerts a modulatory action over the vascular tone, with several studies indicating that melatonin affects vascular smooth muscle contraction. This effect could be homeostatically relevant to counteract the rise in arterial peripheral resistance brought about by neuroendocrine activation in CHF. Increased endogenous melatonin secretion might be associated with lower risk of these diseases; however, the determinants of the amount of endogenous melatonin secretion remain unclear [4].

Physiologically, light exposure is the most important modifier of melatonin secretion in humans. Recently, photosensitive retinal ganglion cells and the photo pigment melanopsin were identified as primary receptors of environmental light, and the mechanisms underlying the association between light exposure and melatonin secretion are being solved [5]. Light exposure at night is a well established and powerful suppressor of nocturnal melatonin secretion through the activation of the Suprachiasmatic Nucleus (SCN) of the hypothalamus, which contains the body's master biological clock. Serum melatonin has a very short half life and is rapidly metabolized, mainly in the liver.

Measurements of the principal metabolite of melatonin, 6-sulfatoxymelatonin (aMT6-s), which is excreted in urine, reflect pineal function; various studies have shown a high correlation between [6]. Measurements of aMT6 in urine and plasma. Endothelial dysfunction is a systematic pathological state of the endothelium and can be defined as an imbalance between vasodilating and vasoconstriction produced by the endothelium. Endothelial dysfunction can result from contribute to several disease processes, as occurs in hypertension like, hypercholesterolemia, diabetes, septic shock and it can also results from environmental factors, such as smoking tobacco products and exposure to air pollution [7]. Studies suggest that endothelial dysfunction is more prevalent in shift workers, a group known to have a higher risk for cardiovascular diseases. Nitric oxide is a molecular and chemical compound with formula of NO. In mammals, nitric oxide is an important cellular signaling molecule involved in many physiological and pathological processes [8]. It is a powerful vasodilator with a short half life in the blood. Long term known pharmaceuticals such as nitroglycerine and amyl nitrite were found to be precursors to nitric oxide more than a century after their first use in medicine [9].

Nitric is essential for hepatic lipid metabolism during starvation. It is a key vertebrate biological messenger, playing a role in a variety of biological processes. Nitric oxide, known as an Endothelium Derived Relaxing Factor (EDRF), is biosynthesized endogenously from L-arginine, and NADPH by Nitric Oxide synthase (NOS) enzymes. Reduced state of inorganic nitrate may also serve to make nitric oxide [10]. The endothelium of blood vessels can use the nitric oxide to signal the surrounding smooth muscle to relax, thus resulting in vasodilation and increasing blood flow. Nitric oxide is highly reactive, yet diffuses freely across membranes. Independent of

nitric oxide synthase, an alternative pathway, coined the nitrate nitrite nitric oxide pathway, elevates nitric oxide through the sequential reduction of dietary nitrate derived from plant based foods [11]. Nitric Oxide (NO) contributes to blood vessel homeostasis by inhibiting vascular smooth muscle contraction and growth, platelet aggregation, and leukocyte adhesion to the endothelium. Human subject with atherosclerosis, diabetes, hypertension often show impaired NO pathways [12].

## MATERIALS AND METHODS

The study design included two groups of human subjects as participants. A total of 98 subjects were studied. Among these 63 were diagnosed hypertensive patients without cause of any secondary hypertension [13]. 35 subjects were control in this study. Hypertension was diagnosed when the systolic and diastolic pressure read 140 mmHg and 90 mmHg, respectively, at two to three random checks. Rajah Muthiah medical college and hospital, Annamalai university, Tamil Nadu, India. All the participants underwent a detailed clinical examination [14]. Different parameters such as age, BP, Hb, and ECG were recorded. The study was approved by both research and ethics committee of our institute. A fully informed consent was obtained from these subjects before participation in the study [15].

### Subject selection criteria

All volunteers enrolled in this study were aged between 30 and 65 years. Subjects of both sexes were included. Women, being on a regular menstrual cycle were included in the study. Subjects with hypertension without any secondary causes were included in this study [16]. Subjects with secondary hypertension, past history of stroke, coronary artery disease, myocardial infarction, and peripheral vascular disease with evidence of tissue injury or loss, renal diseases, diabetes mellitus, after infections, smokers, alcoholic and who with vitamin supplement, known to cause changes in antioxidant and lipid status, were excluded from the study [17]. Serum lipid profile, FRAP Assay and serum TBARS, high sensitivity C reactive protein were estimated in these patients by standard procedures and the values were compared with healthy control subjects. Separated serum was used to analyze lipid profile, uric acid, Malondialdehyde (MDA), hsCRP were estimated in these patients by standard procedures and the values were compared with healthy control subjects [18]. Total cholesterol, High Density Lipoprotein Cholesterol (HDL-C), Triglyceride (TG), was estimated using Erba assay kits. Low Density Lipoprotein Cholesterol (LDL-C) was calculated by Friedewalds formula. MDA was measured using Thio Barbituric Acid (TBA). The mixture was heated at 100°C for 15 min, allowed to cool and centrifuged at 5000 rpm. Total anti oxidant status was assayed by measuring FRAP (Ferric Reducing Activity of Plasma) urine 6-sulfatoxymelatonin performed by ELISA kit method and nitric oxide was done by Griess reagent [19].

### Statistical analysis

Fischer exact test has been used to find the significance of various parameters among cases and controls. Data presented

are mean  $\pm$  SD. Analysis of data was done by student t test. A 'p' value  $<0.05$  was considered significant.

## RESULTS AND DISCUSSION

The compared values of the study group both hypertension and control subjects are shown in the Table 1. Our data clearly show that significant increase in Total Cholesterol, Triglyceride, LDL-C, where as HDL-C, antioxidants, were decreased among hypertensive compared to controls, marking ongoing oxidative stress in hypertensive patients. Table 2 shows a correlation between urinary melatonin and lipid profile. There will be a

positive correlation between LDL cholesterol and Triacylglycerol, There will be a negative correlation between urinary melatonin and nitric oxide. Table 3 shows a correlation between Nitric oxide and lipid profile there will be a positive correlation between LDL cholesterol and Triacylglycerol. The association of blood pressure depend upon the combination of risk factors, such as age, gender, weight, physical activity, smoking, family history, serum cholesterol levels and diabetes mellitus.

**Table 1:** Comparison of study variables in test and control groups.

Variables	Test group	Control group	P-value
FBS	91.30 $\pm$ 9.22	86.66 $\pm$ 5.83	0.002
BMI	25.2 $\pm$ 0.3	24.3 $\pm$ 0.4	0.17
SBP (mmHg)	158.15 $\pm$ 10.46	121.39 $\pm$ 8.03	0.001
DBP (mmHg)	83.37 $\pm$ 10.79	77.80 $\pm$ 3.21	0.001
Triglycerides	126.97 $\pm$ 29.29	111.78 $\pm$ 23.91	0.002
Total cholesterol	189.71 $\pm$ 25.99	175.95 $\pm$ 23.42	0.002
HDL	43.73 $\pm$ 2.79	54.34 $\pm$ 4.72	0.004
LDL	89.28 $\pm$ 19.90	81.83 $\pm$ 13.51	0.022
FRAP assay	180.20 $\pm$ 58.72	223.71 $\pm$ 64	0.001
Nitric oxide	15.68 $\pm$ 4.48	22.74 $\pm$ 9.34	0.0001
Urinary melatonin	224.73 $\pm$ 75.16	289.8 $\pm$ 80.71	0.0002

**Table 2:** Correlation between urinary melatonin and lipid profile.

Variables	R-value	P-value
Urinary melatonin vs. LDL	0.442	0.0001
HDL	-0.287	0.0001
TAG	0.329	0.0008
TC	-0.029	0.0007
Nitric oxide	-0.19	0.0002

**Table 3:** Correlation between nitric oxide and lipid profile.

Variables	R-value	P-value
NO vs. LDL	0.062	0.0001
HDL	-0.069	0.0001
TAG	0.192	0.0008
TC	-0.468	0.0007

The present study aimed to investigate the association of urinary melatonin and endothelial dysfunction in hypertension subjects. Several studies reported that melatonin has antihypertensive effects but not clear with urinary melatonin. Our results clearly show the significant difference and correlation between the study groups. Among 98 subjects 63 were hypertensive and 35 were control subjects included in this study and the results were compared. Our data shows a negative correlation between urinary melatonin with HDL-C, Total cholesterol and nitric oxide. Correlation with Nitric oxide and cholesterol it also shows the same result that HDL-C, Total cholesterol. Whereas LDL-C and triglycerides were shows positive correlation between the control group and to the study group. Therefore High excretion levels of morning urinary melatonin levels and low levels of nitric oxide levels were independently associated with a increased risk of developing hypertension. To our knowledge, this is the first study to demonstrate an association between urinary melatonin levels with nitric oxide in hypertension. In this study, we found that low levels processing of urine samples also had only a modest effect on aMT6-s levels, with slightly lower levels of aMT6-s associated with developing hypertension. Thus, first spot morning urine melatonin measurements are a feasible biomarker in observational studies. Physiologically, light exposure is the most important modifier of melatonin secretion in humans. Recently, photosensitive retinal ganglion cells and the photo pigment melanopsin were identified as primary receptors of environmental light, and the mechanisms underlying the association between light exposure and melatonin secretion are being solved. Serum melatonin has a very short half-life and is rapidly metabolized, mainly in the liver. Measurements of the principal metabolite of melatonin, 6-sulfatoxymelatonin (aMT6-s), which is excreted in urine, reflect pineal function; various studies have shown a high correlation between measurements of aMT6-s in urine and plasma.

The endothelium is not only the inert interface between circulating blood and the vessel wall but is also a major paracrine organ that plays critical roles in controlling vascular tone, inflammation, and smooth muscle cell proliferation. NO production can be controlled by the availability of substrates and cofactors, transcription of eNOS, mRNA stability of eNOS, sub cellular localization of eNOS protein, enzymatic uncoupling, and post-translational modifications. NO is one of the few gaseous signaling molecules known and is additionally exceptional due to the fact that it is a radical gas. It is a key vertebrate biological messenger, playing a role in a variety of biological processes. The production of nitric oxide is elevated in populations living at high altitudes, which helps these people avoid hypoxia by aiding in pulmonary vasculature vasodilatation. Nitric Oxide (NO) contributes to vessel homeostasis by inhibiting vascular smooth muscle contraction and growth, platelet aggregation, and leukocyte adhesion to the endothelium. Humans with atherosclerosis, diabetes mellitus or high blood pressure often show impaired NO pathways. A high salt intake was demonstrated to attenuate NO production in

patients with essential hypertension, although bioavailability remains unregulated.

## CONCLUSION

Our results shows the significant difference between the study group when compare to the control group, low urinary melatonin levels are associated with cause of hypertension and low melatonin production may be a pathophysiological factor for development of hypertension. Endothelial nitric oxide abnormality in hypertensive subjects shows an endothelial nitric oxide deficiency will be results in higher blood pressure. But, in contrast to these findings, we also know that endothelial NO deficiency can be precipitated by a variety of non hypertension related results that increase vascular oxidative stress, such as hypercholesterolemia. More studies are required to explore the potential mechanisms, and additional clinical studies are needed to confirm this association.

## REFERENCES

- Schernhammer ES, Berrino F, Krogh V, Secreto G, Micheli A. Urinary 6-sulphatoxymelatonin levels and risk of breast cancer in premenopausal women: The ORDET cohort melatonin and breast cancer risk. *Cancer Epidemiol. Biomarkers Prev.* 2010;19(3): 729-737.
- Provencio I, Jiang G, de Grip WJ, Hayes WP, Rollag MD. Melanopsin: An opsin in melanophores, brain, and eye. *Proc Natl Acad Sci USA.* 1998;95(1):340-345.
- Zeitler JM, Dijk DJ, Kronauer RE, Brown EN, Czeisler CA. Sensitivity of the human circadian pacemaker to nocturnal light: Melatonin phase resetting and suppression. *J Physiol.* 2000;526(3): 695-702.
- Deanfield J, Donald A, Ferri C, Giannattasio C, Halcox J, Halligan S, et al. Endothelial function and dysfunction. Part I: Methodological issues for assessment in the different vascular beds: A statement by the working group on endothelin and endothelial factors of the European society of hypertension. *Am J Hypertens.* 2005;23(1):7-17.
- Munzel T, Sinning C, Post F, Warnholtz A, Schulz E. Pathophysiology, diagnosis and prognostic implications of endothelial dysfunction. *Ann Med.* 2008;40(3):180-196.
- Hou YC, Janczuk A, Wang PG. Current trends in the development of nitric oxide donors. *Curr Pharm Des.* 1999;5(6): 417-442.
- Gu Q, Yang X, Lin L, Li S, Li Q, Zhong S, et al. Genetic ablation of solute carrier family 7a3a leads to hepatic steatosis in zebrafish during fasting. *Hepatology.* 2014;60(6):1929-1941.
- Dessy C, Feron O. Pathophysiological roles of nitric oxide: In the heart and the coronary vasculature. *Antiinflamm Antiallergy Agents Med Chem.* 2004;3(3):207-216.
- Hernandez C, Abreu J, Abreu P, Castro A, Jimenez A. Nocturnal melatonin plasma levels in patients with OSAS: The effect of CPAP. *Eur Clin Respir J.* 2007;30(3):496-500.
- Berson DM, Dunn FA, Takao M. Phototransduction by retinal ganglion cells that set the circadian clock. *Science.* 2002;295(5557): 1070-1073.

11. Provencio I, Jiang G, de Grip WJ, Hayes WP, Rollag MD. Melanopsin: An opsin in melanophores, brain, and eye. *Proc Natl Acad Sci USA*. 1998;95(1):340-345.
12. Nowak R, McMillen IC, Redman J, Short RV. The correlation between serum and salivary melatonin concentrations and urinary 6-hydroxymelatonin sulphate excretion rates: Two non invasive techniques for monitoring human circadian rhythmicity. *Clin Endocrinol*. 1987;27(4):445-452.
13. Lang U, Kornemark M, Aubert ML, Sizonenko PC. Radioimmunological determination of urinary melatonin in humans: Correlation with plasma levels and typical 24-hour rhythmicity. *J Clin Endocr*. 1981;53(3):645-650.
14. Arendt J, Bojkowski C, Franey C, Wright J, Marks V. Immunoassay of 6-hydroxymelatonin sulfate in human plasma and urine: Abolition of the urinary 24-hour rhythm with atenolol. *J Clin Endocr*. 1985;60(6):1166-1173.
15. Cook MR, Graham C, Kavet R, Stevens RG, Davis S, Kheifets L, et al. Morning urinary assessment of nocturnal melatonin secretion in older women. *J Pineal Res*. 2000;28(1):41-47.
16. Dudzinski DM, Igarashi J, Greif D, Michel T. The regulation and pharmacology of endothelial nitric oxide synthase. *Annu Rev Pharmacol Toxicol*. 2006;46:235-276.
17. Searles CD. Transcriptional and posttranscriptional regulation of endothelial nitric oxide synthase expression. *Am J Physiol Cell Physiol*. 2006;291(5):803-816.
18. Proctor PH. Endothelium-derived relaxing factor and minoxidil: Active mechanisms in hair growth. *Arch Dermatol*. 1989;125(8):1146-1147.
19. Osanai T, Fujiwara N, Saitoh M, Sasaki S, Tomita H, Nakamura M, et al. Relationship between salt intake, nitric oxide and asymmetric dimethylarginine and its relevance to patients with end-stage renal disease. *Blood Purif*. 2002;20(5):466-468.