

## Validation Study in Membrane Chromatography Adsorber and Phenyl Hydrophobic Membrane Chromatography Adsorber for Virus Clearance and Removal of Many Other Components

Hideharu Shintani\*

Chuo University, School of Science, 1-13-27, Kasuga Bunkyo 112-0003 Tokyo, Japan

### Abstract

The paper Clearly shows that film chromatography adsorbers altogether offer an adaptable and financially savvy result for a percentage of the bottlenecks in immunization assembling and infection freedom and are picking up acknowledgement as an elective to customary pitch based chromatography, especially since they kill cleaning and approval expenses, give adaptability in handling prepare plan and scale-up, and could be utilized within viral freedom steps.

Since the favorable circumstances of layer chromatography adsorbers have just been acknowledged in cleaning provisions thus far, the open inquiry is the manner by which to address different bottlenecks, especially the catch go in counter acting agent produce. All the more in particular, what is the part for stuffed cot chromatography in future forms? Four surveys distributed in the course of recent years reach basically the same conclusion, that pressed couch chromatography is still the workhorse in bioseparation and will stay as the standard for in any event the following 5 years in this preservationist industry, yet it can't truly adapt to the tests furnished by expanding maturation titers. As the midway empowering innovation, it has established the framework for downstream handling as an autonomous control. In any case, in spite of the fact that it is staying put for the foreseeable future, it is presently gradually coming to be part of the issue and not the result. The essential address hails from a financial and quality stance, since more advanced in years empowering advances for example precipitation could now furnish a just as satisfactory and a great deal more monetary elective.

With the reach of elective advances now getting accessible, pressed cot chromatography may not involve this hegemonic position sometime later yet might rather be one of a determination of just as feasible results for tomorrow's downstream handling tests.

**Keywords:** Validation study; Membrane chromatography adsorber; Phenyl Hydrophobic

### Introduction

Downstream preparing might never have improved as a single division of the bioprocessing business without chromatography, whose natural straightforwardness and selectivity has made it the crux empowering engineering in all bioseparation methods [1-6].

The most usually utilized chromatography methods incorporate particle trade chromatography (lex), partiality chromatography, and hydrophobic collaboration chromatography/reverse stage chromatography (Hic/rpc). The sap sciences and the arrangement of the steps for utilization are chosen to meet the goals of the sanitization process regarding selectivity, speed, recuperation, and limit [7].

Even though Protein A chromatography has experienced incredible streamlining for throughput in the previous five years, the normal ascent in cell society titers, consolidated with the requirement for bigger masses of antibodies, demonstrates a necessity for further change of the existing Protein A gum to oblige the interest for towering throughput creation [4]. Consequently, there has been huge investment in the infrastructure of new advances that expedite the transforming of multikilogram parcels at the most minimal conceivable cost [1,7]. Around the novel innovations advanced, film chromatography adsorber has an elevated potential, especially for methodology scale monoclonal neutralizer (mab) refinement.

### Membrane Chromatography Adsorbers

Membranes feature a more open structure than resins, so there

is essentially no dissemination restriction, and therefore they offer numerous specialized preferences that have been generally portrayed in the writing [8-11]. Requisitions have been talked over for mab cleansing. As a result of the usually towering isoelectric focuses (pi) of mabs, anion-trade (Aex) saps are regularly utilized as a cleaning go as a part of a course through mode i.e., the item does not tie while the follow contaminations for example DNA, infections, endotoxins, and host cell proteins are held [5]. Granted that the measure of pollutions to tie is exceptionally flat in such operations, accepted cleaning scale-up prompts section oversizing due to the force and dispersion restrictions connected with customary dot based chromatography.

An additional huge utilitarian point of interest of films over saps is that the transport of solutes to their coupling destinations happens essentially by convection, while pore dispersion is insignificant [9]. In view of these hydrodynamic profits, layer chromatography adsorbers can work at much more excellent stream rates than sections,

\*Corresponding author: Hideharu Shintani, Chuo University, School of Science, 1-13-27, Kasuga Bunkyo 112-0003 Tokyo, Japan, Tel: +81425922336; E-mail: [shintani@mail.hinocav.ne.jp](mailto:shintani@mail.hinocav.ne.jp)

Received May 16, 2013; Accepted May 23, 2013; Published May 26, 2013

**Citation:** Shintani H (2013) Validation Study in Membrane Chromatography Adsorber and Phenyl Hydrophobic Membrane Chromatography Adsorber for Virus Clearance and Removal of Many Other Components. Pharm Anal Acta S2: 005. doi:10.4172/2153-2435.S2-005

**Copyright:** © 2013 Shintani H. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

impressively diminishing cradle utilization and shortening the for the most part process time by up to 100-fold [12].

Conversely, a solitary utilize layer chromatography gadget has a convective mode of mass transport, which permits operation at fundamentally higher straight stream rates (shorter habitation times) than sections. Accordingly, single-utilization layer chromatography units can have a much littler volume than that of an accepted segment when utilized as a part of move through mode. This fundamentally lessens support utilization (which could be tossed after single utilize), transforming time, and floor space prerequisites [13].

Film chromatography adsorbers consequently offer a versatile, adaptable, and financially savvy result for a portion of the bottlenecks in counter acting agent make and are picking up acknowledgement as an elective to customary sap based chromatography [13-16]. Due to up to 4-fold lessening qualities realized for host-unit proteins (Hcp), DNA, and numerous infections, film chromatography adsorbers have gotten acknowledged in differentiating the item from idle contaminations and in infection leeway [13,17,18]. Film chromatography adsorber has the preference of various and also single-utilization alternatives, the recent being a standout amongst the most proficient and engaging cleaning methodologies in downstream cleaning. Numerous particle trade cases have been effectually accomplished in different downstream forms (Dsps) for evacuation of adversely charged contaminants [13,19-21].

## Integrity Test for Membrane Chromatography Adsorber

The history of membrane chromatography adsorber shows colossal advancement since 1988, preparing differences of film structures, surface sciences, and setups [22]. Business layers have astounding assembling tolerances, which make them very uniform in thickness and pore-estimate conveyance, with any remaining unevenness nullified by the utilization of stacks holding great amounts of sheets [22]. For cleaning provisions, it has been demonstrated that film chromatography adsorber has a few clear preferences over gums, yet sometime to come it will be fundamental to address certain tests that utmost the more extensive selection of this innovation stage [22].

A dissemination test is suggested by and large as a normal test for location of any gathering disappointment (e.g., welding), major detours and expansive layer absconds (e.g., pinholes through film cot). A practicality testing for the estimation of aggregate particle trade limit might be performed by titration.

At elevated stream rates, the film pores furnish satisfactory tying limit for huge biomolecules for example infections and DNA, and they assume a vital part in the generally speaking viral freedom procedure for immunizer cleansing [17,23]. Practicality and honesty tests show that even with cot statures of less than 1 cm, film stacks are hearty and dependable [7].

An uprightness test was accounted for to perform by the layer diffusivity strategy [13]. The uprightness test was performed prior and then afterward utilization of a film [13]. Briskly, before establishment of the layer unit to the cleaning slide, strip cradle (1m NaCl in 20 mm Tris ph 7.2) was pumped through the case, set up vertically (upward stream), at 225 cm/h until the outlet conductivity matched the gulf's conductivity (pretty nearly 2-3 Dv). This was accompanied with 5 Dv 10 mm Tris cushion of ph 7.2 at 450 cm/h. After the results from channel and outlet matched one another, the pump was ceased and the bay and outlet were clipped off. The qualified film may as well hold a gulf force more amazing than 10 psi (68,948 Pa) for 5 min.

The uprightness and air dissemination test was performed for every producer's method utilizing the Diffusion Test Program for the right size of case. Duplicates of the printout(s) were kept as uprightness capability records for the film chromatography adsorber [13].

## Alternatives to Conventional Chromatography

With a developing pattern to film chromatography adsorber, different plan B have been advanced either to trade segment chromatography or to diminish the burden of contaminations in the feedstream with the intention that one or more chromatography steps might be killed. Some of these options have an association with catch steps while others stand for inventive filtration and chromatography designs that have an association with cleaning operations [7]. Illustrations of higher-end innovations for the reinstatement of segment chromatography incorporate the utilization of charged ultrafiltration layers and film adsorbers. Here we concentrate on the utilization of film chromatography adsorber as a potential result for dodge the restrictions of gums for move through requisitions [24].

Engage in layer chromatography adsorber is developing due to the achievement of single-utilize layer channels, yet there is still an absence of valuation for the numerous focal points layer gadgets offer in downstream handling [13,15]. Rather than the gum based move through courses of action, film chromatography adsorber includes the utilization of slim, engineered microporous or macroporous layers that are stacked some layers profound in a relatively minor cartridge. The foot shaped impression of such units is much more modest than sections with a comparable yield.

This diminishes cradle utilization however increments the stream rate since the couch tallness is much lower and there is a lessened force drop [25]. Regardless of the expanded stream rate, adsorption is effective on the grounds that the transport of solutes to their coupling destinations in a film chromatography adsorber happens basically by convection, while pore dispersion (the dominating instrument in saps) is negligible. These profits lessen transform times to less than 10% of the aforementioned connected with universal stainless-steel sections [12]. An additional essential playing point is the straight scale-up for parameters for example frontal surface territory, cot volume, stream rate, and static tying limit, while standardized dynamic tying limit remains decently steady at 10% or complete achievement [16]. Table 1 shows an illustration of cradle utilization or great scale stuffed bunk chromatography and film chromatography adsorber [26].

Film chromatography adsorbers of distinctive sorts are accessible from diverse suppliers with practical assemblies proportional to the comparing tars, for instance, films holding initiated quaternary ammonium bunches for anion trade.

The utilization of film chromatography adsorbers might be seen as the identical to shortening universal sections to close zero length, permitting hefty scale techniques to run with just a minor force drop

Items	Membrane Chromatography Adsorber (70 ml)	Large Scale Packed Bed Chromatography (40 L)
Pre-equilibration	NA	3 CV = 120 L
Equilibration	10 DV = 3.5 L	5 CV = 200 L
Wash	30 DV = 10.5 L	3 CV = 120 L
Regeneration	NA	3 CV = 120 L
Storage	NA	3 CV = 120 L
Total	14l	680 L

**Table 1:** Comparison of buffer usage for large scale packed bed chromatography and membrane. Chromatography Adsorber.

at extremely heightened stream rates [7]. For instance, cleaning with an anion-trade film could be directed with a couch stature of 4 mm at stream rates of more than 600 cm/h, furnishing a heightened frontal surface zone to mattress tallness proportion [7]. Minor volume single-utilization layer chromatography units can now handle more than 10 L/min/bar/m<sup>2</sup>.

### Pressed Bed still the Preferred Choice for Capturing

Even though there is marked verbal confrontation about this issue, one can reason that tie and-elute operations for monoclonal antibodies are approaching their physical cutoff points [4,27,28]. For the present, there is no elective to pressed bunk chromatography for such operations because of its selectivity and usability. In the long run, restrictions in benefit and accordingly handle economy might better be tended to by level innovation choices for example precipitation, crystallization, or even flocculation.

In spite of the favorable circumstances of layer chromatography adsorber, stuffed couch chromatography still remains the favored operation for catching particles of <200 kda, particularly when crest cutting and inclinations are needed for the partition of nearly identified species [5]. Be that as it may, for bigger particles (counting a large portion of the expected contaminants in immune response make), layer chromatography adsorber offers higher limit and quicker preparing. For instance, course through Aex for counter acting agent cleaning with a layer chromatography adsorber might be directed with a couch tallness of 4 mm at stream rates of more than 600 cm/h, furnishing a much higher frontal surface territory to bunk stature degree than with segments. In spite of the fact that the stream rate is much more terrific than conceivable with a section, there is sufficient maintenance time to decrease DNA, for the most part Hcps, and numerous infections by up to four log diminishment values, permitting film chromatography adsorbers to be utilized not just to divide the item from latent pollutions and yet as an incorporated viral leeway step [2,13,29] (Table 2).

### Applications of Membrane Chromatography Adsorbers

At present, the profits of layer chromatography adsorbers are for the most part clear in course through requisitions on the grounds that limit obligations in maintenance mode, especially at elevated stacking rates, make pitch chromatography a more alluring alternative for catch steps [25]. The introductory catch and refinement step utilizing Protein A fondness chromatography yields an item with more than 90% virtue, however further cleaning is obliged to lessen pollutions and infections to adequate levels and accomodate with administrative guidelines.

### Utilization of Membrane Chromatography Adsorber in the Vaccine Industry

Film chromatography adsorbers have ended up being an exceptionally convenient approach likewise in the viral antibody preparation and DNA purging for gene therapeutics [30-35].

A later report guaranteed that the elevated profit, minimal effort, and effortlessness of utilization made the layer chromatography

adsorber an alluring choice for catching flu infection from unit society supernatants. There was a 5-fold improvement in water powered harsh corrosive (Ha) movement and the normal yield was 72% [36].

An additional study reported complete and combined recuperation of recombinant baculo-infection dependent upon single-utilize film engineering. A middle recuperation yield of 70% was acquired utilizing tangential stream filtration (Tff) and film chromatography adsorber after enhancement of the discriminating working parameters. A generally speaking recuperation yield of 40% was realized, gathering all the untaintedness necessities [37].

Further examinations uncovered the utilization of anion-trade film chromatography adsorber for imposing scale downstream filtration of rotavirus infection like particles (Vlps) to a clinical review at 46% worldwide recuperation yield with almost 100% evacuation of host cell DNA and around 98% evacuation of Hcps [38].

### Infection Clearance utilizing Membrane Chromatography Adsorbers

Keeping in mind the end goal to fulfill the security prerequisites for patients and administrative agreeability from administrative offices while conveying amazingly heightened item throughput, a few single-utilize frameworks have been tried and brought about in hefty scale restorative mab preparation [39]. Distributed information exhibit that film chromatography adsorber can effectively evacuate diverse demonstrate infections at towering stream rates in an expansive extend of operational forces and infection leeway limit might be as towering as 3,000 g/m<sup>2</sup> [13,40].

As of late a study reported the utilization of the particle trade layer chromatography adsorber in a counter acting agent pilot plant run to process material for a lethality study. The scale-down model utilized the 15-layer of the particle trade layer chromatography adsorber. This itemized study showed that anion trade film chromatography adsorber has an ability to uproot Hcp and infections at an unbiased ph and at conductivity beneath 3 ms/cm. At these conditions, great viral evacuation information were accounted for in their huge scale neutralizer methodology: 5.57 log decrease esteem (Lrv) for murine leukemia infection (Mulv), 7.28 Lrv for Reo-3, 6.77 Lrv for moment infection of mice (Mvm), and 5.67 Lrv for pseudorabies infection (Pry). At technique scale, anion-trade layer chromatography adsorber apparatuses have additionally been utilized for endotoxin evacuation [41].

Freedom competencies of particle trade layer chromatography adsorber for nonenveloped infections have been demonstrated to be between 3.56 log<sub>10</sub> for Mvm and more than 6.92 log<sub>10</sub> for porcine parvovirus (Ppv) [42]. It has been showed that the stage tried layer chromatography adsorber, has a procedure limit more amazing than 3,000 g mab/m<sup>2</sup> or 10.7 kg of mab/l with a Log Reduction Values (Lrv) > 5 for four model infections [13]. Mass adjust in viral freedom studies is a different imperative parameter to exhibit productive infection evacuation by layer chromatography adsorbers. A 100% recuperation was showed for Prv, Reo infection sort 3 (Reo-3), and Mvm, when the layers were stripped with 1-M NaCl, outlining productive charge catch for the three model infections while elevated salt medication of the layer chromatography adsorber demonstrated 70% recuperation for Mulv [43]. The infection leeway proficiencie of such innovation is displayed in Table 2 and 3 [13,40].

Infection leeway with single-utilize layer chromatography adsorber is being examined and working systems are being investigated in

Viruses	Size (nm)	LRV(Run 1)	LRV(Run 2)	Recovery (%)
MVM	16-25	6.03 ± 0.21	6.03 ± 0.20	100
Reo 3	75-80	7.00 ± 0.31	6.94 ± 0.24	100
MuLV	80-110	5.35 ± 0.23	5.52 ± 0.27	>70
PRY	150-250	5.58 ± 0.28	5.58 ± 0.22	100

**Table 2:** Log Reduction Values (LRV) of Different Viruses Demonstration Mass Balance upon Adsorption by Ion Exchange Membrane Chromatography Adsorber.

different quarters [20,43]. An expansive go of different operational parameters are almost always tried with a specific end goal to accomplish acceptable and strong infection freedom in mab refinement (Table 4) [43].

Since anion trade film chromatography adsorber is a change-based mechanism, updates in operational ph and conductivity were thought to have an effect on the proficiency of infection leeway. It was therefore watched that the Lrv worth for Mvm was towering when the operational ph was towering. Since the infection particles have an isoelectric focus (pi) of 5, the analyses were performed at a ph of > 6.4. At pH 7.99, and the most elevated Lrv was gotten. Further perceptions sharp out that infection freedom power was marginally diminished when the operational conductivity was expanded (Table 5) [43].

The studies examined decidedly exhibit the capacities of the particle trade layer chromatography adsorber for evacuation of all model infections with suitable enhancement of operational parameters for example pH, conductivity, stream rate, and temperature. The point when the infections were spiked in distinctive food holding different proteins, differed Lrv qualities were gotten because of the macromolecular connections. Rather than filtration, the load based particle trade layer adsorber showed potential viral leeway with controllable characteristics. Numerous research endeavors have exhibited the prevalence of their execution in mab preparation [13,20,40].

From these examinations it is apparent that the particle trade based film chromatography adsorber has as of recently ended up being a capable elective to cleaning segments in move through mode for contaminant evacuation. All the more as of late, new layer chromatography adsorbers have been configured for utilization in other applications.

## Hydrophobic Interaction Chromatography

The present discourse will now concentrate on the exhibition of the new phenyl layer adsorber that has been created for the purging of biomolecules dependent upon hydrophobic connection chromatography (Hic) standards. This item, called Phenyl, joins the

Run	Load Condition	MuLV	PRV	Reo 3	MVM
A	0.49 kg/m <sup>2</sup> at 240 cm/h ~3 ms/cm, pH 7	36.23	35.76	37.53	34.41
B	0.049 kg/m <sup>2</sup> at 240 crn/h	35.57	35.67	37.28	36.67

**Table 3:** Process capacity and virus adsorption capability of ion exchange membrane chromatography adsorber.

Run	Flow Rate	X-MuLV	PRV	Reo 3	MVM
A	3	4.50	5.35	5.58	7.00
B	3	5.52	6.94	-	-
C	3.6	6.00	5.59	-	-

**Table 4:** Virus clearance power of ion exchange membrane chromatography adsorber.

Capacity	Operational pH	Virus Clearance
200	6.4	4.77
	7.22	5.53
	7.99	5.96
3000	4	5.09
	5	4.47
2500	6	4.29

**Table 5:** Effect of operational pH and conductivity on mvm removal capacity of anion exchange membrane. Chromatography Adsorber.

points of interest of film chromatography adsorber with a heightened hydrophobic tying limit for proteins similar to that of routine Hic tars. It stands for another device for downstream preparing, relevant for both tie and-elute utilized forward and move through operations [44].

Various confirmations back Hic as an essential procedure for vast scale refinement of recombinant proteins dependent upon distinctions in their hydrophobicity [45-49]. Hic has additionally been displayed as a productive mode for evacuating dimers and higher atomic weight totals when utilized as a cleaning go within a monoclonal counter acting agent sanitization process [50]. By the by, advancing preparative Hic strategies might be testing, especially with the stream rate and dispersion constraints connected with pressed bunk Hic, which can build the danger of protein denaturation because of long contact time on the hydrophobic surface and the elevated centralization of lyotropic salt around the protein, both of which can cause extreme item misfortunes [51]. The utilization of a convective chromatography strategy, for example film chromatography adsorber, lessens these issues by permitting much quicker preparing time. As the transport of the particles to the coupling destinations happens primarily by convection (while pore dispersion is insignificant), the mass exchange safety is lessened so catch is quick and note impressively free of stream rate. Because of the hydrodynamic profits, layer chromatography adsorber engineering includes much more diminutive gadgets than segments with a comparative throughput. This can altogether diminish cushion utilization, preparing time, and space prerequisites. Course through film chromatography specifically can safeguard up to 95% of support and 66% of procedure time as contrasted and accepted section chromatography [13]. Moreover, layer chromatography adsorbers are accessible in a solitary and prepared to utilize case organize that takes out the necessity for pressing and capability or reuse acceptance in question scale [39,52]. These emphasizes make as lessened process time and multifaceted nature while adding adaptability to the assembling office.

## Phenyl Membrane Chromatography Adsorber for Large-Scale Purification

The as of late presented Phenyl hydrophobic film chromatography adsorber is gathered into 30-layer spiral stream transform containers with a 8-mm cot tallness. It came about because of the combo of a recently composed macroporous layer structure and a covalently connected hydrophobic phenyl ligand [53,54]. The layer structure was configured for elevated stream rates and tying limits. Moreover, the hydrophilic stabilized recovered cellulose film underpin has astounding mechanical and substance strength and shows negligible nonspecific communication with proteins, even at elevated convergances of a lyotropic salt [54]. The functionalized layer shows expanding tying limit in expanding ammonium sulfate focuses, obviously in Hic [54]. Table 6 portrays the coupling limit at 10% leap forward for Phenyl hydrophobic layer adsorber tested with three diverse model proteins [54].

In a later paper, the division lands of the awhile ago referenced Hic layer were assessed and contrasted with those of an accepted Hic tar, phenyl Sepharose Ff (level sub) from Ge Health care [54]. Utilizing a protein mixture of cytochrome C, trypsinogen, and a polyclonal IgG, the Phenyl hydrophobic layer chromatography adsorber showed preferable determination over the Hic tar under the trial conditions utilized for the assessment [54]. Stuck expansion to the tough situation and-elute operation, the creators showed the provision of Phenyl in course through mode for total evacuation in a cleaning process for a



Protein	$[(\text{NH}_4)_2\text{SO}_4]$ (M)	10% DBC (mg/ml)
$\gamma$ -Globulin	0.9	17
Ovalbumin	1.5	28
Lysozyme	1.5	31

All proteins were prepared as 1 g/L solutions in 50 mM phosphate buffer at pH 7.0 with varying amounts of ammonium sulfate, and loading was done at 10 mL/min (18s residence time)

**Table 6:** Dynamic binding capacities (dbc) at 10% breakthrough for phenyl hydrophobic membrane. Chromatography Adsorber (3 ml Membrane Volume) Challenged with Three Different Model Proteins.

recombinant protein. The protein was stacked with shifting measures of ammonium sulfate such that the totals bound while the monomer did not. In the wake of stacking, the layer was washed with deionized water to elute the bound totals. The total evacuation expanded with expanded ammonium sulfate focus [54]. Protein yield was >95%, paying little mind to the salt conditions utilized within the stacking cushion [54].

## Transform Optimization with Hydrophobic Interaction Chromatography

In this area, a human mab (IgG1,  $p_i=8.3$ , 150 kda) was decontaminated utilizing the Phenyl hydrophobic layer chromatography adsorber in move through mode [47]. The protein of investment was handled at some association utilizing the Per.c6 unit line as a part of a bolstered group transform with a synthetically outlined development medium. The unrefined medium was elucidated by centrifugation at 15,000 g accompanied by profundity filtration and sterile filtration. The illuminated media was part of the way cleaned by section chromatography [54].

This material was stacked onto a 3-ml Phenyl nano layer adsorber in 50 mm sodium phosphate cushion, pH 7.0, with ammonium sulfate focuses extending from 0.1 to 0.4 M. The item was gathered in course through divisions and the film was then washed with equilibration cradle to recuperate the whole item. Specimens of the move through portions were examined for totals by size rejection chromatography / gel saturation chromatography (Sec/gpc).

It was watched that the coupling of elevated atomic weight (Hmw) totals enhanced with expanded salt focus in the stacking support [54]. In spite of the fact that 100% leap forward of totals was watched at 0.1 and 0.2 M  $(\text{NH}_4)_2\text{SO}_4$ , immersion was not arrived at 0.3 or 0.4 M  $(\text{NH}_4)_2\text{SO}_4$  when the film was stacked with up to 167 mg-mab/ml layer. Protein recuperation was more terrific than 94%, even at the most elevated salt state of 0.4 M  $(\text{NH}_4)_2\text{SO}_4$ , while total lessening for the aggregate move through pool was pretty nearly 50% [54]. In this study, a stacking limit of no less than 167 mg-Mab/ml layer was accomplished utilizing the Phenyl hydrophobic layer chromatography adsorber with yields of  $\geq 94\%$  and a last total level <1% for a monoclonal IgG1 [54].

## Observation of Hic Membrane with Resin

In an additional report, the decontamination of a monoclonal IgG1 communicated utilizing the Per.c61 human unit line was portrayed thinking about the Phenyl hydrophobic layer chromatography adsorber and Ppg-600m gum [55]. The Phenyl hydrophobic layer chromatography adsorber and Ppg-600m pitch demonstrated practically identical exhibition as far as yield and evacuation of totals and Hcps (Table 7). While the gum had a twofold higher limit, the film adsorber had a six fold shorter process duration. The Phenyl hydrophobic film chromatography adsorber gave another chance in downstream transforming: the capability to refine a recombinant

helpful protein in a bind/elute mode utilizing a film chromatography adsorber at mechanically applicable scales. The Hic layer showed a dynamic tying limit similar to presently ready Hic gums utilized as a part of numerous courses of action and demonstrated phenomenal determination, both with model proteins and a recombinant monoclonal immunizer prepared utilizing the Per. C6 human unit line. In the work talked about, a usable dynamic tying limit of 16 mg/cm<sup>3</sup> was accomplished with yields of 90%, Hcp decrease of give or take 80%, and last total levels beneath 1% for a monoclonal IgG1 [55]. The immune response utilized within the study was truly stable and consequently the total test to the adsorber was not especially domineering. The step was demonstrated to be adaptable over a 50-fold run, and it was additionally showed that the layer could be cycled no less than 10 times if required without any huge change in exhibition. These exhibition studies affirmed that the Phenyl hydrophobic layer chromatography adsorber was fit for pollution evacuation tantamount to that of a pressed couch segment however in an altogether shorter transforming time. The towering stream rate and insignificant dissemination restrictions took into consideration the fast preparing of clusters while disposing of the requirement for the pressing, capability, and cleaning acceptance mulls over connected with pressed couch chromatography. This brought about diminished process intricacy, work, and time while even now administering the needed item untaintedness. This clarifies that around the elective arrangements for refinement unit operations, layer chromatography innovation is starting to make a true effect in the biopharmaceutical business. Notwithstanding the favorable circumstances, the film adsorber additionally gives upgraded procedure adaptability (Table 7) [55].

## New Chemistries in Aex Membranes

One impediment of the layer chromatography adsorber that is predominant in cleaning operations is its affectability to towering salt conditions. Granted that infection freedom works well around unbiased pH and at conductivities up to and even past 10 ms/cm, the coupling limit of layer chromatography adsorber drops dynamically as conductivity expands, prompting wasteful tying of hazardous infections for example Mvm and the early leap forward of Hcps [56]. Sustain streams eluting from the Cex step hence must be weakened to lessen conductivity, expanding handling times, cushion prerequisites, and the measure of anxiety to which proteins are uncovered. Aex with layer chromatography adsorber ligands is a hearty stage that has been approved in many methods, inasmuch as plan B for example blended mode chromatography fail to offer a clear and powerful component that might permit straightforward execution throughout methodology growth [56].

## Salt-Tolerant Interaction Chromatography

The test of cleaning under towering salt conditions was tended to by improving a salt-safe Aex science that met all existing abilities of film chromatography adsorber ligands and could be adjusts to a layer chromatography arrange for cleaning requisitions [56]. The target ionic quality of roughly 16.8 ms/cm was decided to expand the

Features	Nano 3 cm <sup>3</sup>	XK16 9.65 cm <sup>3</sup>
Usable capacity (mg-MAB/cm <sup>3</sup> -membrane)	16	33
Recovery(%)	92	97
Elution [HMW] (%) (load = 0.9%)	0.4	0.7
Elution [HCP] ( $\mu\text{g-HCP/mg-MAB}$ )	1.6	2.1

**Table 7:** Comparison of the capacity, recovery, and impurity reduction of phenyl and PPG-600M.

outline space in procedure infrastructure. Consistent with the necessity determination for another standard in course through cleaning, both base framework and the ligands must be improved separately before distinguishing an optimal arrangement for the innovation [56].


Starting investigation of the variables constraining the exhibition of original layer chromatography adsorber indicated that the united hydrogel layer on the macroporous underpin given way at heightened salt focuses and could never again be entered by macromolecules and infections [57]. Along these lines, a second-era base help network containing a cross-joined, recovered cellulose layer with a novel structure was produced. The new lattice, called the Stic (salt-tolerant cooperation chromatography) had an altogether higher tying limit at elevated salt fixations and was less touchy to expanding salt focuses than layer chromatography adsorber [56].

A later study has indicated that the salt tolerance of an anion-trade network is dead set by the net charge of the ligand, its atomic structure, and the immobilization thickness [58]. It was likewise demonstrated that the amount of free essential amine amasses fundamentally impacted the capacity of a lattice to function in heightened salt focuses, by recompensing for limit impediment through expanded charge thickness [59]. Polycations with numerous  $\text{NH}_2$  bunches have been recognized productive ligands and have been utilized to uproot pathogens from blood [60]. Taking into account the above informative data, a polyallylamine ligand covalently coupled to the twofold permeable film was produced, and its contaminant evacuation exhibition at diverse ph and conductivity qualities was researched [56].

Granted that acidic contaminants for example nucleic acids and endotoxins could be evacuated effectively under definitive process conditions, this was not vitally the situation for host cell proteins and infections. It was consequently vital to recognize suitable most noticeably awful case models to copy the issue of right on time achievement under physiological conditions and explore the conduct of the new material in spiking trials [56]. Taking into account the above qualified data, a polyallylamine ligand covalently coupled to the twofold permeable layer was created, and its contaminant evacuation exhibition at diverse ph and conductivity qualities was examined [56].

Granted that acidic contaminants for example nucleic acids and endotoxins could be uprooted effectively under definitive process conditions, this was not vitally the situation for host cell proteins and infections. It was in this way vital to distinguish suitable most exceedingly awful case models to copy the issue of promptly achievement under physiological conditions and examine the conduct of the new material in spiking trials [56].

Upon examination, it was indicated that an advanced base back film lattice joined with a feeble anion-trade science dependent upon polyallylamine ligands furnished a powerful system for infection leeway at physiological conductivities or more (Figure 1) [56]. Stic and particle trade layer chromatography adsorber display a clear on-off anion-trade system that could be utilized for the first or second cleaning go after catch. Stic did not work with polyvalent cradle frameworks for example phosphate supports, however since polyvalent cushions are not proposed for catching offloads, this ought not be a foundation for concern in for the most part courses of action. Granted that Q chromatography might remain the workhorse in cleaning, Stic permitted cleaning to be done without an interstitial weakening step, which decreased methodology time and kept away from supplemental cradle readiness and hold steps. All the more imperatively, on the whole methodology wellbeing was expanded in view of the more proficient



**Figure 1:** Structure of the covalently attached polyallylamine ligand.

evacuation of less acidic contaminants. Studies are underway to exhibit the evacuation of Hcp under diverse transform conditions.

### Single-Use Option in Membrane Chromatography Adsorber

Single-utilization unit operations are coming to be more vital in bioprocessing as affirmed by an industry study [61]. For numerous unit operations, especially filtration and media/ support space, single-utilization units have been in as something to be shared use for a long while since they recover on cleaning and acceptance costs. Then again, single-utilization steps have different profits: they recover time, furnish adaptability, and streamline process infrastructure [11]. Limit and single-utilization choices are basic variables to acknowledge when figuring unit operation takes for new forms. In spite of the fact that layer mechanisms decidedly have a higher throughput, a straight illustration of pitches and layers dependent upon volume demonstrates that solitary utilization films are as of now more unmanageable. So this should be equalized against the decreased size of layer mechanisms, which likewise decreases support prerequisites, setting aside a few minutes shorter, and carrying along the remaining profits of a solitary utilize innovation [11]. The capacity to swap every module totally makes it simpler to collect transform prepares for new items in existing premises without stressing over cross-defilement, even though there could be extra approval trouble in view of drained materials.

Engage in film chromatography adsorber is developing on account of the triumph of single-utilize layer channels, however there is still an absence of gratefulness despite numerous preferences layer mechanisms offer in downstream handling [13,15]. The utilization of film units brings about the complete end of cleaning and approval, a major cost in downstream preparing. Nourishment and Drug Administration (FDA) regulations need the cleaning, support, and sterilization of settled gear and channeling at fitting interims to counteract glitches and defilement, yet this is unnecessary when fouled or depleted modules can basically be swapped out and displaced with new ones.

Ecological concerns have additionally been referred to as a cause for moving to single-utilize frameworks as a part of biomanufacturing [62]. This stems from the perception that in spite of the fact that solitary utilization frameworks need huge vigor to generate them and create plastic waste, the measure of vigor and water devoured in the generation of the water for infusion (Wfi) and steam utilized as a part of clean-in-place/steam set up operations can balance the waste issue. The point when waste-to vigor plants are acknowledged for transfer of the plastics, the ecological profits of single-utilize advances are further upgraded [63]. By one gauge, the drives of the plant workers explain more than 50% of the aggregate carbon outflows connected with biomanufacturing [62].

A later study depicts an extraordinary decontamination transform for a mab based on single-utilize advances [64]. No stuffed bunk

section chromatography was utilized as a part of the train. Rather, a mix of precipitation and bind/elute and move through (Ft) layer chromatography steps were brought about to decrease the level of Hcps to underneath 50 ppm and item identified debasements (total) to less than 1% fixation for the consequent Cex layer chromatography step. The last Hcp level of this material was <50 ppm for the modest scale analysis and  $\leq 10$  ppm for the middle scale, and total levels were <0.5% at both scales [64].

## Validation in Chromatography

Approval starts with exceptional process growth. It needs prepare planners to comprehend the need to outline a process that will be fit for eventually gathering decided beforehand particulars without being liable to deviations inside a demarcated go of preset working parameters [65].

Cleaning acceptance has been an issue referred to throughout a few assessments. Multiuse offices and chromatography forms appear to draw the most concern. Regularly producers have a tendency to disregard the approval studies for cleaning, reuse, lifetime, and space for the multiuse frameworks, as an aftereffect of which there is intense complaint from administrative firms. An evaluation of remainder and its dangers are significant components that ought to be incorporated in the approval study [65].

## Different Systems

Chromatography segments are utilized for numerous cycles, and it is crucial to exhibit that the cleaning methods after every cycle are sufficient to guarantee negligible danger of an expansion in remainder of tightly bound item or different contaminations from hurry to run. Before improving the cleansing technique, mind must be taken to guarantee that the chromatographic gums and supplies are good with the cleaning regimens set up [66]. Pitch lifetime studies utilizing scale-down models incorporate an assessment of segment honesty, section cleaning, remainder, chromatograms, item recuperation, item immaculacy profile, and contamination leeway (DNA, Hcp, infections). Numerous tests need to be utilized to analyze pre and post-use section honesty and characteristics of the sap [67].

In any case, even after fruition of a chromatography or filtration step, a segment of these pollutions can remain bound to the media and be gradually evacuated upon expanded space or even be continued to the following creation parcel. Further, approval of media life compass needs showing at scale of the focused on number of reuses. These studies are regularly prolonged and exorbitant [68].

## Single-use Systems

The utilization of layer mechanisms brings about the complete disposal of cleaning and approval, a major cost in downstream preparing. FDA regulations need the cleaning, upkeep and purification of settled supplies, and channeling at proper interims to avert glitches and tainting, however this is unnecessary when fouled or depleted modules can essentially be swapped out and traded with new ones [7]. The single-utilization film units might require confirmations of other acceptance studies for example the. Extractables, leachables, pre/post-use respectability tests and also preuse Sip information, which are ordinarily given by the supplier in the acceptance report as well as the quality certification and quality consistence authentications according to the all inclusive guidelines. This saves the end user a lot of time, effort, expense, and man-power, and provides them an opportunity for ready-to-go, plug-and-play systems.

## References

1. Thömmes J, Etzel M (2007) Alternatives to chromatographic separations. *Biotechnol Prog* 23: 42-45.
2. Gottschalk U (2005) New and Unknown Challenges Facing Biomanufacturing. *Biopharm Int* 18: 24-28.
3. BioPlan Associates and BioProcess Technology Consultants (2005) Third Annual Report and Survey of Biopharmaceutical Manufacturing Capacity and Production. Rockville, MD, USA.
4. Low D, O'Leary R, Pujar NS (2007) Future of antibody purification. *J Chromatogr B Analyt Technol Biomed Life Sci* 848: 48-63.
5. Shukla AA, Hubbard B, Tressel T, Guhan S, Low D (2007) Downstream processing of monoclonal antibodies--application of platform approaches. *J Chromatogr B Analyt Technol Biomed Life Sci* 848: 28-39.
6. Przybycien TM, Pujar NS, Steele LM (2004) Alternative bioseparation operations: life beyond packed-bed chromatography. *Curr Opin Biotechnol* 15: 469-478.
7. Gottschalk U (2008) Bioseparation in antibody manufacturing: the good, the bad and the ugly. *Biotechnol Prog* 24: 496-503.
8. Fraud N (2008) Membrane Chromatography: An Alternative to Polishing Column Chromatography. *Bioprocessing J* 7: 26-29.
9. Ghosh R (2002) Protein separation using membrane chromatography: opportunities and challenges. *J Chromatogr A* 952: 13-27.
10. Lim JAC, Sinclair A, Kim DS, Gottschalk U (2007) Economic Benefits of Single-Use Membrane Chromatography in Polishing. *BioProcess Int* 5: 60-64.
11. Mora J, Sinclair A, Delmdahl N, Gottschalk U (2006) Scaling up Disposable Membrane Chromatography. Performance Analysis and Economic Cost Model. *BioProcess Int* 4: 38-43.
12. Walter JK (1998) Strategies and Considerations for Advanced Economy in Downstream Processing of Biopharmaceutical Proteins in Bioseparation and Bioprocessing: Quality and Characterization, Economics, Safety and Hygiene 447-460.
13. Zhou JX, Tressel T (2006) Basic concepts in Q membrane chromatography for large-scale antibody production. *Biotechnol Prog* 22: 341-349.
14. Thömmes J, Kula MR (1995) Membrane Chromatography. An Integrative Concept in the Downstream Processing of Proteins. *Biotechnol Prog* 11: 357-367.
15. Etzel M, Riordan W (2006) Membrane Chromatography: Analysis of Breakthrough Curves and Viral Clearance. In *Process Scale Bioseparations for the Biopharmaceutical Industry*. Taylor & Francis 277-296.
16. Gebauer KH, Thömmes J, Kula MR (1997) Breakthrough Performance of High-Capacity Membrane Adsorbers in Protein Chromatography. *Chem Narrow Sci* 52: 405-419.
17. Norling L, Lute S, Emery R, Khuu W, Voisard M, et al. (2005) Impact of multiple re-use of anion-exchange chromatography media on virus removal. *J Chromatogr A* 1069: 79-89.
18. Farshid M, Taffs RE, Scott D, Asher DM, Brorson K (2005) The clearance of viruses and transmissible spongiform encephalopathy agents from biologicals. *Curr Opin Biotechnol* 16: 561-567.
19. Glynn J, Hagerty T, Pabst T, Annathur G, Thomas K, et al. (2009) The Development and Application of a Monoclonal Antibody Purification Platform. Advancing in Separation and Purification: Purifying High Titrers. *Biopharm Int* 22: 16-20.
20. Arunakumari A, Wang J, Ferreira G (2007) Improved Downstream Process Design for Human Monoclonal Antibody Production. *Biopharm Int* 20: 36.
21. Arunakumari A, Wang J, Ferreira G (2009) Advances in Non-Protein A Purification Processes for Human Monoclonal Antibodies, Advancing in Separation and Purification: Purifying High Titrers. *Biopharm Int Advances in Process Chromatography* 22: 22-26.
22. Curling J, Gottschalk U (2007) Process Chromatography: Five Decades of Innovation. *Biopharm Int* 20: 10-19.
23. Knudsen HL, Fahrner RL, Xu Y, Norling LA, Blank GS (2001) Membrane ion-exchange chromatography for process-scale antibody purification. *J Chromatogr A* 907: 145-154.



24. Curtis S, Lee K, Blank GS, Brorson K, Xu Y (2003) Generic/matrix evaluation of SV40 clearance by anion exchange chromatography in flow-through mode. *Biotechnol Bioeng* 84: 179-186.
25. Giovannoni L, Ventani M, Gottschalk U (2008) Antibody Purification Using Membrane Adsorbers. *BioPharm Int* 21: 48-52.
26. Zhou JX, Tressel T (2005) Membrane Chromatography as a Robust Purification System for Large-Scale Antibody Production. *BioPharm Int* 22: 32-37.
27. Kelley B (2007) Very large scale monoclonal antibody purification: the case for conventional unit operations. *Biotechnol Prog* 23: 995-1008.
28. Kamarck ME (2006) Building biomanufacturing capacity--the chapter and verse. *Nat Biotechnol* 24: 503-505.
29. Gottschalk U (2005) Downstream Processing of Monoclonal Antibodies: From High Dilution to High Purity. *Biopharm Int* 18: 42-58.
30. Specht R, Han B, Wickramasinghe SR, Carlson JO, Czermak P, et al. (2004) Densonucleosis virus purification by ion exchange membranes. *Biotechnol Bioeng* 88: 465-473.
31. Teeters MA, Conrardy SE, Thomas BL, Root TW, Lightfoot EN (2003) Adsorptive membrane chromatography for purification of plasmid DNA. *J Chromatogr A* 989: 165-173.
32. Zhang S, Krivosheyeva A, Nochumson S (2003) Large-scale capture and partial purification of plasmid DNA using anion-exchange membrane capsules. *Biotechnol Appl Biochem* 37: 245-249.
33. Grunwald AG, Shields MS (2001) Plasmid purification using membrane-based anion-exchange chromatography. *Anal Biochem* 296: 138-141.
34. Kuiper M, Sanches RM, Walford JA, Slater NK (2002) Purification of a Functional Gene Therapy Vector Derived from Moloney Murine Leukaemia Virus Using Membrane Filtration and Ceramic Hydroxyapatite Chromatography. *Biotechnol Bioeng* 80: 445-453.
35. Deshmukh RR, Warner TN, Hutchison F, Murphy M, Leitch WE 2nd, et al. (2000) Large-scale purification of antisense oligonucleotides by high-performance membrane adsorber chromatography. *J Chromatogr A* 890: 179-192.
36. Kalbfuss B, Wolff M, Geisler L, Tappe A, Wickramasingh R, et al. (2007) Direct Capture of Influenza A Virus from Cell Culture Supernatant with Sartobind Anion-Exchange Membrane Adsorbers. *J Membrane Science* 299: 251-260.
37. Vicente T, Peixoto C, Carrondo MJ, Alves PM (2009) Purification of recombinant baculoviruses for gene therapy using membrane processes. *Gene Ther* 16: 766-775.
38. Vicente T, Souza MFQ, Peixoto C, Mota JPB, Alves PM, et al. (2008) Anion-Exchange Membrane Chromatography for Purification of Rotavirus-Like Particles. *J Membr Sci* 311: 270-283.
39. Zhou J, Dehghani H (2007) Development of Viral Clearance Strategies for Generous Scale Monoclonal Antibody Production. In *Advances in Generous Scale Biomanufacturing and Scale-Up Production*. 558-586.
40. Zhou JX, Tressel T, Gottschalk U, Solamo F, Pastor A, et al. (2006) New Q membrane scale-down model for process-scale antibody purification. *J Chromatogr A* 1134: 66-73.
41. Neidhardt EA, Luther MA, Recny MA (1992) Rapid, two-step purification process for the preparation of pyrogen-free murine immunoglobulin G1 monoclonal antibodies. *J Chromatogr* 590: 255-261.
42. Diehl T (2006) Presentation "Application of Membrane Chromatography in the Purification of Humanly Monoclonal Antibodies." *Downstream Technology Forum*, King of Prussia.
43. Zhou JX, Solamo F, Hong T, Shearer M, Tressel T (2008) Viral clearance using disposable systems in monoclonal antibody commercial downstream processing. *Biotechnol Bioeng* 100: 488-496.
44. Fraud N, Kuczewski M, Zarbis-Papastoitis G, Hirai M (2009) Hydrophobic Membrane Adsorbers for Large-Scale Downstream Processing. *BioProcess. Int*.
45. Evans DR, Macniven RP, Labanca M, Walker J, Notarnicola SM (2008) Purification of an Fc-fusion biologic: clearance of multiple product related impurities by hydrophobic interaction chromatography. *J Chromatogr A* 1177: 265-271.
46. Gagnon P, Grund E, Lindback T (1995) Large-Scale Process Development For Hydrophobic Interaction Chromatography, Part 1: Gel Selection and Development of Binding Conditions. *LC-GC* 13: 318-328.
47. Gagnon P, Grund E, Lindback T (1995) Large-Scale Process Development For Hydrophobic Interaction Chromatography, Part 2: Controlling Process Variation. *LC-GC* 13: 400-410.
48. Gagnon P, Grund E (1996) Large-Scale Process Development For Hydrophobic Interaction Chromatography, Part 3: Factors Affecting Capacity Determination. *Biopharm Int* 9: 34-39.
49. Gagnon P, Grund E (1996) Large-Scale Process Development For Hydrophobic Interaction Chromatography, Part 4: Controlling Selectivity. *Biopharm Int* 9: 54-64.
50. Li F, Zhou JX, Yang X, Tressel T, Lee B (2005) Current Therapeutic Antibody Production and Process Optimization. *Bioprocess J* 4: 23-30.
51. Jungbauer A, Machold C, Hahn R (2005) Hydrophobic interaction chromatography of proteins. III. Unfolding of proteins upon adsorption. *J Chromatogr A* 1079: 221-228.
52. Gottschalk U, Fischer-Fruehholz S, Reif O (2004) Membrane Adsorbers: A Cutting Edge Process Technology at the Threshold. *Bioprocess Int* 2: 56-65.
53. Liu X, Colling M, Fraud N, Campbell J, Lowenstein I, et al. (2009) Upcoming Technologies to Facilitate More Efficient Biologics Manufacturing. *Biopharm Int* 22: 38-51.
54. Fraud N, Faber R, Kiss C, Demmer W, Hoerl HH, et al. (2009) Hydrophobic-Interaction Membrane Chromatography for Large-Scale Purification of Biopharmaceuticals. *BioProcess Int* 22: 30-35.
55. Kuczewski M, Fraud N, Faber R, Zarbis-Papastoitis G (2010) Development of a polishing step using a hydrophobic interaction membrane adsorber with a PER.C6-derived recombinant antibody. *Biotechnol Bioeng* 105: 296-305.
56. Faber R, Yang Y, Gottschalk U (2009) Salt-Tolerant Interaction Chromatography for Large-Scale Polishing with Convective Media. *BioPharm Int* 11-14.
57. Tatárová I, Fáber R, Denoyel R, Polakovic M (2009) Characterization of pore structure of a strong anion-exchange membrane adsorbent under different buffer and salt concentration conditions. *J Chromatogr A* 1216: 941-947.
58. Riordan W, Heilmann S, Brorson K, Seshadri K, He Y, et al. (2009) Design of Salt-Tolerant Membrane Adsorbers for Viral Clearance. *Biotechnol Eng* 103: 920-929.
59. Johansson B, Belew M, Eriksson S, Glad G, Lind O, et al. (2003) Preparation and Characterization of Prototypes for Multi-Modal Separation Media Aimed at the Capture of Negatively Charged Biomolecules at High Salt Conditions. *J Chromatogr A* 1016: 21-33.
60. Onishi M, Ohwada T, Hadanoshi K, Isehara-shi K, Ohmura Y, et al. (2009) A Pathogenic Substance Removing Material and a Blood Filter Comprising Said Material, European Patent Specification EP 0586268B1.
61. BioPlan Associates and BioProcess Technology Consultants (2006) Fourth Annual Report and Survey of Biopharmaceutical Manufacturing Capacity and Production.
62. Leveen L, Cox S (2008) Analyzing the Carbon Footprint Left by a Company's Use of Disposables. Paper presented at IBC's 5th International Single-Use Applications for Biopharmaceutical Manufacturing, San Diego, CA.
63. Newman D, Walker S (2008) The Environmental Dilemma of Single-Use Manufacturing. Paper presented at IBC's 5th International Single-Use Applications for Biopharmaceutical Manufacturing, San Diego, CA.
64. Kuczewski M, Schirmer E, Lain B, Zarbis-Papastoitis G (2011) A single-use purification process for the production of a monoclonal antibody produced in a PER.C6 human cell line. *Biotechnol J* 6: 56-65.
65. Sofer G (2005) Guidelines to Process Validation. In *Process Validation in Manufacturing of Biopharmaceuticals* 1-11.
66. Adner N, Sofer G (1994) Biotechnology Product Validation. III. Chromatography Cleaning Validation. *BioPharm* 7: 44-48.
67. Godavarti R, Petrone J, Robinson J, Wright R, Kelley B (2005) Scale-Down Models for Purification Processes: Approaches and Applications." In *Process Validation in Manufacturing of Biopharmaceuticals* 69-142.
68. Rathore AS, Sofer G (2005) Life Span Studies for Chromatography and Filtration Media. In *Process Validation in Manufacturing of Biopharmaceuticals*. Taylor & Francis 169-203.