

Vector Control for Mosquito in Ethiopia: A Review Article

Gebrehiwet Tesfahuneygn* and Gebremichael Gebreegziabher

Tigray Health Research Institute, Mekelle, Ethiopia

*Corresponding author: Gebrehiwet Tesfahuneygn, Tigray Health Research Institute, Mekelle, Ethiopia, Tel: +251914240008; E-mail: gtlab2006@gmail.com

Received date: November 22, 2018; Accepted date: January 18, 2019; Published date: January 29, 2019

Copyright: © 2019 Tesfahuneygn G, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License; which permits unrestricted use; distribution; and reproduction in any medium; provided the original author and source are credited.

Abstract

Vector transmitted diseases remains a serious public health burden in the world. The current core methods used for malaria vector control interventions worldwide are Indoor Residual Spraying (IRS) and long-lasting insecticidal nets, with Larval Source Management (LSM) applicable in certain settings where mosquito breeding sites are few, fixed and findable. Mosquitoes transmit many diseases including malaria, dengue fever, Japanese encephalitis virus, West Nile virus, yellow fever virus, and filariasis. Of these, malaria transmitted primarily by *Anopheles gambiae*, dengue transmitted by *Aedes aegypti* and lymphatic filariasis transmitted by *Culex quinquefasciatus* are the most devastating problems in terms of the global number of people affected.

Insecticide-treated bed nets and indoor residual spraying have been widely used as front-line tools against malaria vectors in endemic African regions. These preventive measures are highly effective against malaria vectors, which prefer to bite and rest inside the rooms. Some of the adulticide used for mosquito control include products derived from microorganisms, plants or minerals, synthetic molecules, organophosphates, some natural pyrethrins, or synthetic pyrethroids.

Keywords: Mosquito; Malaria; WHO; Vector control strategies

Introduction

Malaria remains a serious public health burden in endemic regions. The World Health Organization (WHO) estimates that there were 214 million new cases of the disease and more than 445,000 malaria-related deaths in 2016, with 91% of these occurring in Africa [1]. As there are no commercially available vaccines against this disease, vector control remains crucial to reduce disease transmission. Vector control strategies have traditionally focused on killing mosquitoes using a variety of insecticides. Environmental management (through reduction or removal of mosquito breeding sites) has often been used alongside chemical or microbiological ovicides, larvicides, and pupicides areas where endemic mosquito-borne diseases occur. The Global Fund and the President's Malaria Initiative (PMI) are supporting the scaling up of long-Lasting Insecticidal Nets (LLINs) and Indoor Residual Spraying (IRS) in many endemic African countries [2]. Long-Lasting Insecticidal Nets (LLINs) have contributed significantly to the success of malaria control in such malaria-endemic countries. Since 2000, malaria mortality rates have fallen by 66% among all age groups and by 71% among children under 5 years of age [3,4]. According to WHO estimates, the incidence of malaria (i.e. the rate of new malaria cases) fell by 18% between 2010 and 2016 [1].

The current core methods used for malaria vector control interventions worldwide are Indoor Residual Spraying (IRS) and long-lasting insecticidal nets, with Larval Source Management (LSM) applicable in certain settings where mosquito breeding sites are few, fixed and findable [5,6]. Historically, IRS has been the primary intervention method in vector control efforts [5,6]. Since its introduction as a vector control tool in 1945, IRS has proven successful in reducing the prevalence and incidence of malaria by reducing the level of transmission through killing or repelling malaria vectors.

Diseases Spread by Mosquitoes

Mosquitoes transmit many diseases including malaria, dengue fever, Japanese encephalitis virus, West Nile virus, yellow fever virus, and filariasis. Of these, malaria transmitted primarily by *Anopheles gambiae*, dengue transmitted by *Aedes aegypti* and lymphatic filariasis transmitted by *C. quinquefasciatus* are the most devastating problems in terms of the global number of people affected. Approximately half of the world's population are at risk of malaria, with 225 million cases being recorded in 2009 [7]. Dengue fever, a flavivirus transmitted by *Aedes* mosquitoes (mainly *A. gypti* and *A. albopictus*), is an increasingly serious public health problem in over 100 countries with some 2.5 billion people at risk [8]. *C. quinquefasciatus* is a major biting nuisance in the urban tropics, and is a vector for West Nile virus in addition to lymphatic filariasis, which affects 120 million people worldwide, and can lead to genital damage and elephantiasis. Apart from the obvious health issues, the economic and environmental costs are high. Such diseases have detrimental effects on fertility, population growth, saving and investment, worker productivity and absenteeism, premature mortality and medical costs. Countries with high malaria transmissions have had historically lower economic growth than countries without malaria.

Insecticide-Treated Bed Nets (ITNs) and Indoor Residual Spraying (IRS) have been widely used as front-line tools against malaria vectors in endemic African regions [9]. These preventive measures are highly effective against malaria vectors, which prefer to bite and rest inside the rooms [10]. Use of ITNs and IRS over the last decade has led to a significant decrease in malaria transmission in many areas of sub-Saharan Africa.

Malaria vector control strategies target both the immature and adult stages of *Anopheles* populations. Before World War II, anti-larval malaria control was the preferred strategy to control malaria. The intervention strategies predominantly included physical elimination of

mosquito breeding sites, biological control using larvivorous fish and chemical larviciding with Paris green [11]. Today, larval control is used and promoted to limit the development of *Anopheles* larvae in breeding sites [12] using biological larvicides such as *Bacillus thuringiensis israelensis* (Bti) that were first developed 30 years ago [13]. However, larval control is not particularly effective, especially in case of some *Anopheles* species for which breeding sites are extremely numerous and inaccessible.

Larval Source Management

Larval Source Management (LSM) refers to the targeted management of mosquito breeding sites, with the objective to reduce the number of mosquito larvae and pupae. LSM is only recommended as a supplementary malaria vector control measure; it should not be used to replace core vector control interventions, such as Long-Lasting Insecticidal Nets (LLINs) and Indoor Residual Spraying (IRS). LSM includes 1) habitat modification, which means a permanent alteration to the environment, e.g. land reclamation or surface water drainage; 2) Habitat manipulation, which refers to a recurrent activity e.g. water-level manipulation, flushing of streams, the shading or exposure of habitats; 3) larviciding, which involves the regular application of a biological or chemical insecticide to water bodies; and 4) biological control, which refers to the introduction of natural predators into water bodies, for example, predatory fish or invertebrates.

In general, LSM programmes need to be fully tailored to local environmental conditions and should be based on comprehensive feasibility and cost-effectiveness studies. Similarly to core vector control interventions, the management of larval habitats is a major financial and technical undertaking, requiring both community support and long-term political commitment.

Adulticides technique is usually less efficient for mosquito control. However, it is the only way to kill adult mosquitoes and is the last line of defense in reducing mosquito populations. Some of the adulticide used for mosquito control include products derived from microorganisms, plants or minerals, synthetic molecules, organophosphates, some natural pyrethrins, or synthetic pyrethroids [14].

Chemical Insecticides

Chemical insecticides have represented the widest method used to control mosquito-borne vectors. However, the effects of chemical insecticides on mosquito vector populations are usually transitory because vectors can rapidly develop resistance against them. Some of the disadvantages that generate when using only chemical products are (a) the selection of new insecticide resistance in pest populations; (b) the resurgence of already treated populations; (c) the generation of waste, risks, and legal complications; (d) the destruction of beneficial species; and (e) the high costs in equipment, labor, and material. In addition, the highly toxic and no biodegradable properties of insecticides and waste generated in the soil, water, food, and crops that affect public health are additional reasons to search new methods to help solve the problems caused by chemical insecticides [15]. Consequently, the concept of integrated control arises, a method in which pest and diseases control are performed using chemicals, useful organisms, and cultural practices.

The progress of science and the chemical industry in the nineteenth century, with the discovery of DDT, made possible the development and emergence of new conventional insecticides or so-called of

synthesis [16]. The most used of these insecticides of synthesis are modulators of sodium channels (organochlorines, pyrethroids, and pyrethrins), acetylcholinesterase inhibitors (carbamates and organophosphates), and the chloride channel antagonists regulated by the Gamma-Aminobutyric Acid or also known as GABA (organochlorine cyclodiene phenylpyrazoles).

Using these conventional insecticides gave positive results against insect's disease vectors at first. However, due to its massive use, insects soon began to develop resistance to them. Thus, an insecticide that initially was effective, just being useless in the long term. In response to this problem, new-generation insecticides also called bio-rational insecticides have been developed, whose research strategy is based on a good understanding of the physiological processes or mechanisms specific communication of insects, and in obtaining agents that are able to affect them. These products are divided into the following: those who are analogs of juvenile hormone and molting, inhibitors tissue formation, pheromones, insecticides that prevent hatching, and biological insecticides [17].

Plants and their Derivates

For centuries, nature has created several active substances that, when applied correctly, can control insect pests such as mosquito in an efficient manner. The use of plants by a man with insecticide purposes dates back to early human history. Due to their environmental advantages, the use of insecticides of vegetable origin in pest management has been increasing [18]. Among plants with potential activity against mosquitoes, Nim or Neem (*Azadirachta indica*) causes stunted growth, loss of appetite, reduction of fertility, molting disorders, morphological defects, and behavioral changes [19]. Moreover, it has been demonstrated that raw or partially purified plant extracts are most effective for mosquito control in place of the purified compounds or extracts [20]. The snuff (*Nicotiana tabacum*) is used, thanks to its insecticide and insect repellent action, where nicotine acts on the nervous system of insects through breathing, ingestion, and contact [21]. Other plants from which oils are extracted are garlic (*Allium sativum*) and cinnamon (*Cinnamomum osmophloeum*), which have been shown to have insecticidal properties against larvae and adults of *A. albopictus*, *Culex quinquefasciatus*, and *Armigeressubalbatus*.

Biological Agents

Among biological agents used for mosquito control can be mentioned derivatives of viruses, bacteria, and fungi. The entomopathogenic virus spreads from one insect generation to the next causing paralysis and eventually died on mosquito larvae being more effective in the first stage of development [22]. Within bacteria, only reports of *Bacillus thuringiensis*, *B. sphaericus*, and *B. popilliae* with possibilities to exercise control over dipterans insects currently exist. These bacteria, during the sporulation process, produce protein crystals with insecticidal effect and/or some toxins with the same effect [23]. *Bacillus* initially causes diarrhea and intestinal paralysis in the mosquito, giving rise to a decrease in body movements, convulsions, and general paralysis. Internally, within the mosquito stomach, *B. thuringiensis* releases toxic crystals that paralyze the insect gut stopping peristalsis, causing that the insect stop feeding and die by starvation. Within the gut, bacteria multiply until they break the epithelium and invade the rest of the insect body. However, its use for mosquito control is scarce and presents some drawbacks as its duration in the environment is limited, its dispersion is rather inefficient, and

the susceptibility to bacterial infection in the pest population is very heterogeneous. There are very sensitive individuals and other highly resistant. Fungi are other microorganisms that may be used to control mosquito vectors, of which 400 species are known with insecticide potential. About 20 of them have been given more attention, including those in the *Lagenidium*, *Entomophaga*, *Neozygites*, *Entomophobia*, *Erynia*, *Aschersonia*, *Verticillium*, *Nomuraea*, *Hirsutella*, *Metarhizium*, *Beauveria*, and *Paecilomyces* genera [24]. Although entomopathogenic fungi are not as specific as bacteria or viruses, spores persist and infect insect successive generations, so that when the infection is established, its effects can last several years. Infection occurs by adhesion of the spores on the insect cuticle, where these germinate and penetrate the cuticle leading to insect colonization by mycelium. Cuticle penetration occurs through the use of an enzyme complex that the fungi use to feed. The entomopathogenic fungus most used in controlling mosquito infestations is *Beauveria bassiana*, which produces various active ingredients such as beauvericin [25].

The biological control of mosquito larvae with predators and other biological control agents could be a more effective and environmentally friendly strategy, thus avoiding the use of synthetic chemicals and the consequent environmental damage [26]. Among them, some insects and vertebrates such as fish, amphibians, and some mammals have the potential to control mosquito disease-vector populations. Within vertebrates, amphibians, bats, and fish have been used to control populations of mosquito. For example, using larvivorous fish species, control of mosquito larvae in deposits used to store water has been achieved [27]. Moreover, bats are responsible for capturing flying insects such as mosquitoes at night; similarly, toads and frogs consume large numbers of insects, slugs, worms, and other invertebrates [28]. However, the use of frogs and tadpoles for disease vector control is still largely unexplored.

Conclusion

Mosquito-borne diseases remain a major cause of morbidity and mortality across tropical regions. The current core methods used for malaria vector control interventions worldwide are indoor residual spraying and long-lasting insecticidal nets, with larval source management applicable in certain settings where mosquito breeding sites are few, fixed and findable.

Competing Interests

The authors declare that they have no competing interests.

Authors' Contribution

GT and GG wrote and edited the manuscript. All authors read and approved the final manuscript.

References

- World Health Organization (2017) World malaria report WHO, Geneva.
- Bhattarai A, Ali AS, Kachur SP, Martensson A, Abbas AK, et al. (2007) Impact of artemisinin-based combination therapy and insecticide-treated nets on malaria burden in Zanzibar. PLoS Medicine 4: e309.
- Ceesay SJ, Casals-Pascual C, Nwakanma DC, Walther M, Gomez-Escobar N, et al. (2010) Continued decline of malaria in The Gambia with Implications for elimination. PLoS One 5: e12242.
- WHO (2016) World malaria report. World Health Organization, Geneva.
- WHO (2014) World Malaria Report. World Health Organization, Geneva.
- WHO (2012) Global plan for insecticide resistance management in malaria vectors: World Health Organization Global Malaria Programme.
- WHO (2010) World Malaria report. Geneva, Switzerland.
- WHO (2008) Dengue and dengue haemorrhagic fever.
- Hemingway J, Ranson H, Magill A, Kolaczinski J, Fornadel C, et al. (2016) Averting a malaria disaster: Will insecticide resistance derail malaria control? The Lancet 387: 1785-1788.
- Scott TW, Takken W (2012) Feeding strategies of anthropophilic mosquitoes result in increased risk of pathogen transmission. Trends Parasitol 28: 114-121.
- Mabaso MLH, Sharp B, Lengeler C (2004) Historical review of malarial control in Southern African with emphasis on the use of indoor residual house-spraying. Trop Med Int Health 9: 846-856.
- Dambach P, Schleicher M, Stahl H-C, Traoré I, Becker N, et al. (2016) Routine implementation costs of larviciding with *Bacillus thuringiensis israelensis* against malaria vectors in a district in rural Burkina Faso. Malar J 15: 380.
- Becker N, Margalit J (1993) Use of *Bacillus thuringiensis israelensis* against mosquitoes and blackflies. *Bacillus thuringiensis*. In: Entwistle PF, Cory JS, Bailey MJ, Higgs S, editors. *Bacillus thuringiensis*, an environmental biopesticide: Theory and practice. Chichester: Wiley and Sons pp. 147-170.
- Fathy KH (2012) Ecosmart biorational insecticides: alternative insect control strategies. Insecticides-Adv Integr Pest Manage InTech.
- Koul O, Walia S, Dhaliwal GS (2008) Essential oils as green pesticides: potential and constraints. Biopesticide Int 4: 63-84.
- Díez DM (2013) Effect of metoprenol on metamorphosis of the cockroach, *Blattellagermanica*. Vizcaya: University of the Basque Country.
- Mamatha DM, Kanji VK, Cohly H, Rajeswara RM (2008) Juvenile hormone analogues, methoprene and fenoxycarb dose dependently enhance certain enzyme activities in the silkworm *Bombyx mori* (L). Int J Environ Res Public Health 5: 120-124.
- Leyva M, MarquettiMdel C, French L, Montada D, Tiomno O, et al. (2013) Effect of a turpentine oil obtained from *Pinustropicalis* Morelets on the biology of a strain of *Aedes* (*Stegomyia*) *aegypti* Linnaeus, 1762 resistant to insecticides. Ann Biol 35: 75-84.
- Wandscheer CB, Duque JE, da Silva MAN, Fukuyama Y, Wohlke JL, et al. (2004) Larvicidal action of ethanolic extracts from fruit endocarps of *Melia azedarach* and *Azadirachta indica* against the dengue mosquito *Aedes aegypti*. Toxicon 44: 829-835.
- Abdelouaheb A, Nassima R, Noureddine S (2009) Larvicidal activity of a neem tree extract (*Azadirachtin*) against mosquito larvae in the Republic of Algeria. Jordan J Biol Sci 2: 15-22.
- Brechelt A. Manejo Ecológico de Plagas y Enfermedades (2004) Network of action in plagicidasysus alternativaspara américa latina (RAP-AL). Santiago, Chile, Chile 4-35.
- Primo YE. Mundi press ed. (1991) Ecologychemistry: New methods of fight against insects. Madrid pp: 191.
- Badii MH, Abreu JL (2006) Control biológico una forma sustentable de control de plagas. Daena: Int J Good Conscience 1: 82-89.
- Van Driesche RG, Bollows TS (1996) Biological control. Chapman and Hall, New York pp: 539.
- García GC, Tamez GP (2012) Bioinsecticide market in Mexico. Course on organic and sustainable agriculture. Produce Sinaloa Foundation pp: 99-114.
- Kumar R, Hwang JS (2006) Larvicidal efficiency of aquatic predators: a perspective for mosquito biocontrol. Zoo Stud 45: 447-466.
- Connelly CR, Bolles E, Culber D, De Valerio J, Donahoe M, et al. (2014) Florida Resident's Guide to mosquito control, integrated pest management for mosquito reduction around homes and neighborhoods. University of Florida, USDA-NIFA 22-37.

28. Mahr LD, Whitaker P, Ridgway N (2008) Biological control of insects and mites: An introduction to beneficial natural enemies and their use in pest management, UW Extension, Wisconsin Madison.