



Migraine: Understanding Its Causes, Symptoms, and Management

Sterling Beaumont*

Department of Neurological Sciences, Western Pacific University, Seattle, United States

DESCRIPTION

Migraine is a neurological condition that can cause repeated episodes of moderate to severe headache along with a range of other physical and sensory symptoms. It affects people of many ages and can interfere with employment, education, family responsibilities, sleep, and social activities. A migraine attack may last for several hours and, in some cases, continue for a few days. Although headache is one of its most recognized features, migraine is not simply an ordinary headache. It involves changes in how the nervous system processes pain and sensory information [1].

Migraine symptoms can vary considerably from one person to another. Some individuals experience a throbbing or pulsating pain, often on one side of the head, although pain can occur on both sides. Physical movement may make the discomfort worse. Nausea, vomiting, sensitivity to light, sensitivity to sound, and sensitivity to smells are also common. Some people prefer to rest in a quiet, dark room because ordinary environmental stimuli can become difficult to tolerate during an attack [2].

Certain individuals experience an aura before or during a migraine episode. Aura consists of temporary neurological symptoms that may affect vision, sensation, speech, or other functions. Visual changes are common and may include flashes of light, zigzag patterns, blind spots, or changes in the field of vision. Some people experience tingling or numbness, while others may have difficulty finding words. Aura usually develops gradually and resolves within a limited period. Because similar symptoms can occur with other neurological conditions, new or unusual neurological symptoms should receive medical evaluation [3].

Stress is often connected with migraine, but this does not mean that migraine is simply a psychological condition. Emotional stress can influence the nervous system and may contribute to an attack in susceptible individuals. Interestingly, some people report an attack after a stressful period ends rather than during the stressful event itself. Relaxation exercises, regular physical

activity, sufficient sleep, and healthy routines may help some people manage factors associated with migraine [4].

Treatment generally has two major goals: Reducing symptoms during an attack and decreasing the frequency or severity of future attacks. The appropriate approach depends on the person's symptoms, attack frequency, medical history, and other medicines being taken. For occasional attacks, healthcare professionals may recommend medicines designed to relieve pain or other symptoms. Some people with more frequent or disabling migraine may benefit from preventive treatment taken regularly [5].

Several types of medicines may be used for migraine. Depending on the individual situation, options can include certain pain-relieving medicines, triptans, anti-nausea medicines, and preventive medicines from several drug classes. Newer preventive treatments that target calcitonin gene-related peptide pathways are also available in some countries. The choice of treatment should be made with a qualified healthcare professional because medicines can have side effects, interactions, and specific restrictions [6].

Medication use itself requires attention. Frequently taking acute headache medicines can contribute to medication-overuse headache in some individuals. This can create a cycle in which headaches become more frequent and medicines are used increasingly often. People who need headache medicine on many days should discuss their pattern with a healthcare professional rather than repeatedly increasing self-treatment [7].

Lifestyle habits can provide useful support alongside medical care. Maintaining a consistent sleep schedule, drinking sufficient fluids, eating regular meals, and engaging in suitable physical activity may help reduce the likelihood of attacks for some individuals. Excessive caffeine can be associated with headaches in some people, while abrupt withdrawal can also produce headache. Therefore, people who consume caffeine regularly may benefit from discussing gradual changes with a healthcare professional if they wish to reduce intake [8,9].

Correspondence to: Sterling Beaumont, Department of Neurological Sciences, Western Pacific University, Seattle, United States, E-mail: sterling.beaumont27@archmail.us

Received: 12-May-2026, Manuscript No. BDT-26-32569; **Editor assigned:** 14-May-2026, PreQC No. BDT-26-32569; **Reviewed:** 27-May-2026, QC No. BDT-26-32569; **Revised:** 03-Jun-2026, Manuscript No. BDT-26-32569; **Published:** 10-Jun-2026, DOI: 10.35248/2168-975X.26.15.338

Citation: Beaumont S (2026). Migraine: Understanding Its Causes, Symptoms, and Management. Brain Disord Ther. 15:338.

Copyright: © 2026 Beaumont S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Diagnosis is usually based on the person's medical history, symptom pattern, and physical or neurological examination. There is no single routine laboratory test that confirms migraine. In some cases, imaging or other investigations may be recommended when symptoms are unusual or when the clinician needs to exclude another condition. Accurate diagnosis is useful because several headache disorders can appear similar but require different approaches [10].

CONCLUSION

Migraine is a complex neurological condition with symptoms that can extend far beyond head pain. Although its effects can be disruptive, appropriate diagnosis and treatment can help many people reduce the frequency, intensity, and impact of attacks. Paying attention to symptoms, maintaining regular health habits, recording possible patterns, and seeking professional medical advice can support better long-term management. Each person's migraine experience is different, so care should be based on individual symptoms, health needs, and professional assessment rather than assumptions about what works for everyone.

REFERENCES

1. Pietrobon D, Moskowitz MA. Pathophysiology of migraine. *Annu Rev Physiol.* 2013;75:365-391.
2. Lipton RB, Diamond S, Reed M, Diamond ML, Stewart WF. Migraine diagnosis and treatment: Results from the American Migraine Study II. *Headache.* 2001;41(7):638-645.
3. Goadsby PJ, Lipton RB, Ferrari MD. Migraine-current understanding and treatment. *N Engl J Med.* 2002;346(4):257-270.
4. Burstein R, Nosedà R, Borsook D. Migraine: Multiple processes, complex pathophysiology. *J Neurosci.* 2015;35(17):6619-6629.
5. Viana M, Linde M, Sances G, Ghiotto N, Guaschino E, Allena M, et al. Migraine aura symptoms: Duration, succession and temporal relationship to headache. *Cephalalgia.* 2016;36(5):413-421.
6. Silberstein SD. Practice parameter: Evidence-based guidelines for the migraine headache (an evidence-based review) and [RETIRED] Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology.* 2000;55 (6): 754-762.
7. Eigenbrodt AK, Ashina H, Khan S, Diener HC, Mitsikostas DD, Sinclair AJ, et al. Diagnosis and management of migraine in ten steps. *Nat Rev Neurol.* 2021;17(8):501-514.
8. Silberstein SD, Holland S, Freitag F, Dodick DW, Argoff C, Ashman E. Evidence-based guideline update: Pharmacologic treatment for episodic migraine prevention in adults: Report of the Quality Standards Subcommittee of the American Academy of Neurology and the American Headache Society. *Neurology.* 2012;78(17):1337-1345.
9. Ashina M, Katsarava Z, Do TP, Buse DC, Pozo-Rosich P, Özge A, et al. Migraine: Epidemiology and systems of care. *Lancet.* 2021;397(10283):1485-1495.
10. Guyuron B, Kriegler JS, Davis J, Amini SB. Comprehensive surgical treatment of migraine headaches. *Plast Reconstr Surg.* 2005;115(1): 1-9.