



Tauopathy: Understanding Tau-Related Brain Disorders and Their Clinical Effects

Renjiro Tsukishima*

Department of Neurological Sciences, Hoshizora Center for Brain Research, Yokohama, Japan

DESCRIPTON

Tauopathies are a group of neurological disorders associated with abnormal changes involving a protein called tau. Tau is normally found inside nerve cells, where it helps maintain the structure of microtubules. Microtubules are part of the internal transport system of neurons and help move materials within these cells. In tauopathies, tau can become abnormally modified, change its normal behavior, and accumulate inside nerve cells or supporting brain cells. These changes can interfere with cellular function and contribute to progressive damage in affected areas of the nervous system.

Several neurological diseases are classified as tauopathies. Alzheimer's disease is associated with tau accumulation as well as other pathological changes. Other conditions in which tau plays a major role include progressive supranuclear palsy, corticobasal degeneration, Pick's disease, and some forms of frontotemporal dementia. Although these disorders share abnormalities involving tau, their symptoms, affected brain regions, progression, and microscopic features can differ considerably.

Tau exists in different molecular forms. In the adult human brain, tau proteins can contain either three or four microtubule-binding repeats, commonly referred to as 3R and 4R tau. Different tauopathies can show different proportions of these forms. Progressive supranuclear palsy and corticobasal degeneration are commonly associated with predominantly 4R tau, while Pick's disease is associated mainly with 3R tau. Alzheimer's disease contains both 3R and 4R forms. These differences are important in research because they may help explain why particular disorders affect different parts of the brain.

Under normal conditions, tau contributes to the stability of neuronal structures. Various biological processes regulate tau, including phosphorylation, which involves the addition of phosphate groups to proteins. Tau normally undergoes controlled changes in phosphorylation. In certain disease

conditions, abnormal phosphorylation and other chemical modifications can alter the protein's behavior. Abnormal tau may detach from microtubules, accumulate within cells, and form structures known as aggregates.

Alzheimer's disease is another major neurological condition in which abnormal tau is present. In Alzheimer's disease, tau accumulates inside neurons and can form neurofibrillary tangles. The distribution of tau pathology is associated with cognitive decline as the disease progresses. Alzheimer's disease also involves amyloid-beta abnormalities, meaning that it is not classified as a pure tauopathy in the same sense as certain other disorders. Nevertheless, tau remains an important subject of research into disease mechanisms and treatment.

Diagnosis of tau-related disorders can be challenging because symptoms may overlap among several neurological diseases. Doctors usually consider the person's medical history, neurological examination, cognitive assessment, and disease progression. Magnetic resonance imaging may show patterns of brain volume loss that support a particular diagnosis. Other investigations can help exclude conditions with similar symptoms. In specialized centers, biomarker testing using cerebrospinal fluid or advanced imaging may provide additional information about abnormal proteins.

Research into tauopathies is expanding across several areas. Scientists are studying how tau changes inside neurons, how abnormal tau spreads between cells, why certain brain regions are more vulnerable, and how different tau structures relate to specific diseases. Researchers are also investigating biomarkers that may help identify disease before severe symptoms develop and treatments designed to reduce abnormal tau production, modification, aggregation, or spread.

Clinical studies are evaluating several approaches directed toward tau. These include antibody-based therapies, vaccines, antisense medicines, and other methods intended to influence tau biology. Results have varied across studies, and no single approach has yet become a universal treatment for all tau-related

Correspondence to: Renjiro Tsukishima, Department of Neurological Sciences, Hoshizora Center for Brain Research, Yokohama, Japan, E-mail: renjiro.t.1865733@hoshizora.jp

Received: 12-May-2026, Manuscript No. BDT-26-32579; **Editor assigned:** 14-May-2026, PreQC No. BDT-26-32579; **Reviewed:** 27-May-2026, QC No. BDT-26-32579; **Revised:** 03-Jun-2026, Manuscript No. BDT-26-32579; **Published:** 10-Jun-2026, DOI: 10.35248/2168-975X.26.15.344

Citation: Tsukishima R (2026). Tauopathy: Understanding Tau-Related Brain Disorders and Their Clinical Effects. Brain Disord Ther. 15:344.

Copyright: © 2026 Tsukishima R. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

disorders. Continued research is needed to determine which strategies are effective for particular diseases and stages.

Tauopathies represent a diverse group of neurological conditions connected by abnormal tau biology. Their effects can include memory loss, behavioral changes, language difficulties, impaired movement, balance problems, swallowing difficulties, and progressive loss of independence. Accurate assessment can help distinguish these disorders from other neurological and psychiatric conditions. Although current treatment is mainly focused on symptom control and supportive care for many primary tauopathies, ongoing research into tau structure,

biomarkers, disease mechanisms, and targeted therapies may improve future options.

For individuals and families affected by tau-related disease, regular neurological care, rehabilitation, safety planning, nutritional support, emotional assistance, and practical family support remain important. Understanding the specific pattern of symptoms can help healthcare teams provide appropriate care as the condition changes. While tauopathies can be serious and progressive, comprehensive support can help preserve dignity, comfort, communication, and functional abilities for as long as possible.